UMASS/AMHERST \$312066005806256



DATE	DUE	

UNIVERSITY OF MASSACHUSETTS LIBRARY

5 73 374 no.601-780 1939-53

4		





MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 601

January, 1939

Meteorological Observations

FOR

JANUARY 1939

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments 67 ft. Time used, 75th Mer.

		Ter	nperatu	re,			9	W	ind.		Pr	ecipitat	ion.	
	Max	mum	Mini	mum.			urs of unschi	g. E	nt	s/hr*				
Day.	Degrees	Time	Degrees	Time.	Mean	Mean Relative Humidity	No. Hours of Bright Sunshine	Prevailing Direction	Total Movement	Max miles/hr*	Began	Ended	Rain	Snow.
$egin{array}{cccccccccccccccccccccccccccccccccccc$	- 43	2 A A P 2 P 2 P 3 P 4 P 3 P 2 P 2 P 3 P 4 P 4		8 P 12 P 8 A A A A A A A A A A A A A A A A A A	28.5 36.5 21.5 22.0 24.0 38.5 35.5 43.5 43.5 43.5 27.0 20.0 21.5 24.5 20.0 17.5 18.5 19.5 9.0 19.5 36.6	57 56 75 89 88 97 81 55 63 72 60 57 62 62 63 45 84 63 45 84 63 73 84 84 85 86 87 87 88 88 88 88 88 88 88 88 88 88 88	6.8 6.0 3.8 0.0 4.5 7.5 6.5 4.8 9.4 6.5 7.8 6.5 7.8 6.5 7.8 6.5 7.8 6.5 9.7 4.0 9.7 4.0 9.7 9.7 9.7 9.7 9.7 9.7 9.7 9.7 9.7 9.7	w w nw nw s nw		$\begin{array}{cccccccccccccccccccccccccccccccccccc$	11-30 P 0 A 0 A 2 A 2-30 P 6 A	12 P 12 M 1 8 A 11 A 4 A 4-30 P 8 P	.01	.23 1.00 .2

^{*}Based on least time required to blow one mile.

MASSO MASS.

NOTE-Beginning Mayor and relative hamility is based on readings taken at 8 A. M., cr x pox or 18 P. A. Prior to that time readings were taken at 8 A. M. and 8 P. M.

F. W. WHITTE FORE, Observer

January 1959	Normal	Extremes
Barometer, in inches	1	
(Readings reduced to sea level)		
Maximum 30.52, 4th	30.70	31.00, 1927
Minimum 29.20, 22nd	29.20	28.55, 1913
Mean semi-daily 30.041	30,07	
Range 1.32	1.51	2.18, 1913; 0.97, 1896
Air Temperature, in degrees F.	1.01	2.10, 1010, 0.01, 111
Highest	51.1	66.0, 1932
Lowest4, 27th	-7.1	-26.0, 1904
Mean	24.2	34.2, 1913; 13.9, 1918
Monthly range 56	58.2	54.2, 1515, 15.5, 1516
	+2O+m	
Highest mean daily	1	
Lowest mean daily 8.0, 26th		
Mean maximum 33.0	-	
Mean minimum 16.7		
Greatest daily range 32, 22nd		
Least daily range 7, 11th, 31st	1 1	
Precipitation, in inches		
Precipitation 2.21	3.61	7.15, 1898; 1.07, 1896
Snow 4.75	13.24	33,1898, 1923; 1.75, 11
Maximum precipitation in		
24 hrs 1.34, 5th, 6th	1	
Number of days with .01 or more 8	1 1	
Wind, in miles		
Total movement 5079	5055	7770, 1908; 2896, 1895
Greatest daily movement 389, 23rd	-	
Least daily movement 25, 16th	1	
Mean hourly velocity 6.8	6.8	
Maximum velocity 44, 22nd	32.7	47, 1938
Wind, direction	.,,	11, 11
Prevailing direction NW	WNW	
North, days 10	77.74.74	
Northeast, days 0	1	
East, days 0		
Southeast, days 0		
South, days 7		
Southwest, days 0		
West, days 4		
Northwest, days 10		
Weather		
Mean relative humidity, percent 69.9	70.1	
Mean cloudiness, percent 57	55	78, 1932; 37, 1908
Number of clear days 17	()	16, 1932; 2, 1914
Number of fair days 10	()	18, 1890, 1926; 3, 1946
Number of cloudy days 4	13	22, 1923, 1931; 4, 1920
	137	214, 1920; 74, 1932
Number hours bright sunshine (5)		
Number hours bright sunshine189 Percent of possible hours of	,	, , , , , ,
Percent of possible hours of bright sunshine64.3	46.7	

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1938.

The weather during January was pleasant, with less rain and snow than usual and with more sunshine than normal. The mean temperature for the month was 24.9 degrees which is .7 degrees higher than normal. The lowest was —1 degrees on the 27th. The average low temperature for January is —7.1 degrees, and the extreme was —26 degrees in 1904. The domestic heating load for the month was 1245 degree-days, the normal for January being 1265 degree-days. The total heating load for the season is 3541 degree-days, the normal being 3814 for this period. Only 4.75 inches of snow fell in January, while the normal is 13.24 inches. The total precipitation for the month was 2.21 inches compared to a normal of 3.61 inches. There were 189 hours of bright sunshine, with 17 days classified as "clear". The normal hours of sunshine for January is 137 hours. A wind velocity of 44 miles per hour occurred on the 22nd and there was thunder and lightning on that afternoon.

NOTF: The weather observatory completed 50 years of records with the December 1938 bulletin. A new set of normals has been calculated based on records from 1889 to 1938 inclusive.

Two errors were made in the December bulletin. The relative humidity on December 1 should read 52 percent making the mean monthly humidity 73.3 percent, and the maximum barometer for December was 30.67 inches on the 16th.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 602

February 1939

Meteorological Observations

FOR

FEBRUARY 1939

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments 67 ft. Time used, 75th Mer.

		Ten	pera	ture					Wind			Preci pit atio	on
	Maxim	num	Min	imum		>	rs of unshine	S C	ı t	les/br*			
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	No. Hours of Bright Sunshine	Prevailing Direction	Total Movement	Max. miles/hr*	Began	Ended	Rain
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 223 224 225 226 27 28	24 1 34 32 31 29 34 41 40 32 35 32 50 38 56 30 31 1 46 38 62 35 32 28 28 39 35 34 42	4 P P P P P P P P P P P P P P P P P P P	2 14 24 16 11 9 25 15 15 17 10 28 27 29 4 4 0 26 30 34 30 17 11 15 16 11 17 17 19 25 16 16 17 17 17 18 18 18 18 18 18 18 18 18 18 18 18 18	3 A A O A A 12 P A A A A A A A A A A A A A A A A A A	18.00 19.00 29.00 21.00 19.00 29.5 28.00 25.00 25.00 25.00 26.00 32.5 42.5 17.00 332.5 42.5 17.00 34.00 48.00 32.5 25.5 32.00 33.00 34.00 35.00 36.00 36.00 37.00	77 94 97 61 65 80 89 80 48 95 64 46 91 92 43 73 73 74 49 68 58 68 68 68 68 68 68 68 68 68 68 68 68 68	10 0 0 0 5 3 3 3 3 9 9 0 0 10 11 6 3 5 .8 8 .4 1 10 .0 0 7 .7 7 10 3 9 .8 6 .5 0 .0 0 10 .6 6 3 .2 2 8 0 .0 0 7 .5 9 .0 0 3 .8 8 3 .3 3 9 5 0 .0 0 9 .3 0 .0 0	nw nw nw nw s w n s nw n nw s s s w n nw s nw n nw s s nw n m w n w n n m w n n m w n m w n m w n m w n m m m m	82 53 55 282 83 51 105 108 129 202 162 50 144 433 147 725 111 120 173 257 141 204 115 101 144 60	11 9 15 48 9 6 9 15 18 9 39 18 20 25 8 20 11 32 48 20 25 8 20 11 32 48 21 32 48 33 48 48 48 48 48 48 48 48 48 48	0 A 0 A 0 A 0 A 10 P 0 A 0 A 0 A 12 M	3 A 12 P 12 P 4 A 12 P 1 P 1 P 8 A 9 P 9 P 8 8:30 A	.02 T T T .02 .50 .16 1.50 .28 3.00 .41 .10 .45 .27 .03 .17 1.00

^{*}Based on least time required to blow one mile.

February 1939	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum30.68, 16th, 17th	30.63	31.05, 1920
Minimum	29.24	28.56, 1895
Mean semi-daily	30.04	20.30, 1073
Range	1.38	1.89, '00, '08; .88, '13,'31
Air Temperature, in degrees F.	1.50	1.00, 00, 00, .00, 10, 01
Highest	50.5	65.0, 1930
Lowest	-7.4	-22.5, 1918
Mean	23.7	22.3, 1710
Range	58	
Highest mean daily		
Lowest mean daily 15 . 5, 17th		
Mean maximum		
Mean minimum		
Greatest daily range		
Least daily range		
Precipitation, in inches		
Precipitation	3.19	8.12, 1900; 0.62, 1901
Snow	13.99	48.75, 1893; 0.50, 1937
Maximum precipitation in 24 hrs0.97, 26th	13.99	40.73, 1093, 0.30, 1937
Number of days with .01 or more13	10	15, '93, '20; 4, 1901
Wind, in miles	10	13, 93, 20, 4, 1901
Total movement	4776	6445, 1896; 3438, 1892
Greatest daily movement433, 16th		, , , , , , , , , , , , , , , , , , , ,
Least daily movement50, 14th		
Mean velocity	7.1	
Maximum velocity48, 4th, 16th	31.7	48, 1934, 1937
Wind, direction		,
Prevailing direction	WNW	
North, days5		
Northeast, days0		
East, days0		
Southeast, days1		
South, days		
Southwest, days		
West, days1		
Northwest, days		
Weather		
Mean relative humidity, percent 73 . 3	66.6	
Mean cloudiness, percent 59	50.9	66, '90, '27; 31, 1905
Number of clear days	10	17, '19, '24; 2, 1927
Number of fair days6	8	16, 1920; 2, 1936
Number of cloudy days7	10	16, 1894; 2, 1920
Number hours bright sunshine173	158	221, 1924; 110, '27, '38
Percent of possible hours of bright		, , , , , , , , , , , , , , , , , , , ,
sunshine	53.4	
Thunder and lightning		

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1938.

The weather during February was warmer than usual, with many rainy days. The walks and roads were either slushy or icy throughout most of the month. The mean temperature for the month was 27.8 degrees, the normal being 23.7 degrees. On the 20th the temperature reached 62 degrees, a temperature exceeded only twice in February during the last 50 years. The lowest temperature was zero on the 17th. The heating load during the month was 1042 degree-days, compared to a normal of 1156 degree-days. The total heating load for the season up to February 28 is 4583 degree-days, while the normal for this period is 4970 degree-days. The total precipitation for the month was 3.62 inches, compared to a normal of 3.19 inches. Only 6 inches of snow fell, the average snowfall for February being 14 inches. The total snowfall for the winter up to the present time is 27.25 inches, compared to a normal of 38.35 inches for this period. The total wind movement during the month was 4097 miles, the normal being 4776. High winds reaching 48 miles per hour were recorded on the 4th and 16th. There were 173 hours of bright sunshine, the normal being 158 hours.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 603

March 1939

Meteorological Observations

FOR

MARCH 1939

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, 75th Mer.

		Γen	npera	ture					Wind			Precipitati	on	
	Maxi	mum	Min	imum		>	rs of unshine	× .	l ut	les/hr*				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	No. Hours of Bright Sunshine	Prevailing Direction	Total Movement	Max. miles/hr*	Ведап	Ended	Rain	Snow
1 2 3 4	44 32 44 48	9 A 4 P 4 P 1 P	33 22 16 31	12 P 12 P 7 A 0 A	38.5 27.0 30.0 39.5		3.5 11.2 11.2 5.8	nw nw s	139 291 116 211	40 28 20 20	1 A	2 а	.03	
4 5 6 7 8	44 55 36 30	1 P 4 P 0 A 2 P	36 36 20 12	7 A 12 P 12 P 7 A	40.0 45.5 28.0 21.0	96 82 48 35	0.0 1.0 9.8 11.5	w s nw nw	52. 170 478 307	12 25 48 22	4 A	9-30 a	T .49	
9 10 11 12 13	25 33 24 32 35	2 P 1 P 3 P 12 M 2 P	20 11 4 22 25	4 A 12 P 7 A 0 A 8 A	22.0 14.0 27.0	70 87	0.0 11.6 8.5 1.2 6.5	n n nw nw n	89 198 119 154 378	10 32 22 17 39	7 A 4 P 0 A 0 A	8 P 12 P 12 P 8 A	.60 .16 .73	2.0 8.0 2.0
13 14 15 16 17 18	39 37 39	2 P 3 P 2 P 2 P 12 M 1 P	26 17 25 19	6 A 6 A 12 P 7 A 7 A		57 79 83 60	11.8 8.2 4.5 11.8 10.0	n n nw nw	261 66 125 136 126	25 32 25 20 20	4 P 0 A	12 P 5 A	.83	7.0 T
19 20 21 22	29 40 33 35	2 P 5 P 4 P 4 P	12 18 22 16	6 A 2 A 7 A 6 A	20.5 29 0 27.5 25.5	49 88 58 57	12.0 1.0 12.1 12.1	n s nw nw	93 135 280 137	14 32 32 32 22				T
23 24 25 26 27	42 59 52 47 45	3 P 3 P 4 P 4 P 12 M	13 28 31 35 34	6 A 6 A 7 A 4 A 12 P		62 68 77 96 75	9.5 11.2 8.0 3.2 0.0	sw n n se w	58 56 107 93 72	8 11 22 8 13	1:30 A 8 A	8 A 9 A	.13	
28 29 30 31	36 43 35 45	4 P 4 P 4 P 2 P	27 22 31 34	11 P 6 A 5 A 0 A	31.5 32.5 33.0 39.5	93 42 97 71	5.7 12.5 0.0 1.6	n nw se w	175 109 126 137	13 13 9 12	1 A 3.30 A	11 A 6 P	.53	1.5 0.5

^{*}Based on least time required to blow one mile.

March 1939	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum30.56, 8th, 19th	30.57	30.96, 1904
Minimum	29.24	28.47, 1914
Mean semi-daily	30.00	
Range1.16		2.10, 1014; 0.85, 1915
Air Temperature, in degrees F.		2.10, 1 11, 0.00, 1210
Highest	63.8	79.5, 1907, 1921
Lowest	6.1	-7.5, 1906
Mean	34 4	
Range	57.6	1
Highest mean daily		
Lowest mean daily11.0, 11th		
Mean maximum38.9		
Mean minimum		
Greatest daily range		
Least daily range		
Precipitation, in inches		
Precipitation4.19	3.70	7.13, 1899; 0.12, 1915
Snow	7.47	27, 1899; 0, 1921
Max precipitation in 24 hrs. 1.02, 15th 16th		
Number of days with 01 or more12	11	17, 1890, 1913, 1936; 3, 1915
Wind, in miles		
Total movement	5944	8182, 1896; 3006, 1905
Greatest daily movement478, 7th		
Least daily movement52, 5th		
Mean velocity	7 7	
Maximum velocity	31.2	48, 1932
Wind. direction		, ,
Prevailing directionNW	WNW	
North, days10		
Northeast, days0		
East, days		
Southeast, days		
South, days4		
Southwest, days		
West, days		
Northwest, days11		
Weather		
Mean relative humidity, percent 69 1	64.2	i
Mean cloudiness, percent48	51.8	
Number of clear days	11	22, 1924; 3, 1901
Number of fair days	10	17, 1900; 4, 1895, 1897, 193b
Number of cloudy days9	10	21, 1901; 1, 1915
Number hours bright sunshine217	199	292, 1924; 93, 1901
Percent of possible hours of bright		
sunshine	53.6	
Thunder and lightning		

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1938

The weather during March was cold and wintry with snow on the ground throughout most of the month. The mean temperature for the month was 30.9 degrees, the normal for March being 34.4 degrees. It is interesting to note that the mean temperature for February of this year was 27.8 degrees while the mean temperature during March is normally 10.7 degrees higher than in February. The domestic heating load for March was 1057 degree-days while the normal for March is 949 degree-days. The total heating load for the winter to the end of March is 5640 degree-days, the normal for this period being 5919 degree-days.

A total of 20 inches of snow fell during the month. The storm on the 11th, 12th, and 13th was generally recognized as the heaviest snowfall in March since the great storm of 1888. Twelve inches of snow fell during the days mentioned and this was preceded with 6 inches on the 9th. The total precipitation during the month was 4.49 inches, the normal being 3.70 inches. There were 217 hours of bright sunshine which is well above the average of 199 hours. A high wind velocity of 48 miles per hour was recorded on the 7th.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 604

April 1939

Meteorological Observations

FOR

APRIL

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, 75th Mer.

-		Тетр	erati	ıre					Wind		P	recipitatio	on	
Day	Maxim		Degrees	Time	Mean	Mean Relative Humidity	No. Hours of Bright Sunshine	Prevailing Direction	Total Movement	Max. miles hr*	Ведап	Ended	Rain	Snow
1 2 3 4 5 6 7	43 41 42 49 49 1	0 A 3 4 P 4 P 4 P 4 P 4 P 4 P 4 P 4 P 4 P 4	33 33 31 29 26 27 28	8 A 12 P 4 A 6 A 4 A 12 P	44.5 38.0 36.0 35.5 37.5 38.0 36.0	84 64 62 55 79	9.8 1.0 3.0 9.4 12.7 2.5 8.5		101 292 273 243 168 106 318	13 28 22 22 14 20 28	3 A	11 A	.90	2.00 T
8 9 10 11 12 13 14	50 47 48 57	3 P 5 P 6 P 4 P 4 P 3 P 5	24 30 26 33 28 25 33	6 A		67 92 98 72 47 54	2.0 5.3 0.0 0.1 3.5 9.8 10.0	se w nw nw w nw sw	86 195 71 66 236 329 232	14 32 10 15 32 30 39	8 A 7 A 0 A	8 P 12 P 2 A	.08 .59 .19	Т
15 16 17 18 19 20 21	55 42 49 57 65	4 P 2 P 2 P 6 P 3 P 3 P 3 P	35 27 28 35 39 38 32	12 P 4 A 6 A 6 A 6 A 5 A 6 A	38.5 44.0 47.5 48.5	38 5 61 6 97 0 98 6 66 5 59	8.5 11.8 9.0 0.0 0.0 8.3 13.0	n nw nw s	263 123, 110 119 78 144 127	25 12 11 9 10 18 17	2:30 A 6 P 0 A 2 A	3 A 12 P 12 M 8 P	.03 .26 .40 .56	
22 23 24 25 26 27 28 29 30 31	65 59 64 82 53 51 49 61 59	5 P 4 P 5 P 4 P 3 P 6 P 2 P	45 43 37 41 43 42 37 32 31	7 A 12 P 4 A 5 A 7 A 12 P 12 P 6 A 6 A	51.0 50.5 61.5	0 62 5 63 5 65 0 94 5 92 0 83 5 51	8.0 6 5 7.5 12.5 0.0 0.0 1.8 13.8 10.7	e ne	187 167 143 56 105 62 129 113 69	22 25 22 17 13 11 12 17 13	6 A 6 A 7 A 0 A	7 A 12 P 8 P 6 A	T .09	

^{*}Based on least time required to blow one mile.

April 1939	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum	30.48	30.71, 1911
Minimum29.33, 2nd		28 99, 1907, 1929
Mean semi-daily	29.99	4 53 4030 52 4040
Range	1 11	1.52, 1930; .72, 1919
Air Temperature, in degrees F.	70.4	90 F 1904
Highest	79.4	88,5, 1896
Lowest	22.0 45.7	8.5, 1923 52, 1921; 41,2, 1926
Mean	57.4	32, 1921; 41,2, 1920
Range	374	
Lowest mean daily		
Mean maximum51.9		
Mean minimum		
Greatest daily range		
Least daily range		
Precipitation, in inches		
Precipitation	3 35	6.89, 1929; .76, 1892
Snow	2.17	11, 1891; 0, 1910, 1934
Max. precipitation in 24 hrs 1.23, 2nd	_,,,	, , . , , ,
Number of days with 01 or more13	11	18, 1909; 3, 1892
Wind, in miles		
Total movement	5404	8208, 1908; 3853, 1917
Greatest daily movement329, 13th		
Least daily movement56, 25th		
Mean hourly velocity6.5	7.5	
Maximum velocity	31.4	40, 1935, 1938
Wind. direction	****	
Prevailing direction	HXH.	
North, days 5		
Northeast, days2		
East, days		
Southeast, days		
South, days		
West, days4		
Northwest, days		
Weather		
Mean relative humidity, percent 68.8	61.6	
Mean cloudiness, percent69	51.8	75, 1901; 34, 1925, 1927
Number of clear days10	11	19, 1911; 3, 1898, 1902
Number of fair days 8	9	18, 1915; 2, 1894, 1901
Number of cloudy days	10	22, 1901; 3, 1899, 1924
Number hours bright sunshine189	220	296, 1911; 103, 1901
Percent of possible hours of bright		
sunshine	54 7	
Thunder and lightning26th		
Frosts after the 15th None		
Last snow		

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1938

The weather during April was wet, cold, and disagreeable. The total precipitation for the month was 4.56 inches, the normal being 3.35 inches. Two inches of snow fell during the month, all on April 2. The total precipitation since January 1 is now 14.88 inches. The normal for this period is 13.85 inches.

The mean temperature for the month was 42.5 degrees. This is the coldest April since 1926, when the mean was 41.2 degrees. The mean temperatures for April 1898, 1907, and 1917 were also lower than that of the past month. The domestic heating load was 675 degree-days during April, compared to a normal of 579 degree-days. The total heating load for the season is 6315 degree-days; the normal for this period is 6498 degree-days. The total snowfall during the past winter was 49.25 inches, the normal being 47.73 inches. It is interesting to note that 23.5 inches fell during the two storms of November 24-25 and March 11-13.

There were 189 hours of bright sunshine, compared to a normal of 220 hours.

Dr. J. K. Shaw gives the following report on vegetation: "In contrast to the very early season of 1938, the present spring will be recorded as very late. Vegetation is about a month behind that of last year and nearly two weeks later than average. However, a few very warm days would enable vegetation to catch up; the late spring will have little if any effect on the maturity of most fruit crops. A late spring is on the whole favorable, for it shortens the probable period of danger from spring frosts.

"Prospects for a peach crop are good as there was no cold severe enough to kill the fruit buds. We have observed considerable injury to apple wood, probably caused by the unseasonable cold of the last week in November before the wood had completely matured. The temperature of -4° on November 26, 1938 is $7\frac{1}{2}$ degrees lower than any previously recorded for this month at this station, and no lower temperature was recorded during the winter. It will not be surprising if this most unusual cold caused some injury to fruit plants."

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 605

May 1939

Meteorological Observations

FOR

MAY

1939

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, 75th Mer.

		Ten	iperat	ture					Wind			Precipitat	ion	
	Maxi	imum	Mini	imum			s of nshine	50	ıt	es, hr*				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	No. Hours of Bright Sunshine	Prevailing Direction	Total Movement	Max. miles, hr*	Ведап	Ended	Rain	Snow
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 22 1 22 23 24	61 59 68 70 82 89 75 67 71 63 55 66 67 73 63 80 79 74 57	12 M 1 P 1 P 4 P 4 P 4 P 5 P 5 P 4 P 4 P 4 P 4 P 4 P 4 P 4 P 4 P 5	34 36 38 43 51 55 45 57 55 45 57 32 36 32 36 32 36 37 37 36 37 37 36 37 37 37 37 37 37 37 37 37 37 37 37 37	6 A A 6 A A 6 A A 6 A A 6 A A 6 A A 12 P 6 A A 6 A A 5 A A 6 A A A 6 A A A 6 A A A 6 A A A 6 A A A 6 A A A 6 A A A 6 A A A 6 A A	46.5 48.5 48.5 55.5 50.5 72.0 60.0 66.5 72.0 60.0 52.5 49.0 49.0 51.5 54.5 55.5 54.0 55.5 54.0 66.5 54.0	52 51 52 48 57 59 50 73 96 46	7.88.0 3.55 6.03 14.1 14.1 7.00 14.4 9.3 0.5 5.5 14.0 11.5 10.5 6.0 0.0 0.0 10.0		118 922 1977 2200 168 67 180 1178 154 212 195 98 65 172 98 88, 172 118 132 88, 144 126 146 146 147 148 148 148 148 148 148 148 148 148 148	14 8 20 22 25 9 28 14 15 18 21 17 11 10 18 15 10 20 13 13 9 6 9 10 10 10 10 10 10 10 10 10 10	10:30 A 6 A 6:30 P	8:30 P	Т	
25 26 27 28 29 30 31	81 78 81 89 82 82 89	6 P 3 P 1 P 3 P 3 P 4 P	49 56 55 65 60 55 64	4 A 5 A 6 A 1 A 12 P 5 A 4 A	65.0 67.0 68.0 77.0 71.0 68.5 76.5	72	8.0 14.6 9.6 9.4 14.4 14.9 14.8	s se se nw w	83 81 149 183 159 135 93	8 10 18 39 25 18 12	2 P	11 Р	1.40 T	

^{*}Based on least time required to blow one mile.

May 1939	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum30.30, 13th	30.39	30.62, 1936
Minimum29.43, 9th	29.51	29.10, 1938
Mean semi-daily	29.96	
Range	.88	
Air Temperature, in degrees F.	İ	
Highest	86.1	94.5, 1896, 1911
Lowest	31.2	24.0, 1900
Mean59.7	57.1	
Range	54.9	
Highest mean daily		
Lowest mean daily46, 13th		
Mean maximum		
Mean minimum47.2		
Greatest daily range		
Least daily range		
Precipitation, in inches		
Precipitation	3.60	7.44, 1931; .48, 1903
Snow	0,00	111, 1101, 110, 1700
Max, precipitation in 24 hrs 1.40, 28th		
Number of days with .01 or more 4	12	20, 1901; 5, 1903
Wind, in miles	12	20, 1701, 3, 1703
Total movement	4504	5946, 1907; 2180, 1894
Greatest daily movement	1501	0 10, 1207, 2100, 1071
Least daily movement		
Mean hourly velocity5.4	6.1	
Maximum velocity	28.8	45, 1935
Wind. direction	20.0	43, 1733
Prevailing direction	11.	
North, days	**	
North past days		
Northeast, days		
East, days0		
Southeast, days		
South, days		
Southwest, days4		
West, days		
Northwest, days		
Weather	60.7	
Mean relative humidity, percent 60 5		70 1001, 20 1032
Mean cloudiness, percent	52.1 11	70, 1901; 30, 1923
Number of clear days	11	20, 1923; 0, 1927
Number of fair days	9	17, 1907, 1926; 5, 1923
Number of cloudy days		20, 1927; 3, 1903, 1922 334, 1922; 137, 1927
Number hours bright sunshine284	252	334, 1924; 137, 1927
Percent of possible hours of bright	E E 6	
sunshine	55.6	
Thunder and lightning 9th, 21st, 28th	M 12	I 9 1022. Apr. 32 1004
Last frost	may 12	June 8,1932; Apr. 23, 1904
h		

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1938.

The weather during May was warm and dry. The mean temperature for the month was 59.7 degrees compared to a normal of 57.1 degrees. This is the highest mean temperature for May since 1922 with the exception of 1933 when it was 60.5 degrees. The highest temperature was 89 degrees which occurred on three days; the 7th, 28th, and 31st. The lowest temperature during the month was 32 degrees on the 14th and 16th. The total heating load for the month of May was 166 degree-days, making the heating load for the season 6481 degree-days, the normal being 6743 degree-days for this period.

The total rainfall for the month was 2.15 inches; the normal, 3.60 inches. In 1931, 7.44 inches fell in May and in 1903, .48 inch. There was a heavy thunderstorm with high wind on the 28th, 1.40 inches being recorded. The total precipitation since the first of the year is now 17.03 inches, compared to a normal of 17.45 inches for this period. Four days of rain were recorded during the month, which is the least number of rainy days on record for May since 1889. Only one day of rain, the 9th, was recorded from April 29 to May 20.

There were 284 hours of bright sunshine compared to a normal of 252 hours for May. The mean humidity for the month was nearly normal. The last frost occurred on May 16, the normal date being May 12.

Dr. J. K. Shaw of the Pomology Department makes the following report on vegetation: "Vegetation made rapid progress during May and apple trees were in full bloom on about May 18-22, approximately one week later than normal. Slight frost did some injury to early blooming strawberries and tender plants, but the amount of injury was insignificant. Some winter injury has been noted on nursery-grown apple trees of tender varieties, probably from the unusual cold of late November. Raspberry canes were badly killed, probably from the same cause. This will, in many plantations, severely reduce the crop.

"The weather during the blooming period of tree fruits was favorable. McIntosh apple trees bloomed heavily and should have a good crop. Other varieties were variable.

"Most of the month was rather dry but heavy showers toward the end of the month renewed the water supply and trees will have water enough to meet their needs for some time."

GOODELL LIBRARY MASS.STATE COLLEGE AMHERST.MASS. MASSACHUSETTS

10 37

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 606

June 1939

Meteorological Observations

FOR

JUNE 1939

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, 75th Mer.

		Гen	perat	ture					Wind			Precipitation	o n	
	Maxi	mum	Min	ımum			s of inshine	20 _	at	es. hr*				
Lay	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	No. Hours of Bright Sunshine	Prevailing Direction	Total Movement	Max. miles hr*	Began	Ended	Rain	Snow
1 2 3	77 77	3 P 5 P	56 49	5 A 5 A	63.0	50	8.0 12.5	W.	112 92	8 13	1 A	2 A	.08	
	77 76 81 68	3 P 5 P 4 P 7 A	46 51 48 54	5 A 3 A 6 A 12 P	64.5	80 49	13.8 6.1 12.2 10.0	w w n	106 130 127 255	12 17 20 32	7:30 A	10:30 A	.17	
	79 83 88	5 P 4 P 5 P	42 52 65	6 A 5 A 6 A	60.5 67.5 76.5	56 82 76	13.3 4.5 11.4	W	86 82 82	9 11 20	7 Р	10 р	.43 T	
	85 90 82 80	3 P 5 P 1 P 3 P	62 67 63 56	5 A 5 A 12 P 5 A	78.5 72.5	71 59	6.7 15.0 14.3 10.6	SW S	58 117 156 72	10 13 14 8	10 р	12 P	T T	
	74 77 76	3 P 2 P 4 P 5 P	58 49 61	12 P 4 A 4 A	66.0 63.0	76 57	$ \begin{array}{c c} 10.0 \\ 2.5 \\ 10.0 \\ 14.1 \end{array} $	SW DW SW	158 118 261	17 13 20	0 A	4:30 P	.63 T	
	58 73 75	4 P 4 P 3 P	51 46 47	12 P 4 A 5 A	54.5 59.5 61.0	91 63 67	0.0 9.0 9.5	nw	101 57 119	48 9 13	2 A	8 A	.80	
	79 84 81	3 P 3 P 3 P	62 62 55	8 A 6 A 5 A	73.0 68.0	54 52	0.6 15.3 11.7	s nw s	71 154 86	6 15 7	7 A	9:30 A		
	73 79	6 P 3 P	57 50	4 A 5 A			0.5	se	102	17 14	2:30 A 11:30 P	4:30 P 12 P	.20	
	70	3 P 5 P	57	5 A 6 A			4.3	s n	103	17	0 A	8 P		
	70 80 79	3 P 5 P 4 P	59 57 58	5 A 4 A 5 A	64.5 68.5 68.5	82 08 80	1.5 12.5 6.8	n w s	83 45 79	10 7 8	6 A	8 A	.01 T	
)	83 78	2 P 4 P	64 67	5 A			0.8	S	149 211	14 11	1:30 A	11 P		

^{*}Based on least time required to blow one mile.

June 1939	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum	30.30	30.54, 188
Minimum		29.24, 190.
Mean semi-daily		23.21, 130.
Range	0.75	
Air Temperature, in degrees F.	0.70	
Highest	91.2	101, 1919
Lowest	40.0	34, 189
Mean66.7	65 7	01,10
Range		
Highest mean daily78.5, 11th		
Lowest mean daily		
Mean maximum		
Mean minimum55.7		
Greatest gaily range37, 7th	1	
Least daily range		
Precipitation, in inches		
Precipitation	3 51	9.68, 1922; 0.76, 1908
Snow	1 331	7.00, 1722, 0.70, 1900
Max precipitation in 24 hrs80, 17th		
		17 1022.1 1009
Number of days with 01 or more12 Wind, in miles	1 1	17, 1922; 4, 1908
	3585	4571 10 8, 1100 1007
Total movement		4571, 19. 8; 1409, 1906
Greatest daily movement		
Least daily movement45. 27th		
Mean hour's velocity4 8		10 103
Maximum velocity48, 17th	24.6	40, 1931
Wind direction	11.011.	
Prevailing direction	WSW	
North, days		
Northeast, days0		
East, days0	i i	
Southeast, days		
South, days9		
Southwest, days5		
West, days7		
Northwest, days4		
Weather		
Mean relative humidity, percent 69 1	66.9	
Mean cloudiness, percent60		71, 1903; 28, 1908
Number of clear days	10	22, 1908; 1, 1927
Number of fair days11	12	23, 1912; 4, 1895, 1903
Number of cloudy days	8	22, 1903; 1, 1923
Number hours bright sunshine254	257	362, 1908; 102, 190,
Percent of possible hours of bright		
sunshine	54.0	
Thunder and lightning1, 11, 13, 17, 24, 25		
Last frost		June 8,1932; Apr. 23, 1904
		J ., _, _, _, _ , _ , ,

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1938.

The weather during June was very nearly normal in most respects. The month generally was considered dry, but apparently there were a number of local showers which gave more rain at the weather observatory than at nearby points. The total rainfall for the month was 3.21 inches which is .30 less than normal. This rain fell on 12 days. In addition, there were traces of rain with a fall of less than .01 inch on 6 additional days. The total rainfall since January 1 is now 20.24 inches; the normal for this period is 20.96 inches. The mean temperature for the month was 66.7 degrees which is one degree higher than normal. The highest temperature was 90 degrees on the 11th and the lowest, 42 degrees on the 7th. The domestic heating load for the month was 40 degree-days, compared to a normal of 69 degree-days. The total heating load for the season was 6521 degree-days and the normal, 6812 degree-days.

There was a heavy windstorm on the 17th, with considerable damage to trees. The maximum velocity recorded was 48 miles an hour. There were 254 hours of bright sunshine which is 3 hours less than normal.

Dr. J. K. Shaw reports that "all fruit plants made normal progress during June. Tree fruits have set fair to good crops. Strawberries produced fair to good crops but raspberries and blackberries will be light due to winter injury to the canes. Rainfall has been rather short but timely showers have thus far prevented real injury to fruit crops. Scab control on McIntosh apple trees has not been as difficult as last year because of fewer humid periods, and thorough and timely spraying has been effective in control of the disease."

GOODELL LIBRARY MASS.STATE COLLEGE AMHERST.MASS.

10 37

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 607

July 1939

Meteorological Observations

FOR

JULY

1939

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, 75th Mer.

		Ten	apera	ture					Wind	į	P	Precipitation			
	Max	imum	Min	imum			s of nshine	-0		ss/hr*					
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	No. Hours of Bright Sunshine	Prevailing Direction	Total Movement	Max. miles/hr*	Began	Ended	Rain	Snow	
123456789012345678901223456789001	81 74 81 89 90 83 93 93 90 86 80 87 77 74 80 81 83 83 81 85 90 92 92 84 86 88 88 88 88 88 88 88 88 88 88 88 88	3 P 4 P 7 5 P 2 P 2 P 2 P 2 P 4 P 7 5 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P	61 52 45 67 68 69 61 68 66 65 44 47 52 47 53 49 55 47 53 49 56 68 69 67 68 67 68 68 68 68 68 68 68 68 68 68 68 68 68	12 P P S A A A A A A A A A A A A A A A A A	71.0 63.0 63.0 70.0 78.5 77.5 81.0 79.0 79.0 66.5 60.0 63.0 63.0 67.0 67.0 67.0 67.0 72.5 80.0 77.5 75.0 77.5 75.0	58 60 63 68 75 72 71 66 60 60 54 72 59 62 59 52 55 60 64 58 63 66 73 86 73 86 73 75 75 75 75 75 75 75 75 75 75 75 75 75	15.0 13.5 15.2 15.2 15.2 15.9 15.8 9.0 11.7 11.7 11.7 13.3 14.8 13.7 11.0 13.0 1	nw nw se se se s w nw n sw s mw nw nw sw se nw nw nw sw se nw nw sw se nw nw sw nw nw nw sw nw nw sw nw nw sw sw nw nw sw sw nw nw sw sw nw nw sw nw nw sw nw nw nw sw nw nw nw nw nw sw nw	170 179 71 66 122 151 108 124 62 123 141 101 123 160 87 730 48 98 68 57, 78 59 49 167 74 102 77 69 83	13 20 10 9 12 17 9 22 5 18 17 25 11 13 18 6 8 13 5 11 11 8 15 9 7 13 13 13 14 15 15 16 16 17 18 18 18 18 18 18 18 18 18 18 18 18 18	2:30 P 3.30 A 2:30 P 0 A 3 P 3 P 11:30 A	3 P 2 P 2 P 2 A 8 P 6 P 1:30 P	.18		

^{*}Based on least time required to blow one mile.

Baromeler, in inches (Readings reduced to sea level) Maximum .30.28, 21st, 22nd Minimum .29.58, 14th Mean semi-daily .29.97 Range .0.70 Air Temperature, in degrees F. Highest .93, 7th, 8th Lowest .45, 3rd Mean .71.2 Range .48 Highest mean daily .81, 7th Lowest mean daily .60, 13th, 1 th Mean maximum .83 5 Mean minimum .58 8 Greatest daily range .38, 4th Least daily range .12, 31st Precipitation, in inches Precipitation 2.30 Snow Max. precipitation in 24 hrs. .76, 14th Number of days with 01 or more .7 Wind, in miles	30.27 29.59 29.96 .68 93.9 46.4 70.8 47.7	30.50, 1892 29.27, 1933 .97, 1892; .47, 1938 104, 1911 40, 1890 ,1898 14 51, 18)7; .70, 1929 20, 1915; 4, 1929
Maximum 30.28, 21st, 22nd Minimum 29.58, 14th Mean semi-daily 29.971 Range 0.70 Air Temperature, in degrees F. Highest 93, 7th, 8th Lowest 45, 3rd Mean 71.2 Range 48 Highest mean daily 81, 7th Lowest mean daily 60, 13th, 1 th Mean maximum 83 5 Mean minimum 58 8 Greatest daily range 12, 31st Precipitation, in inches Precipitation, in inches Precipitation 2.30 Snow Max. precipitation in 24 hrs. .76, 14th Number of days with 01 or more .7 Wind, in miles	29.59 29.96 .68 93.9 46.4 70.8 47.7	29.27, 19397, 1892; .47, 193. 104, 191 40, 1890 ,1893
Minimum. 29.58, 14th Mean semi-daily 29.971 Range. 0.70 Air Temperature, in degrees F. Highest 93, 7th, 8th Lowest 45, 3rd Mean 71.2 Range 48 Highest mean daily 81, 7th Lowest mean daily 60, 13th, 1 th Mean maximum 83 5 Mean minimum 58 8 Greatest daily range 12, 31st Precipitation, in inches Precipitation 2.30 Snow 36, 14th Number of days with 01 or more 7 Wind, in miles 7	29.59 29.96 .68 93.9 46.4 70.8 47.7	29.27, 193 .97, 1892; .47, 193 104, 191 40, 1890 ,189
Mean semi-daily 29.971 Range 0.70 Air Temperature, in degrees F. Highest Highest 93, 7th, 8th Lowest 45, 3rd Mean 71.2 Range 48 Highest mean daily 60, 13th, 1 th Mean maximum 83.5 Mean minimum 58.8 Greatest daily range 12, 31st Precipitation, in inches Precipitation 2.30 Snow 38, 4th Mumber of days with 01 or more 76, 14th Number of days with 01 or more 7 Wind, in miles	29.96 .68 93.9 46.4 70.8 47.7	.97, 1892; .47, 193 104, 191 40, 1890 ,189 14 51, 18)7; .70, 192
Range	.68 93.9 46.4 70.8 47.7	104, 191 40, 1890 ,189 14 51, 18)7; .70, 192
Range	93.9 46.4 70.8 47.7	104, 191 40, 1890 ,189 14 51, 18)7; .70, 192
Highest 93, 7th, 8th Lowest 45, 3rd Mean 71.2 Range 48 Highest mean daily 81, 7th Lowest mean daily 60, 13th, 1 th Mean maximum 83 5 Mean minimum 58 8 Greatest daily range 12, 31st Precipitation, in inches Precipitation 2.30 Snow 38, 4th Max. precipitation in 24 hrs 76, 14th Number of days with 01 or more 7 Wind, in miles	46.4 70.8 47.7	40, 1890 ,189 14 51, 18)7; .70, 192
Lowest 45, 3rd Mean 71, 2 Range 48 Highest mean daily 81, 7th Lowest mean daily 60, 13th, 1 th Mean maximum 83 5 Mean minimum 58 8 Greatest daily range 38, 4th Least daily range 12, 31st Precipitation, in inches Precipitation 2.30 Snow Max. precipitation in 24 hrs. 76, 14th Number of days with 01 or more 7 Wind, in miles	46.4 70.8 47.7	40, 1890 ,189 14 51, 18)7; .70, 192
Mean 71.2 Range 48 Highest mean daily 81, 7th Lowest mean daily 60, 13th, 1 th Mean maximum 83 5 Mean minimum 58 8 Greatest daily range 38, 4th Least daily range 12, 31st Precipitation, in inches Precipitation 2.30 Snow Max. precipitation in 24 hrs. 76, 14th Number of days with 01 or more 7 Wind, in miles	70.8 47.7	14 51, 18)7; .70, 192
Range .48 Highest mean daily .81, 7th Lowest mean daily .60, 13th, 1 th Mean maximum .83 5 Mean minimum .58 8 Greatest daily range .12, 31st Precipitation, in inches Precipitation 2.30 Snow Max. precipitation in 24 hrs. .76, 14th Number of days with 01 or more .7 Wind, in miles	4 10	
Highest mean daily 81, 7th Lowest mean daily 60, 13th, 1 th Mean maximum 83 5 Mean minimum 58 8 Greatest daily range 38, 4th Least daily range 12, 31st Precipitation, in inches Precipitation 2.30 Snow Max. precipitation in 24 hrs. .76, 14th Number of days with 01 or more Wind, in miles	4 10	
Lowest mean daily 60, 13th, 1 th Mean maximum 83 5 Mean minimum 58 8 Greatest daily range 38, 4th Least daily range 12, 31st Precipitation, in inches Precipitation 2.30 Snow Max. precipitation in 24 hrs. .76, 14th Number of days with 01 or more Wind, in miles		
Mean maximum. 83 5 Mean minimum. 58 8 Greatest daily range. 38, 4th Least daily range. 12, 31st Precipitation, in inches Precipitation. 2.30 Snow.		
Mean minimum		
Greatest daily range 38, 4th Least daily range 12, 31st Precipitation, in inches 2.30 Precipitation 2.30 Snow Max. precipitation in 24 hrs .76, 14th Number of days with 01 or more Wind, in miles		
Least daily range		
Precipitation, in inches Precipitation		
Precipitation		
Snow		
Max. precipitation in 24 hrs	11	20, 1915; 4, 192
Number of days with 01 or more7 Wind, in miles	11	20, 1915; 4, 192
Wind, in miles	11	20, 1915; 4, 192
Total movement	3422	5,097, 1909; 1,109, 189
Greatest daily movement179, 2nd		
Least daily movement48, 18th		
Mean hourly velocity4.1	4.6	
Maximum velocity25, 12th	28.7	44, 193
Wind, direction	CIVI	
Prevailing direction	SW	
North, days4		
Northeast, days0		
East, days		
Southeast, days		
South, days5		
Southwest, days		
West, days		
Northwest, days8		
Weather	60.1	
Mean relative humidity, percent 65.1	68.4	(0.1011.21.10)
Mean cloudiness, percent	50.9	69, 1914; 31, 192
Number of clear days	10	22, 1923; 0, 191
Number of fair days14	14	24,'09,'21;5,'89,'23,'28,'3
Number of cloudy days4	268	18, 1889; 0, 191
Number hours bright sunshine283	200	371, 1910; 180, 193
Percent of possible hours of bright	58	
sunshine	30	
I nunder and lightning1, 8, 14, 24, 25, 31		

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1938.

The weather during July was very dry until the last few days of the month. Only .95 of an inch fell until July 27. This light rainfall in July with a deficiency of 1.45 inches in May and .30 in June made a dry growing season with considerable damage to crops. The total rainfall for the month was 2.30 inches compared to a normal of 4.10 inches. The least rainfall we have had in July was .70 inch in 1929. The total precipitation since January 1 is now 22.54 inches. The normal precipitation for this period is 25.06 inches.

The mean temperature for the month was 71.2 degrees, the normal being 70.8 degrees. The highest temperature during the month was 93 degrees which occurred on the 7th and 8th. The lowest was 45 degrees on the 3rd. The total wind movement during the month was 3072 miles, the normal being 3422 miles. The maximum velocity was 25 miles on the 12th. There were 283 hours of bright sunshine compared to a normal of 268 hours.

Dr. J. K. Shaw of the Pomology Department gives the following report: "The period of dry weather, which began in June, continued until July 27 with only one or two showers. During the last few days of July there were several showers, bringing sufficient water to relieve conditions for a week or two. The drouth did not become severe enough to do great damage to fruit crops though it must have interfered with the development of full size on early varieties ripening in late July and early August. Lawns and shallow-rooted crops suffered more or less severely. If timely rains come during August, there will be, on the whole, little drouth injury to fruit crops during this season. The dry weather has made weed control easier and has been unfavorable to the development of diseases and certain insects, and should favor the development of fruit buds for next season's crop."

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 608

August 1939

Meteorological Observations

FOR

AUGUST

1939

C. 1. GUNNESS

OBSERVATORY

Latitude, 42 23' 48.5" N. Longitude, 72 31' 10" W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, 75th Mer.

		Γen	perat	ture					Wind Precipitation					1	
	Maximum Minimum					s of nshine			s h*						
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	No. Hours of Bright Sunshine	Prevailing Direction	Total Movement	Max. miles	Began	Ended	Rain	Cusus	
	87 85 88 78 83 88	3 P 4 P 4 P 10 A 3 P 3 P	65 55 66 65 61 54	12 P 6 A 4 A 12 P 6 A 6 A	76.0 70.0 77.0 71.5 72.0 71.0	56 78 94 68 63	14.5 14.5 10.3 1.3 12.0 14.3	w se se sw nw sw	165 58 169 73 79 43	17 8 17 14 13 8	6 P 10:30 A	10:30 Р 4:30 Р	.23		
	87 86 88 83 87 86	4 P 2 P 1 P 3 P 3 P 2 P	63 69 62 58	5 A 6 A 2 A 5 A 5 A 5 A	75.0 74.5 78.5 72.5 72.5 72.0	70 82 63 69	10.1 10.5 5.0 14.2 14.1 11.1		64 110 146 124 51 108	8 10 12 17 8 10	3 Р	4 P	.04		
	85 85 89 87 88 88	4 P 3 P 3 P 1 P 4 P	69 63 60 65 64 60	12 P 12 P 6 A 6 A 4 A 5 A	77.0 74.0 74.5 76.0	91 75 62 73 73	3.9 14.0 14.0 10.6 10.2 10.0	s nw s se se	189 83 51 134 67 74	25 10 7 28 6 9	6 P 0 A	7 P 1 A	.43 .07 T		
	84 84 87 89 86	1 P 2 P 1 P 2 P 3 P	72 67 75 67 60 59	4 A 6 A 6 A 12 P 6 A	78.0 75.5 81.0 78.0 73.0 73.5	73 90 86 73 64	8.1 2.8 2.7 10.7 12.0 10.5	nw s s sw sw	124 178 112 103 50 72	10 18 11 10 5	2 A 1:30 P	4 P 2:30 P	.87 .09		
	88 80 82 82 82	3 P 12 M 2 P 3 P 2 P	59 66 64 58 50	6 A 5 A 12 P 6 A 5 A	73.0 73.0 73.0 70.0	85 67 71	2.4 8.9 9.9 10.0		74 122 122 111	5 14 10 11	12 м	5 Р	.16		
	73 76 70	11 A 4 P 3 P	56 60 63	6 A 5 A 7 A	64.5 68.0 66.5	69 67	10.2 3.4 0.8	ne n n	118 192 169	13 20 14			T		

^{*}Based on least time required to blow one mile.

August 1939	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum30.30, 12th	30.32	30.50, 193-
Minimum	29,61	28.87, 1936
Mean semi-daily	30.00	
Range	.71	
Air Temperature, in degrees F.		
Highest	91.6	100, 191
Lowest	43.4	37, 1894, 190
Mean	68.6	0., 10, 1, 1, 1,
Range	48.1	
Highest mean daily	-0	
Lowest mean daily		
Mean maximum84 2		
Mean minimum		
Greatest daily range		
Least daily range		
Precipitation, in inches		
Precipitation	4 08	9 10 1029, 21 190
	4 03	8 40, 1928; .31, 189-
Snow		
Max. precipitation in 24 hrs 2 23, 3rd, 4th	1.1	16 1903 1022 4 1906
Number of days with 01 or more8	11	16, 1892, 1933; 4, 1899
Wind, in miles	2127	1 271 1012 1 020 100
Total movement	3127	4,271, 1910; 1,920, 1894
Greatest daily movement192, 30th		
Least daily movement50, 23rd		
Mean hourly velocity4.5	4 2	
Maximum velocity28, 16th	22.7	35, 191.
Wind, direction	0111	*
Prevailing directionSW	SW	
North, days4		
Northeast, days1		
East, days		
Southeast, days4		
South, days		
Southwest, days6		
West, days		
Northwest, days6		
Weather		
Mean relative humidity, percent72.9	70.2	
Mean cloudiness, percent57	49 7	67, 1901; 27, 1923
Number of clear days21	9	17, 1921, 1936; 0, 1915, 1929
Number of fair days3	13	25, 1912; 5, '93, '33, '36
Number of cloudy days	9	18, 1901, '28; 2, '10, '2.
Number hours bright sunshine287	237	317, 1921; 152, 1915, 1929
Percent of possible hours of bright		
sunshine	55.2	
Thunder and lightning 3, 4, 9, 13, 16, 17		
18, 21, 25		
,,		

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1938

The weather during August was considerably warmer than usual with a mean temperature of 73.5 degrees. This is the highest mean for August in Amherst beginning in 1837. In August 1937 the mean was 73.3 degrees. The normal mean temperature for August is 68.6 degrees. Not only was the temperature unusually high but it was accompanied with high humidity on many days. The mean relative humidity for the month was 72.9 percent compared to a normal of 70.2 percent. The mean temperature for the three summer months June, July and August was 70.5 degrees. This has been exceeded only three times since 1837: in 1870 the mean temperature for the three months was 71.7 degrees; in 1876, 71.8 degrees; and in 1937, 70.6 degrees. The last three years have had the hottest summers since observations have been taken at the State College, beginning in 1889.

The rain fall during the month was 3.89 inches, the normal being 4.08 inches. Most of the rain fell in the form of local showers which were accompanied with an unusual amount of thunder and lightning.

There were 287 hours of bright sunshine compared to a normal of 237 hours.

Dr. J. K. Shaw of the Pomology Department gives the following report: "The weather during August continued reasonably favorable for fruit development. Timely showers prevented serious drought, and midseason and later varieties are sizing up well and developing good color. Probably due to the dry weather of late June and July, considerable internal cork is developing in certain varieties of apples and will cause serious losses in some orchards. The crop of McIntosh in the College orchard will be heavy and that of most other orchards fair to good. It is quite free of diseases but many apples show hail marks from the hail storm of carly summer."

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 609

September 1939

Meteorological Observations

FOR

SEPTEMBER 1939

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, 75th Mer.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

DAILY RECORDS

		Ten	npera	ture					Wind		1	Precipitati	on
	Max	imum	Min	imum			s of nshine		ļ ţ	s/hr*			
	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	No. Hours of Bright Sunshine	Prevailing Direction	Total Movement	Max. miles/hr*	Began	Ended	Rain
	82 86 80 81	3 P 3 P 3 P 3 P	54 61 61 61	5 A 6 A 1 A 5 A	68.0 73.5 70.5 71.0	78 88	10.3 7.4 4.8 8.2	se s s	95 128 114 145	9 28 11 10	3 Р	3:30 р	.39 T
-	76 69 74	3 P 3 P 4 P	56 49 51	12 P 6 A 0 A	66.0 59.0 62.5	82	4.4 10.5 12.8	s nw nw	183 190 142	21 25 18	9 A	12 P	.30
	60 74	10 A 4 P	51 46	0 A 7 A	55.5	87	$0.0 \\ 11.0$	sw	55 81	13 12	9 A	5 P	.51
	67 67 69 76 79 87	6 P 4 P 12 M 3 P 3 P 3 P 2 P	56 47 42 44 48 59 62	12 P 12 P 4 A 6 A 6 A 6 A	61.5 57.0 55.5 60.0 63.5 73.0	84 61 65 69 65 67	0.0 7.9 7.0 12.4 12.5 12.5 12.2	nw nw w se	104 220 78 57 133 112 114	32 25 9 8 11 8	11:30 A	4 P	.38
	74 67 71	12 M 5 P 3 P	48 41 35	12 P 12 P 5 A	61.0 54.0 53.0	70 62 68	5.0 12.2 10.0	n n se	228 157 89	25 14 12	7	0.20	Т
	74 66 73 82 76 76	3 P 3 P 3 P 2 P 4 P 2 P	45 45 37 48 48 54	4 A 12 P 6 A 6 A 7 A 3 A		67 75	7 3 6.0 9.7 10.0 7.5 2.8	s n sw sw ne s	114 103 61 175 57 161	13 11 9 17 8 20	7 P	9;30 Р	.12
	65 46 69 65	8 A 8 P 5 P 3 P	42 41 44 48	12 P 7 A 7 A 1 A	53.5 43.5 56.5 56.5	82 97 83	0.0 0.5 8.0 0.0	n n nw	181 104 60 98	22 25 8 9	12 m 3 A	11 P 10 P	.11 1.08
	82	3 P 2 P	56	0 A	69.0		5.1	se	188	13	9 P	10 P	.08

^{*}Based on least time required to blow one mile.

	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum30.56, 19th		30 65, 1924
Minimum		28 41, 1938
Mean semi-daily30.038		
Range	.88	1.99, 1938; .57, 1910
Air Temperature, in degrees F.	0.7.	
Highest		97, 1895, 1929
Lowest		24.5, 1914
Mean		
Range		
Highest mean daily		
Lowest mean daily		
Mean maximum73.5		
Mean minimum49.3		
Greatest daily range36, 19th, 22nd		
Least daily range	1 1	
Precipitation, in inches	1 26	14 77 4020 72 404
Precipitation2.97	4.26	14 55, 1938; .52, 1914
Snow		
Max precipitation in 24 hrs 1.08, 27th		16 1022 2 1002
Number of days with .01 or more8	10	16, 1933; 3, 1903
Wind, in miles	2271	1 606 1006 1 414 1004
Total movement		4 686, 1896; 1,414, 1894
Greatest daily movement228, 17th		
Least daily movement		
Mean hourly velocity		80 1030
Maximum velocity32, 10th Wind, direction	23.9	80, 1938
Prevailing directionSSW	WSW	
North, days		
Northeast, days		
East, days		
Southeast, days		
South, days		
Southwest, days		
West, days		
Northwest, days5		
Weather		
Mean relative humidity, percent 73.9	73 6	
Mean cloudiness, percent		70, 1934; 27, 1908, 1914
Number of clear days	10	19, 1932; 2, 1907, 1928
Number of fair days11	10	19, 1908; 3, 1889
Number of cloudy days6	10	20, 1934; 4, 1908, 1916
Number hours bright sunshine218		255, 1916; 106, 1934
Percent of possible hours of bright		,,0, 1001
sunshine	54.9	
Thunder and lightning 10th, 27th		
FrostsNone		
First Frost		Aug. 22, '94, '95; Oct. 13, '09

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1938.

The weather during September was nearly normal in most respects. There was less rain than usual, with 2.97 inches, the normal being 4.26 inches. The total rainfall since January 1 is now 29.40 inches compared to a normal of 33.40 inches for this period. The mean temperature for the month was 61.4 degrees, while the normal mean temperature for September is 61.7 degrees. The highest temperature was 91 degrees on the 16th and the lowest, 35 degrees on the 19th. There have been no killing frosts at the thermometer shelter, although there have been frosts on low ground in Amherst. The total wind movement for the month was slightly higher than normal and the highest wind velocity was 32 miles per hour on the 15th. There were 218 hours of bright sunshine which is 17 hours more than normal.

Dr. J. K. Shaw of the Pomology Department gives the following report: "The weather during September has continued to be favorable for fruit development and ripening. Rainfall was rather deficient but not seriously so. Apples have sized up well where trees were not overloaded and are developing good color. No general frosts have occurred. The peach crop has been practically all harvested and was good on the few trees that survived the severe winters of a few years ago. Our Baldwin apple trees have not borne well since the cold winter of 1933-1934. Winter injury is probably responsible, at least

in part, for the short crops.

GOODELL LIBRARY MASS.STATE COLLEGE AMHERST.MASS.

MASSACHUSETTS

10 3

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 610

October 1939

Meteorological Observations

FOR

OCTOBER 1939

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 262.3 ft. Height of wind instruments, 67 ft. Time used, 75th Mer.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

DAILY RECORDS

		Гen	pera	ture		İ			Wind			Precipitat	ion
	Max	imum	Min	imum		>	rs of inshine	8 t	nt	les/hr*			
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	No. Hours of Bright Sunshine	Prevailing Direction	Total Movement	Max. miles/hr*	Began	Ended	Rain
1	62 61	12 M	48 45	12 P 7 A	55.0 53.0		1.0 1.5	n n	183 230	18 17	2 в	3 в	.03
3	54 69	1 P 4 P	48 48	12 P 12 P	51.0 58.5	75 77	0.0 6.5	n n	193 123	13 10	9 A	5:30 г	.42
	76 75 70	3 P 2 P 3 P	46 54 48	6 A 12 P 6 A	61.0 64.5 59.0	90	6.8 4.5 11.5	sw se nw	44 117 116	7 13 14	6 1	7 1	T .50
	66 78	3 P 2 P	43 56	7 A 7 A	54.5 67.0	80 77	0.0 5.0	SW	53 112	10 22	11 F 0 A		1 1
	89 68 58	4 P 0 A 3 P	55 45 40	7 A 12 P 4 A	72.0 56.5 49.0	63	10.8 6.0 0.5	nw sw	106 228 74	32 22 8			
	56 56	4 P 3 P	40	7 A	52.0	62	11.0 5.0	N N N N N N N N N N N N N N N N N N N	93	13 32			T
	50 64	4 P 4 P	29 24	12 P 7 A	39.5 44.0	54 61	$\frac{11.0}{11.0}$	nw	227 92	20 8			
	56	12 M 4 P	32 26	7 A 6 A	37.0	47	3.3 10.0	nw nw	199 187	28 18			
	68 70	3 P	32 40	10 A 12 P		6.3	7.8 8 0	w.	228 165	18 17			
	75 64	4 P	34 42	7 A 12 P		86	3.8	s s	98	20 28	9:30 F 0 A		
	49 44 52	1 P 4 P 3 P	34 28 24	12 P 7 A 6 A	36.0	47	3.0 10.5 5.5	nw nw s	267 309 113	32 28 11	9 1	12 1	.09
	55	4 P	45	0 A	50.0	95	0.0	n	52	5	0 A	8 A	.03
	53 64 47	12 P 8 A 3 P	46 34 26	7 A 12 P 12 P		73	$0.0 \\ 4.7 \\ 10.3$	s w nw	54 275 240	7 39 25	8 A 0 A		.12
	47 45 44	3 P 4 P 12 M	23 42	5 A 8 A	34.0	87	0.0	nw	60 225	8 18	5 F 0 A		

^{*}Based on least time required to blow one mile.

October 1939	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum	30.53	30.68, 1929
Minimum	29.42	29 00, 1926
Mean semi-daily	30.06	2. 33, 1.2
Range1.13	1.11	1.47, 1926; .76, 1899
Air Temperature, in degrees F.		1.11, 1, 20, 1,0, 10,
Highest	79.4	90.5, 1908
Lowest		17, 1930
Mean	50.5	56.4, 1920; 43.2, 1890
Range	56.0	00.1, 1720, 43.2, 1090
Highest mean daily	00.0	
Lowest mean daily		
Mean maximum		
Mean minimum		
Greatest daily range		
Least daily range		
Precipitation, in inches		0.01 1011 01 100
Precipitation4.55		8.81, 1911; .01, 1924
Snow		
Max. precipitation in 24 hrs 175, 31st		
Number of days with 01 or more13	9	15, 1913; 1, 1897, 1924
Wind, in miles		
Total movement4792	4074	5,467, 1910; 2,540, 1894
Greatest daily movement309, 24th		
Least daily movement		
Mean hourly velocity 64		
Maximum velocity	29 5	42, 1937
Wind. direction		·
Prevailing direction	11.	
North, days5		
Northeast, days0	i	
East, days0		
Southeast, days		
South, days5		
Southwest, days		
West, days3		
Northwest, days		
Weather		
Mean relative humidity, percent 72.0	69 ()	
Mean cloudiness, percent	48	67, 1932; 18, 1924
Number of clear days9	10	21, 1938; 1, 191
Number of fair days	10	17, 1924; 3, 1938
Number of cloudy days	11	19, 1896; 3, 193
Number hours bright sunshine159		232, '23, '38; 91, 1913
Percent of possible hours of bright	110.0	202, 20, 30, 91, 1910
	51.3	
sunshine	31.3	
Thunder and lightning	Cont 21	Aug 22 '01 '05. Oct 12 100

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1938

The weather during October was seasonable in most respects. The mean temperature for the month was 50.2 degrees, the normal being 50.5 degrees. The lowest temperature was 23 degrees on the 30th. The first frost of the season occurred on October 15 which is the latest date for first frost recorded at this station. The normal date for the first frost is September 21. The domestic heating load for October was 460 degree-days. This, together with 155 degree-days in September, makes the heating load for the two months 615 degree-days. The normal for this period is 604 degree-days.

There was a deficiency in rainfall during each of the summer and fall months, beginning with May. As a result many wells went dry and town water supplies were threatened. This shortage of water continued through October until the 25th. Only 1.70 inches of rain fell in October until that date. Rains during the last week brought the total for the month to 4.55 inches. The normal rainfall for

October is 3.29 inches.

There were 159 hours of bright sunshine compared to a normal of 175 hours. The highest wind velocity was 39 miles per hour on the 28th.

Dr. J. K. Shaw of the Pomology Department makes the following comment: "The month of October marks the close of the 1939 season for fruits. The season has been generally favorable for the crops. The apple crop was very heavy, especially of McIntosh, and that of other fruits, fair to good. While the rainfall was deficient at times, causing some injury to shallow-rooted crops, fruits suffered but little from lack of water. There were no unseasonable spring or fall frosts. Hail caused damage in some orchards. The fruit grower should be well satisfied with the season of 1939, except in the very important matter of prices which are discouragingly low."

GOODELL LIBRARY MASS.STATE COLLEGE AMHERST.MASS.

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 611

November 1939

Meteorological Observations

FOR

NOVEMBER 1939

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 262.3 ft. Height of wind instruments, 67 ft. Time used, 75th Mer.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

1

DAILY RECORDS

		Геп	ıpera	ture					Wind		F	recipitati	on	
	Max	imum	Min	imum			s of nshine		1	es/hr*				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	No. Hours of Bright Sunshine	Prevailing Direction	Total Movement	Max. miles/hr*	Began	Ended	Rain	Snow
1 2 3 4 5 6 7 8 9 10 11 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 30 30 30 30 30 30 30 30 30 30 30 30	52 54 50 38 46 50 55 40 55 40 55 40 45 45 40 38 49 40 46 48 48 43	3 P 4 P 2 P 8 P 11 A 3 P 0 A 3 P 11 A 3 P 0 A 3 P 2 P 0 A 3 P 2 P 0 A 3 P 2 P 11 A 3 P 2 P 2 P 11 A 3 P 2 P 11 A 3 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2	32 27 23 28 35 34 36 26 23 11 23 23 23 23 21 26 31 26 24 24 26 24 26 26 26 26 26 26 26 26 26 26 26 26 26	12 P 12 P 12 P 15 A 10 P 11 P 11 A 12 P 12 P 12 P 12 P 12 P 6 A 7 A 12 P 7 A 12 P 6 A A 7 A 12 P 7 A 8 A 12 P 8 P 12 P 12 P 12 P 12 P 12 P 12 P	40.5 36.5 36.5 33.0 40.5 42.0 40.0 44.0 38.0 41.5 28.0 41.5 39.5 39.5 30.0 30.5	81 67 72 96 74 67 65 59 63 57 69 69 63 64 63 60 82 74 77 77 77 77 73 72 49 40	5.3 5.0 2.0 10.2 0.0 4.5 3.0 0.5 6.2 3.3 6.0 7.0 8.0 7.0 7.0 9.5 9.5 9.5 9.5 9.5 9.5 9.5 9.6 9.6 9.6 9.6 9.6 9.6 9.6 9.6	w nw nw nw nw w w w nw nw w w w nw nw w w w ne n ne n ne n nw nw w w w	1722 488 1155 51 1447 2433 1588 165 321 1522 343 3224 1588 115 1457 488 115 15221 666 237 2388 2322 1866 90 30	15 6 14 5 25 22 14 25 40 18 32 20 28 32 7 25 32 14 13 20 20 22 21 5 22 21 21 21 21 21 21 21 21 21 21 21 21	1 P 0 A 9 A	12 P 1 A	.92 .01 .05 T	Т

^{*}Based on least time required to blow one mile.

November 1939	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum	30.59	30 87, 1932
Minimum	29.30	28 73, 1904
Mean semi-daily	30.05	,
Range1.12	1.29	1.84, 1904; .94, 1905
Air Temperature, in degrees F.		
Highest	66.0	75, 1938
Lowest	12.9	-4, 1938
Mean	38 9	44.1, 1931; 33.6, 1901
Range43	53.0	
Highest mean daily45.5, 8th		
Lowest mean daily28, 14th		
Mean maximum46.7		
Mean minimum26.9		
Greatest daily range39, 16th		
Least daily range		
Precipitation, in inches		
Precipitation	3.41	8 64, 1927; 63, 1917
SnowTrace	2.34	13.50, 1938
Max. precipitation in 24 hrs93, 5th, 6th		
Number of days with 01 or more 3	9	22, 1921; 2, 1904
Wind, in miles		
Total movement	4531	5 978, 1906; 2 589, 1889
Greatest daily movement343, 11th		
Least daily movement30, 30th		
Mean hourly velocity 6 9	6.3	
Maximum velocity	30.3	44, 1938
Wind direction		
Prevailing directionNW	WNW	
North, days6		
Northeast, days2		
East days0		
Southeast, days0		
South, days1		
Southwest, days		
West, days9		
Northwest, days		
Weather		
Mean relative humidity, percent 67 8	70.6	
Mean cloudiness, percent	55.1	72, 1900, 1927; 34, 1917
Number of clear days	6	15, 1903; 1, 1900, 1911, 1927
Number of fair days	9	16, '12; 4, '89, '30, '32, '35
Number of cloudy days	15	24, 1927; 9, 1905, 1917
Number hours bright sunshine	121	182, 1903; 66, 1927
Percent of possible hours of bright		
	41.3	
sunshine53.9		I .
Thunder and lightning		
	Nov. 6	Oct. 10, '25; Nov. 27, '31

Note —The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1938

The outstanding features of November weather were the large amount of sunshine and the lack of precipitation. Only .98 inch of precipitation was recorded during the month, the normal for November being 3.41 inches. In the storm on the 5th, .93 inch of rain fell. This is the least precipitation ever recorded in November at this station, with the exception of November 1917, when .63 of an inch fell. No snow fell during the month, with the exception of a trace on the 9th and the 13th. The normal snowfall for November is 2.34 inches and the normal date for first snow is November 6. Last year 13.50 inches fell in November, which was the heaviest on record for November.

The mean temperature for the month was 36. 8 degrees, the normal being 38.9 degrees. The lowest temperature was 18 degrees on the 14th, compared with a normal minimum of 12.9 degrees for November.

The domestic heating load for November was 846 degree-days; the normal, 783 degree-days. The load for the heating season thus far is 1461 degree-days, while the normal for this period is 1387 degree-days.

There were 158 hours of bright sunshine, the normal for November being 128 hours. Only three other years since 1889 had more hours of sunshine in November, viz., 1936, 161 hours; 1917, 170, and 1903, 182.

GOODELL LIBRARY MASS.STATE COLLEGE AMHERST.MASS.

10 37

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 612

December 1939

Meteorological Observations

FOR

DECEMBER 1939

C I GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 262.3 ft. Height of wind instruments, 67 ft. Time used, 75th Mer.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

DAILY RECORDS

	Т	empera	ture					Wind		1	Precipitati	on	
	Maximun	Min	imum			s of inshine	50 _	nt	es/hr*				
Day	Degrees	Degrees	Time	Mean	Mean Relative Hunidity	No. Hours of Bright Sunshine	Prevailing Direction	Total Movement	Max. miles/hr*	Began	Ended	Rain	Snow
1 2 3 4 5 6 7 8	53 12 53 0 43 0 40 1 42 12 1 45 1 43 0	P 19 P 37 A 40 A 33 P 30 D 27 P 27 A 23 P 21	7 A 2 A 5 P 7 P 5 A 11 P 6 A 12 P 4 A	45.0 46.5 38.0 35.0 34.5 36.0 33.0	100 84 66 55 74 86 57	9.3 0.0 0.5 0.0 3.0 1.0 0.5 3.8 1.3	w w sw nw nw sw s nw se	35 170 317 164 46 220 473 290	5 9 22 22 15 5 22 39 22	11 P 3 A 1 A	12 P 12 P 5 P	·03 1.46 .09	Т
9 10 11 12 13 14 15	38 0 32 4 37 2 34 6 30 2	33 A 16 P 15 P 17 A 18 P 5	4 A 2 A 12 P 8 A 4 A 12 P 7 A 0 A	38.5 27.0 23.5 27.0 26.0 17.5	80 89 69 83 74 86	0.0 0.0 9.1 0.0 5.5 9.0 1.2	s	64 258 284 70 163 74 143	8 28 25 9 20 5	3.30 P 0 A 11 A	12 P 2 P 10 P	.23 .19 .28	
16 17 18 19 20 21 22 23	39 10 43 2 38 1 38 6 41 11 31 0 31 12	P 30 P 30 P 26 P 32 A 30 A 21 M 16	6 A 12 P 7 A 6 A 11 P 8 A 12 P	34.5 36.5 32.0 35.0 35.5 26.0 23.5	100 73 91 100 81 59 56	0.0 0.0 2.0 0.0 2.0 3.0 9.0	s nw n n w	64 140 62 135 191 375 376	6 17 5 14 15 28 39	10 A	11 A 6 P	.11 T 1.23	T
24 25 26 27 28 29 30 31	32 2 26 2 19 3 31 3 25 8 35 2	P 12 P 19 P 8 P 1 P 5 P 9 P 16 P 14	6 A 1 A 12 P 7 A 8 A 5 A 12 P 11 P	19.0 25.5 17.0 10.0 18.0 17.0 25.5 23.5	6 62 54 0 67 0 47 0 76 5 97	9 0 5 0 9.0 0.0 9.2 0.0 5.6 2.0		256 253 300 167 144 42 63 138	25 32 22 18 13 4 13 17	1 A	9 а	.27	3.00

^{*}Based on least time required to blow one mile.

December 1939	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum30.45, 15th	30.66	30,96, 1889
Minimum 29.27, 7th, 8th	29.27	28 85, 1913
Mean semi-daily	30.06	, ,
Range1.18	1.38	1.78, 1895; 1.01, 189;
Air Temperature, in degrees F. Highest	54.3	65.5, 190
Lowest	-19	- 22.5, 191
Mean28.9	27.5	36.9, 1891; 17.1, 191
Range	56.2	30.9, 1891, 17.1, 191
Highest mean daily		1
Lowest mean daily		
Mean maximum		
Mean minimum		
Greatest daily range		
Least daily range		
Precipitation, in inches	2.20	# ## 4004 OF 400
Precipitation3.89		7.77, 1901; .87, 1890
Snow	8.50	26.50, '02; t, 189
Max. precipitation in 24 hrs1 48, 2nd, 3rd		
Number of days with 01 or more 9	10	17, 1902; 4, 1892
Wind, in miles		
Total movement5519	4710	6,694, 1925; 3,239, 1918
Greatest daily movement 473, 8th		
Least daily movement		
Mean hourly velocity	6.3	
Maximum velocity39, 8th, 23rd	31.4	48, 193
Wind. direction		
Prevailing direction	WNW.	
North, days 5		
Northeast, days0		
East, days0		
Southeast, days1		
South, days6		
Southwest, days		
West, days5		
Northwest, days		
Weather		
Mean relative humidity, percent 75.3	69.4	
Mean cloudiness, percent 56		71, 1929; 39, 191
Number of clear days	8	15, 1890; 2, 1922, 193.
Number of fair days	9	16 '00:4 '80 '30 '31 '36 '3
Number of cloudy days	14	16, '09;4, '89, '30, '31, '36, '36, '37, '09, '23, '33; 7, '09, '23, '33, '33; 7, '09, '23, '33, '33, '33, '33, '33, '33, '33
Number hours bright sunshine		172, 1896; 63, 193.
Percent of possible hours of bright	120	172, 1070, 03, 193
sunshine	45.2	
Suisinie	13.2	
		I

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1938.

ANNUAL SUMMARY

Annual 1939	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum30.68, Feb. 16 and 17	30.81	31.05, 1920
Minimum	28.95	28.41, 1938
Mean semi-daily	30.01	
Range1.48	1.85	2.47, 1938; 1.38, 1933
Air Temperature, in degrees F.		
Highest	95.7	104 1911
Lowest	-12.2	26, 1904
Mean48.0	47.4	49.8, 1921; 44, 1904
Range97	107.8	
Highest mean daily 81, July 7, Aug. 21	1	
Lowest mean daily		
Mean maximum	57.8	
Mean minimum	36.5	
Greatest daily range	50.5	
Least daily range		
Description in inches		
Precipitation, in inches Precipitation	43.70	59.00, 1938; 30.68 1908
27.50		89.00, 1893; 24.50, 1919
Snow	31.10	09.00, 1093, 24.00, 1919
Max. precipitation in 24 hrs. 2.23, Aug. 3, 4	1.2.1	144, 1902; 96, 1924
Number of days with .01 or more 110	124	144, 1902; 90, 1924
Wind, in miles		(2.571 1000 24.357 1001
Total movement 51,734	52,223	63,571, 1908; 36,257, 1894
Greatest daily movement478. Mar. 7		
Least daily movement	- 0	
Mean hourly velocity 5.9	5.8	90 1029
Max. velocity, 48. Feb.4 16 Mar.7. June17	39.5	80,1938
Wind, direction	11.	
Prevailing direction WNW	11.	
North, days		
Northeast, days		
East, days		
Southeast, days		
South, days		
Southwest days	1	
West, days45		
Northwest, days		
Weather		
Mean relative humidity, percent 69.8	67.6	
Mean cloudiness, percent	51.7	60, '98, '01, '02; 4 1, '08, '24
Number of clear days	116	169, 1936; 59, 1927
Number of fair days 123	123	182, 1912; 64, 1936
Number of cloudy days 87	126	179, '01, '02; 71, 1910
Number hours bright sunshine2511	2.353	2838, 1924; 1864, 1902
Percent of possible hours of bright		
sunshine	52.8	
Last frost May 16	May 14	Apr. 23, '04; June 8, '32
Girst frost Oct. 15	Sept. 21	Aug. 22, '94, '95; Oct. 13,'09
	1 1 7	1 Nov. 14 '10: Nov. 14 '07
Last snow April 9 First snow Nov. 9	Apr. 15	Oct. 10, '25; Nov. 27, '31

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1938

GOODELL LIBRARY
MASS.STATE COLLEGE
AMHERST.MASS.

10

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 613

January 1940

Meteorological Observations

FOR

JANUARY

1940

C. I. GUNNESS

OBSERVATORY

Latitude, 42–23′ 48.5″ N. Longitude, 72–31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 262.3 ft. Height of wind instruments, 67 ft. Time used, 75th Mer.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

DAILY RECORDS

		Ten	ipera	ture					Wind		P	recipitati	on	
	Maxi	imum	Min	imum			s of nshine		- L	s/hr*				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	No. Hours of Bright Sunshine	Prevailing Direction	Total Movement	Max. miles/hr*	Векап	Ended	Rain	Snow
1 2 3 4 5 6 7 8 9 10 11 12 3 14 15 16 17 18 9 21 22 23 24 5 26 27 8 2 2 30 31	25 21 31 26 22 28 32 28 31 30 35 37 42 31 15 18 22 23 33 30 35 22 22 23 31 32 23 31 32 23 31 31 24 25 31 26 27 27 28 28 28 31 28 28 31 28 28 31 31 31 31 31 31 31 31 31 31 31 31 31	2 P 11 A 2 P 2 P 3 P 3 P 3 P 3 P 3 P 4 P 6 A 1 A 4 P 8 P 2 P 2 P 1 P 2 P 2 P 2 P 4 P 6 P 1 P 4 P 2 P 4 P 4 P 6 P 7 P 8 P 8 P 9 P 9 P 9 P 9 P 9 P 9 P 9 P 9	7 11 18 8 3 11 3 12 5 -3 -1 22 29 3 3 3 2 11 3 4 4 4 0 9 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	8 A A A A A A A A A A A A A A A A A A A	16.0 24.5 17.0 12.5 19.5 14.0 15.5 12.3 15.0 26.0 32.0 37.0 21.0	67 69 62 50 81 74 62 68 89 90 88 90 88 91 97 88 97 88 97 88 97 87 87 87 87 87 87 87 87 87 8	9.11 7.00 8.5 9 2 0 1 0 1 8.00 0.00 0.00 0.00 0.00 7.5 5.00 0.00 0.	nw n se nw nw nw sw n s sw sw n s w nw nw nw nw nw nw	152 207 307 233 39 294 156 146 145 33 33 50 82 165 176 368 263 175 192 221 110 268 122 256 182 147	12 25 22 28 9 11 11 3 3 4 5 18 28 18 10 14 11 17 25 14 20 22 32 18 33 5 28	1.30 P 7.30 A 3 A 0 A 3 P 0 A	8 P 8 P 12 P 8 A 12 P 10 A	.07 .05 .31 .05 .89 1.22	1.00

^{*}Based on least time required to blow one mile.

	 I	
January 1940	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum30.53, 10th	30.70	31.00, 1927
Minimum	29.20	28.55, 1913
Mean semi-daily	30.07	
Range1.00	1.50	2.18, 1913; 0.97, 1896
Air Temperature, in degrees F.		
Highest	51.1	66.0, 1932
Lowest	-7.1	-26.0, 1904
Mean	24.2	34.2, 1913; 13 9, 1918
Range45	58.2	
Highest mean daily		
Lowest mean daily		
Mean maximum		
Mean minimum		
Greatest daily range32, 11th		
Least daily range		
Precipitation, in inches		
Precipitation	3.61	7.15, 1898; 1 07, 1896
Snow	13 24	
Max. precipitation in 24 hrs 2 11, 14th, 15th	15 24	33,1898, 1923; 1.50, '08
	11	15 103 120 1 1001
Number of days with 01 or more 7	11	15, '93, '20; 4, 1901
Wind. in miles	5055	5550 1000 2007 1005
Total movement	5055	7770, 1908; 2896, 1895
Greatest daily movement		
Least daily movement 33, 10th, 11th		
Mean hourly velocity	6.8	
Maximum velocity39, 28th	32.7	47, 1938
Wind. direction		
Prevailing directionWNW	WNW	
North, days		
Northeast, days0		
East, days		
Southeast, days		
South, days		
Southwest, days		
West, days		İ
Northwest, days 19		
Weather		
Mean relative humidity, percent 74 8	70.1	
Mean cloudiness, percent 42	5.5	78, 1932; 37, 1908
Number of clear days	9	17, 1939; 2, 1914
Number of fair days. 9	0	18, 1890, 1926; 3, 1916
Number of cloudy days 8	1.3	22, 1923, 1931; 4, 1920, 1939
Number hours bright sunshine	137	214, 1920; 74, 1932
Percent of possible hours of bright	107	219, 1920, 79, 1932
sunshine	46.7	
эмпэник	10.7	
		•

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1939

The weather during January was featured by the continued cold spell which began the last week in December. January was the coldest since 1920, with a mean temperature of 18.5 degrees compared to 15.2 degrees in 1920. The highest temperature was 42 degrees on the 15th, compared to a normal high for January of 51.1 degrees. The lowest temperature was -3 degrees on the 10th, compared to a normal low of -7.1 for January. The mean maximum temperature for the month was 27.1 degrees and the mean minimum, 9.8. Only one other day, the 11th, was below zero with a reading of -1: but only 5 days were above 32 degrees. While January was cold, it had a total of 171 hours of sunshine compared to the normal of 137 hours. This is the highest number of hours recorded in the month of January since 1924. The domestic heating load for the season to the end of January is now 4021 degree-days, whereas the normal number of degree-days for this period is 3814, giving an excess of about 5 percent. The total precipitation for the month was 2.63 inches. The normal for January is 3.61 inches. A total of 5.75 inches of snow fell during the month, compared to the normal of 13.24 inches. The total wind movement for the month was 5091 miles, the normal being 5055 miles.

GOODELL LIBRARY MASS.STATE COLLEGE MASSACTUSETTS

10 37

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 614

February 1940

Meteorological Observations

FOR

FEBRUARY

1940

C. L. GUNNESS

OBSERVATORY

Latitude, 42 23' 48.5" N. Longitude, 72 31' 10" W.

Height of barometer above ground, 36 ft. Above sea level, 262.3 ft.

Height of wind instruments, 67 ft. Time used, 75th Mer.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

DAILY RECORDS

1		Гетрегатиге							Wind		Precipitation					
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	No. Hours of Bright Sunshine	Prevailing Direction	Total Movement	Max. miles · hr*	Векап	Ended	Rain	Snow		
1 2 3 4 5 6 7 8 9 0 1 1 2 3 1 4 5 6 7 8 9 0 1 1 2 3 1 4 5 6 7 8 9 0 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	34 30 31 33 40 37 40 38 32 41 46 3 5 33 31 42 41 35 33 37 37 34 41 35 35 36 37 37 37 38 38 37 37 37 37 37 37 37 37 37 37 37 37 37	1 P 4 P 2 P 3 P 2 P 12 M 4 P 12 M 14 P 12 P 12 M 4 P 12 P 14 A 11 A 1 A 2 P 4 A 1 A 1 A 1 A 1 A 1 A 1 A 1 A 1 A 1 A 1	10 17 11 13 20 19 24 27 29 24 32 25 19 20 16 15 18 32 20 10 11 15 17 7	3 A 6 A 8 A 12 P 8 A	23.0 30.0 28.0 36.5 33.5 28.0 36.5 39.0 32.0	79 67 71 87 93 75 60 60 100 80 70 68	5.8 6.00 8.5 5.9 1 5.3 2.00 1.0 8.0 7.0 0.0 9.0 3.1 5.7 0.0 9.0 9.0 10.6 5.0 10.9 6.5 0.5 4.4 11.0 6.0 5.0 11.0	nw nw nw nw n s n n	81 204 257 130 34 76 188 83 201 174 263 572 359 91 146 3153 1483 153 146 134 200 233 66 98 47	7 22 28 11 3 13 13 25 25 17 9 25 18 17 40 39 18 25 18 29 20 20 8 9 5 5	2 P 0 A 3 A 0 A 5 A 8 A 0 A 0 A 8 P	12 P 3 A 12 P 7 A 10 P 12 P 2 P 2 A	.38 .02 .51 .22 1.01 .37 .14 .02	T 2.5 9.0 1.0 T 0.5 T		

^{*}Based on least time required to blow one mile

February 1940	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum	30.63	31 05, 1920
Minimum	29.24	28 56, 1895
Mean semi-daily	30.04	
Range	1.38	1.89, '00, '08; .88, '13, '31
Highest	50.5	65.0, 1930
Lowest	-7.4	=22.5, 1918
Mean	23.7	,
Range	58	
Hignest mean daily 39, 12th		
Lowest mean daily		
Mean maximum	32.8	
Mean minimum18.4	14.5	
Greatest daily range		
Least daily range		
Precipitation, in inches		
Precipitation	3,19	8.12, 1900; 0.62, 1901
Snow	13.99	48.75, 1893; 0.50, 1937
Max. precipitation in 24 hrs 1.01, 14th		
Number of days with .01 or more 9	10	15, '93, '20; 4, 1901
Wind, in miles	"	10, 50, 20, 1, 1501
Total movement	4776	6445, 1896; 3438, 1892
Greatest daily movement572, 15th	17.0	0110, 1070, 0100, 1072
Least daily movement		
Mean hourly velocity	7.1	
Maximum velocity	31.7	48, 1934, 1937
Wind. direction		10, 1701, 1701
Prevailing directionNW	WNW	
North, days9		
Northeast, days		
East, days 0		
Southeast, days		
South, days		
Southwest, days		
West, days		
Northwest, days		
Weather		
Mean relative humidity, percent	66,6	
Mean cloudiness, percent 54	50.9	66, '90, 27; 31, 1905
Number of clear days	10	17, '19, '24; 2, 1927
Number of clear days	8	16, 1920; 2, 1936
Number of cloudy days9	10	16, 1894; 2, 1930
Number hours bright sunshine161	158	221, 1924; 110, '27, '38
Percent of possible hours of bright	100	221, 172x, C10, 21, 30
synshine	53.4	
5.T.S	.,,,,,,	
	1	

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1939

The month of February was warmer than normal and was marked by a severe blizzard on the 14th when 9 inches of snow fell and much drifting was caused by high winds. In the Boston area and other castern parts of the state 14 inches of snow fell which drifted badly and caused serious traffic tie-ups.

The mean temperature was 26.7 degrees, while the normal is 23.7 degrees. The maximum temperature of 46 degrees occurred on the 12th and the minimum was -6 degrees on the 27th. The normal maximum temperature for February is 50.5 degrees and the normal minimum temperature, -7.4 degrees. The total precipitation of 2.72 inches was under the normal of 3.19 inches. A total of 15.50 inches of snow fell as compared to the normal of 13.99 inches. The heating load for the month was 1112 degree-days. The normal load for February is 1156 degree-days. The heating load thus far is 5133 degree-days; and the normal load for this same period is 4970 The total wind movement was 5459 miles. degree-days. normal for February is 4776 miles. This marks the largest wind travel for February since 1933 when 6029 miles were recorded. February 1918 had 5632 miles of wind travel. The maximum wind velocity of 40 miles per hour was recorded on the 14th, the day of the blizzard. A total of 161 hours of bright sunshine was slightly higher than the February normal of 158 hours.

10 3

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 615

March 1940

Meteorological Observations

FOR

MARCH

1940

C. I. GUNNESS

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft.

OBSERVATORY

Heights of wind instruments, 67 ft. Time used E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

MONTHLY WEATHER SUMMARY

	Temperature						Wind]	Precipit	ation		
Day	Maximum	Time	Minimum	Time	Mean	Humidity	Sunshine	Direction	Movement	Maximum	Began	Ended	Water	Snow
1 2 3 4 5 6 6 7 8 9 10 11 12 13 14 15 16 17 18 20 21 22 23 24 25 27 28 29 30 31 31 31 31 31 31 31 31 31 31 31 31 31	33 35 36 33 38 39 39 40 43 31 41 41 41 37 39 47 47 36 22 24 28 39 34 46 44 43 59 46 47 47 47 47 47 47 47 47 47 47 47 47 47	2 P 4 P 5 P 4 P 1 P 1 A 12 M 2 P 4 P 4 P 3 P 4 P 4 P 1 P 4 P 1 P 4 P 1 P 4 P 4 P 1 P 4 P 1 P 4 P 4 P 4 P 4 P 1 P 4 P 1 P 4 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P 1	18 5 29 29 28 31 22 28 22 19 21 11 13 17 29 21 22 23 27 23 19 11 10 11 15 23 23 23 23 24 25 26 27 27 27 27 27 27 27 27 27 27	12 P 6 A 12 P 3 A 12 P 7 A 6 A 12 P 7 A 7 A 7 A 12 P 6 A 12 P 6 A 12 P 6 A 12 P 6 A 12 P 6 A 12 P 6 A 12 P 6 A 12 P 12 P 12 P 13 A 14 A 15 P 16 A 17 A 18 A 18 A 18 A 18 A 18 A 18 A 18 A 18	25.5 20.0 32.5 31.0 33.0 35.0 32.5 24.5 20.0 27.0 29.0 31.0 29.5 38.5 36.0 27.5 16.5 17.0 21.5 31.0 29.5 37.0 38.0	63 64 87 91 94 89 82 73 75 66 64 57 61 73 88 87 60 85 71 60 85 77 62 79 65 66 65 77 67 67 67 67 67 67 67 67 67 67 67 67	6.5 11.2 0.0 0.0 2.2 0.0 5.3 8.0 8.5 10.0 11.6 11.6 3.3 0.0 7.5 8.0 7.5 8.5 10.3 2.0 11.5 10.5 8.4 12.3 0.0 0.0 5.5 10.0 5.5 8.5	n w w w n w n w n w n w n w n w n w n w	127 80 55 147 194 97 133 85 190 326 384 140 49 270 156 103 31 23 31 226 150 354 16 17 18 27 19 19 10 10 10 10 10 10 10 10 10 10 10 10 10	14 75 15 13 8 15 11 18 18 28 21 15 15 16 21 21 21 21 21 21 21 21 21 21	9:30	1 8 A 8 A 8 A 8 A 8 A 10 P 1 P A 1 P A 12 P	.06 1.42 .03 .18 .05	T .50 3.00 1.00 .50 .25 T T .25

^{*}Based on least time required to blow one mile.

March 1940	Normal	Extremes
Barometer, in inches		•
(Readings reduced to sea level)		
Maximum30.59, 2nd	30.57	30.96, 1904
Minimum	29.24	28.47, 1914
Mean semi-daily	30.00	
Range	1.34	2.10, 1914; 0.85, 1915
Air Temperature, in degrees F.		, , , ,
Highest	63.8	79.5, 1907, 1921
Lowest	6.1	-7.5, 1906
Mean	34.4	,
Range	57.6	
Highest mean daily 48.5, 31st		
Lowest mean daily		
Mean maximum	43.2	
Mean minimum	25.4	
Greatest daily range	2.7.1	
Least daily range		
Precipitation, in inches	3.70	7.13, 1899; 0.12, 1917
Precipitation	7.47	
Snow	1.41	27, 1899; 0, 1921
Maximum precipitation in 24 hrs.		
1.48, 15th, 16th		17 1000 1010 1000 0 101
Number of days with .01 or more 13	11	[17, 1890, 1913, 1936; 3, 1915]
Wind, in miles	W () 4 4	0100 1000 0000 100
Total movement	5944	8182, 1896; 3006, 1903
Greatest daily movement 384, 11th		
Least daily movement 49, 14th		
Mean velocity	7.7	
Maximum velocity	31.2	48, 1932, 1939
Wind, direction		
Prevailing direction WNW	HXH	
North, days $\dots \dots		
Northeast, days		
East, days0		
Southeast, days0		
South, days		
Southwest, days		
West, days		
Northwest, days		
Weather		
Mean relative humidity, percent 73.6	64.2	
Mean cloudiness, percent		
Number of clear days		22, 1924; 3, 190
Number of fair days		17, 1900; 4,1895, 1897, 1930
Number of cloudy days		21, 1901; 1, 191
		292, 1924; 93, 190
Number hours bright sunshine	199	202, 1024, 00, 100
Percent of possible hours of bright	59 C	
sunshine53.6	53.6	
Thunder and lightning		

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1939.

March was much colder than normal, with more rain and snow than usual. The only warm, pleasant day was the last day of the month.

The mean temperature for the month was 30.1 degrees, compared to the normal of 34.4 degrees for March. This was the coldest March since 1926 when the mean temperature was 29.4 degrees. Last year the mean temperature for March was 30.9 degrees, with the month almost identical in character with March of this year.

The highest temperature was 59 degrees on the 31st and the

lowest, 5 on the 2nd.

The mean temperature for the winter months December through March was 26 degrees compared to a normal of 27.5 degrees for this same period. The lowest temperature recorded during the winter was -6 on February 27.

The domestic heating load for March was 1082 degree-days, compared to the normal load of 949 degree-days. The total load for the months September through March is 6215 degree-days, while the normal for this same period is 5919 degree-days.

The total precipitation was 5.58 inches, while the normal for March is 3.70 inches. The snowfall for the month was 9.00 inches, compared to the normal of 7.47 inches. The total snowfall for the season thus far is 35.00 inches; the normal for this period is 45.54 inches.

The total hours of sunshine were 199, which is the normal for March. The wind velocity was 5493 miles for the month, compared to the normal of 5944 miles.

GOODELL LIBRARY MASS.STATE COLLEGE MASSAMERSTETITS:

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 616

April 1940

Meteorological Observations

FOR

APRIL

1940

C. I. GUNNESS

C. I. GUMMESS

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

OBSERVATORY

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

10

MONTHLY WEATHER SUMMARY

		Ter	mpera	ture				Wind			F	recip i t	ation	
Day	Maximum	Time	Minimum	Time	Mean	Humidity	Sunshine	Direction	Movement	Maximum	Began	Ended	Water	Snow
1 2 3 4 4 5 6 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 223 224 25 26 27 28 29 30	47 52 42 49 51 55 47 58 55 53 33 40 51 57 60 58 42 39 38 49 51 57 60 58 47 64 77 77 77 77 77 77 77 77 77 77 77 77 77	1 P 4 P 2 P 2 P 2 P 2 P 4 P 2 P 2 P 11 A M 2 P 2 P 1 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P	34 27 32 33 30 34 36 41 33 24 20 32 33 32 33 32 33 33 30 31 33 31 31 31 32 31 31 31 31 31 31 31 31 31 31 31 31 31	12 P 6 A 8 A 11 P 4 A 3 A 6 A 12 P 5 A 12 P 12 P 5 A 6 A 12 P 12 P 2 A 4 A 3 A 6 A 12 P 5 A 6 A 12 P 5 A 6 A 12 P 12 P 12 P 12 P 12 P 12 P 12 P 12 P	40.5 39.5 40.5 41.0 41.5 44.5 44.5 44.0 28.5 39.0 49.5 40.0 43.5 35.5 35.0 41.5 45.0 45.0 45.0 45.0 56.5		10.3 12.7 9.0 0.0 6.5 12.9 13.0 0.0 2.6 13.1 11.3 0.0 8.4 11.0 10.5 0.0 9.5 0.0 0.0 6.5 1.5 0.0 9.5 1.5 0.0 1.5 0.0 1.5 0.0 1.5 0.0 1.5 0.0 1.5 0.0 1.5 0.0 0.0 1.5 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0	nw w nw nw nw nw nw nw se nw nw se nw nw se n n n n n n n n n s se	390 122 90 82 256 339 185 97 156 269 143 220 374 180 146 131 129 78 288 349 311 79 97 106 138 173 90 164	$\begin{array}{c} 32\\ 15\\ 13\\ 8\\ 28\\ 40\\ 17\\ 15\\ 22\\ 20\\ 15\\ 18\\ 32\\ 15\\ 8\\ 17\\ 8\\ 17\\ 15\\ 22\\ 25\\ 15\\ 25\\ 15\\ 25\\ 18\\ 8\\ 9\\ 17\\ \end{array}$	8 A A 3 A A 0 A A 0 A	1 A 12 P 4 A 12 P 8 A 8 P 8 P 12 P 8 P 12 P	.055.666.077 .833.511 1.333.32 .055 .744 .855.22 .666 .033.05 T T	.50 2.00 1.50 2.50

^{*}Based on least time required to blow one mile.

April 1940	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum	30.48	30.71, 1911
Minimum		28.99, 1907, 1929
Mean semi-daily	29.99	20100, 1001, 100
Range	1.11	1.52, 1930; .72, 1919
Air Tomporature in degrees F	1.11	1.02, 1000, .,2, 1010
Air Temperature, in degrees F.	79.4	00 5 1000
Highest		88.5, 1896
Lowest	22.0	8.5, 1923
Mean	45.7	52, 1921; 41.2, 1926
Range	57.4	
Highest mean daily		
Lowest mean daily $\dots 28.5$, 13th		
Mean maximum51.6	56.8	
Mean minimum	34.8	
Greatest daily range 40, 29th		
Least daily range 5, 20th		
	1	
Precipitation, in inches	9.95	6 90 1020 76 1909
Precipitation	3.35	6.89, 1929; .76, 1892
Snow	2.17	11, 1891; 0, 1910, 193
Maximum precipitation in 24 hrs.		
1.66, 12th, 13th		
Number of days with $.01$ or more 14	11	18, 1909, 3, 1892
Wind, in miles		
Total movement	5404	8208, 1908; 3853, 1917
Greatest daily movement		, , , ,
Least daily movement		
Mean hourly velocity 7.6	7.5	
Maximum relacity	31.4	40, 1935, 193
Maximum velocity	31.4	40, 1955, 1956
Wind, direction	33737337	
Prevailing directionNNW	WNW	
North, days8		
Northeast, days1	1	
East, days	i	
Southeast, days		
South, days4		
Southwest, days		
West, days1		
Northwest, days		
Weather		
	61.6	
Mean relative humidity, percent 67.3	61.6	7- 1001. 24 1025 1025
Mean cloudiness, percent	51.8	75, 1901; 34, 1925, 1927
Number of clear days13		19, 1911; 3, 1898, 1902
Number of fair days	9	18, 1915; 2, 1894, 190
Number of cloudy days10	10	22, 1901; 3, 1899, 192
Number hours bright sunshine 207	220	296, 1911; 103, 190
Percent of possible hours of bright		
sunshine	54.7	
Thunder and lightning		
Last snow	Apr 15	Mar. 14, 1910: May 11, 1907
	TARDER KID	interest and a state of a state o

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1939.

The past month was one of the coldest and wettest Aprils on record at this station.

The mean temperature for the month was 42.1 degrees, compared to the normal of 45.7 degrees. It was colder than this only twice since 1889, the year the station was first established, viz., 41.2 in 1926 and 41.7 in 1907. In 1917 it was also 42.1 degrees. The highest temperature was 74 on the last day of the month, and the lowest, 20 on the 14th. The lateness of the season was indicated by the fact that the winter ice on the college pond remained until the 10th.

The domestic heating load for the month was 686 degree-days, while the normal for April is 579 degree-days. The total heating load for the season thus far is 6902 degree-days, compared to the normal of 6498 for this period.

The precipitation for the month totaled 6.37 inches, the normal being 3.35 inches. This is the heaviest for April since 1929 and was exceeded only three times in the past 50 years. A total of 6.50 inches of snow fell, compared to the normal of 2.00 inches for April. The snow fell in two storms; the last one totalling 4 inches occurred on April 21 and 22. This is the heaviest snowfall on record at such a late date. The total snowfall for the season was 41.50 inches compared to the normal seasonal fall of 47.71 inches.

The sunshine and wind travel for the month were very nearly normal.

GOODELL LIBRARY MASS.STATE COLLEGE

MASSACHUSETTS: AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 617

May 1940

Meteorological Observations

FOR

MAY

1940

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

> Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

10

MONTHLY WEATHER SUMMARY

	Temperature								Wind		P	recipit	ation	
Day	Maximum	Time	Minimum	Time	Mean	Humidity	Sunshine	Direction	Movement	Maximum	Began	Ended	Water	Snow
1 2 3 4 4 5 6 6 7 8 9 10 11 12 13 144 15 6 17 188 19 20 22 23 24 25 26 27 28 29 30 31	71 62 70 54 68 70 69 68 67 78 63 67 82 67 68 63 75 77 68 64 69 65 76 68 65 76 69 69 69 69 69 69 69 69 69 69 69 69 69	3 P P 4 P 4 P P 4 P P 4 P P 4 P P 4 P P 4 P P 4 P P 4 P P 4 P P 4 P P 4 P P 4 P P 4 P P 4 P P 4 P	43 55 53 49 48 40 49 43 45 43 43 43 43 43 43 41 55 55 49 50 50 50 50 50 50 50 50 50 50 50 50 50	6 A A 6 A 6 A 6 A 6 A 6 A 6 A 6 A 6 A 6	57.0 58.5 61.5 58.0 55.0 56.5 57.0 56.5 51.0 55.0 65.0 58.0 66.5 62.5 55.0 66.5 66.5 66.5 66.5 57.5 57.0 58.0	$\begin{array}{c} 57 \\ 93 \end{array}$	6.5 0.0 2.0 0.0 13.0 8.3 13.6 11.3 14.4 6.0 8.5 9.3 2.2 13.0 0.0 0.1 1.5 5.5 14.0 4.4 2.0 0.2 0.3 0.0 0.0 1.5 5.5 14.0 0.0 0.0 0.0 0.0 1.5 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0	s se se s n n n s mw n m n n sw n mw n mw sw e se n me n e n e n e n e n e s s e n w sw s se e n me n e s s e n m m e s s e n m m e s s e n m m e s s e n m m e s s e n m m e s s e n m m e s s e n m m e s s e n m m e s s e n m m e n e s e n m m e n e n e s e n m m e n e n e n e n e n e n e n e n	755 2266 1844 1400 1755 3144 94 134 134 112 2207 66 38 131 114 246 65 95 141 173 167 130 187 71 145 76 96	20 17 10 17 15 11 40 10 18 11 20 9 14 13 8 11 20 9 14 13 8 7 11 15 11 11 11 11 11 11 11 11 11 11 11	7 P O A A 11:30 A 3 A 5:30 A	8 A 7 P 8 A A	.133 .511 T .100 .04 .299 T T T .055 .044	

^{*}Based on least time required to blow one mile.

May 1940	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
$Maximum \dots 30.27, 30 th$	30.39	30.62, 193
Minimum	29.51	29.10, 193
Mean semi-daily	29.96	
Range	0.88	
Air Temperature, in degrees F.		
Highest	86.1	94.5, 1896, 191
Lowest	31.2	24.0, 190
Mean	57.1	
Range	54.9	
Highest mean daily 66.5, 20th		
Lowest mean daily 50.5, 10th		
Mean maximum		
Mean minimum		
Greatest daily range		
Least daily range 5, 4th, 23rd		
Precipitation, in inches	1	
Precipitation 5.67	3.60	7.44, 1931; .48, 1903
Maximum precipitation in 24 hrs. 2,58, 31st	0.00	1111, 1001, 110, 101
Number of days with .01 or more 12	12	20, 1901; 5, 1903
Wind, in miles		20,1001,0,100
Total movement	4504	5946, 1907; 2180, 189
Greatest daily movement 314, 7th	1001	5510, 1501, 2100, 185
Least daily movement		
Mean hourly velocity	6.1	
Maximum velocity	28.8	45, 193
Wind, direction	20.0	40, 199
Prevailing direction ESE	W	
North, days		
Northeast, days		
East, days		
Southeast, days		
South, days		
Southwest, days		
West, days		
Northwest, days ϵ		
Weather	00.5	
Mean relative humidity, percent 70.3		5 0 1001 90 100
Mean cloudiness, percent	52.1	70, 1901; 30, 192
Number of clear days		20, 1923; 0, 192
Number of fair days	11	17, 1907, 1926; 5, 192
Number of cloudy days	9	20, 1927; 3, 1903, 192
Number hours bright sunshine 182	252	334, 1922; 137, 192
Percent of possible hours of bright		
sunshine40	55.6	
Thunder and lightning	1	_
Last frost	May 14	June 8, 1932; Apr. 23, 190-
·		
	1	

The weather during May was marked with rain above normal and less sunshine than usual. The total rainfall for the month was 5.67 The rainfall during the storm on the 31st was 2.58 inches. The normal rainfall for May is 3.60 inches. This makes a total rainfall of 17.62 inches for March, April and May. This has been exceeded only once in the 50-year history of the station. In 1901 there was a total rainfall of 18.52 inches in these three spring There were only 182 hours of bright sunshine compared to the normal of 252 hours. Most of the deficiency occurred in the latter half of the month. Only three times during the last 50 years has there been less sunshine in May than the past month. there were 137 hours; in 1916, 181; and in 1901, 159. The temperature for the month was nearly normal with a mean of 57.2 degrees. The normal mean temperature for May is 57.1 degrees. There were no heavy frosts, the lowest temperature being 33 degrees on the 13th. The highest temperature was 82 degrees on the 15th. domestic heating load was 242 degree-days, the normal being 245 degree-days. The total heating load for the season is now 7143 degree-days, the normal for this period being 6743 degree-days.

Dr. J. K. Shaw of the Pomology Department makes the following report: "Although the past winter was a severe one, fruit plants came through with little injury. Peach fruit buds survived and the bloom was abundant. Raspberry canes suffered much less injury than in the previous winter. Apple varieties blossomed moderately or heavily, especially considering the heavy crop of last year which might be expected to limit fruit bud formation. Quite likely the dry sunny weather of last June favored fruit bud formation. season of bloom was nearly a week later than the average and, because of much cold, cloudy weather, was much prolonged. were a few sunny days favorable to bee flight and pollination. cannot yet be said how well apples have set. The abundance of wet weather during the early season of growth favored scab infection and made scab control difficult."

MASS.STATE COLLEGE AMHERST.MASS.

10 37

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 618

June 1940

Meteorological Observations

FOR

JUNE

1940

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

MONTHLY WEATHER SUMMARY

		Ter	npera	ture				,	Wind		Pr	ecipitat	ion	
Day	Maximum	Time	Minimum	Time	Mean	Humidity	Sunshine	Direction	Movement	Maximum	Ведап	Ended	Water	Snow
1 2 3 4 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 22 12 22 32 4 22 5 26 27 28 29 30	79 79 79 85 89 89 82 86 57 75 88 79 83 75 77 78 83 65 64 70 74 62 73 76 79 74	4 P 3 P 4 P 2 P 4 P 5 P 12 M 4 P 9 A A 1 P P 12 M 4 P 12 M 4 P 12 M 4 P 12 M 4 P 14 P 15 P 14 P 15 P 14 P 15 P 15 P	56 59 53 61 63 59 49 56 56 57 56 66 53 57 50 48 48 49 49 49 49 49 49 49 49 49 49 49 49 49	5 A 12 P 5 A 6 A 12 P 7 5 A A 12 P 5 A A 12 P 5 A A 12 P 5 A A 12 P A 5 A A 12 P A 5 A A 12 P A 5 A A 12 P	67.5 69.0 69.0 75.0 76.0 76.5 67.5 57.5 69.5 57.0 66.0 71.5 58.0 54.5 54.5 58.0 56.0 54.5 60.5 54.5 57.0 60.5 60.5 60.5 60.5 60.5 60.5 60.5 60	72 65 51 66 64 49 90 94	10.2 14.0 15.1 12.5 11.5 15.2 15.2 15.2 17.7 3.4 0.0 0.0 2.6 6.9 15.3 2.5 12.5 12.5 12.5 12.5 12.5 12.5 12.5	sw w w w sw nw nw w sw nw nw nw nw nw w sw nw nw nw nw nw w sw nw w sw w w sw w w sw w w w	S5 1066 83 777 799 2699 118 153 1077 1077 40 52 210 135 170 132 116 178 164 268 107 143 57 771 83 129 125 179 127 179 179 179 179 179 179 179 179 179 17	$\begin{array}{c} 9 \\ 10 \\ 13 \\ 8 \\ 11 \\ 28 \\ 13 \\ 13 \\ 22 \\ 10 \\ 4 \\ 8 \\ 28 \\ 13 \\ 22 \\ 13 \\ 14 \\ 14 \\ 22 \\ 0 \\ 17 \\ 18 \\ 9 \\ 7 \\ 10 \\ 18 \\ 14 \\ 25 \\ 20 \\ \end{array}$	7 A 4 A 0 A 1 P 7 A 12:30 P 11 A 7 A 5 A 1:30 P 4 P 0 A 9:30 P	12 P 2 A	.10 .34 .06 .05 .22 .12 .14 .45 .19 .14 .19	

^{*}Based on least time required to blow one mile.

June 1940	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum	30.30	30.54, 1889
Minimum	29.55	29.24. 1909
Mean semi-daily	29.95	
Range	0.75	
Air Temperature, in degrees F.	01.0	101 1014
Highest	91.2	101, 1919
Lowest	40.0	34, 189
Mean	65.7	
Range	51.3	
Highest mean daily		
Lowest mean daily 53.5, 21st	l i	
Mean maximum	1	
Mean minimum		á
Greatest daily range	!	
Least daily range	1	
Precipitation, in inches Precipitation	3.75	9.68, 1922; 0.76, 1908
Maximum precipitation in 24 hrs 49,	0.10	<i>5.00, 1522, 0.70, 100.</i>
24th, 25th	1	
Number of days with .01 or more	11	17, 1922; 4, 1908
Wind, in miles		17,1022, 1,1000
Total movement	3585	4571, 1908; 1409, 1906
Greatest daily movement 269, 6th		
Least daily movement		
Mean hourly velocity 5.4	5.0	
Maximum velocity	24.6	48, 1939
Wind, direction		
Prevailing direction	WSW	
North, days		
Northeast, days		
East, days		
Southeast, days1		
South, days		
Southwest, days		
$\underbrace{ ext{West}}_{, ext{days}}, \underbrace{ ext{days}}_{, ext{.}}$		
Northwest, days	1	
Weather Mean relative humidity, percent	66.9	
Mean cloudiness, percent		71, 1903; 28, 1903
Number of clear days		22, 1908; 1, 192
Number of fair days9	1 7 2 1	23, 1912; 4, 1895, 190
Number of cloudy days		22, 1903; 1, 192
Number hours bright sunshine		362, 1908; 102, 190
Percent of possible hours of bright	~	33_,, -3_, -3
sunshine	54.0	
Thunder and lightning4th, 5th.		
13th, 15th, 19th, 27th, 28th, 29th.		
	1	l .

The weather during June was in reverse seasonal order. The first seven days were warm and clear, with a mean temperature for the week of 70.6 degrees as compared to the normal of 62.4 degrees for that week. This period was followed by several days with nearly normal temperatures but with many showers, the month closing with a ten-day period of some of the coldest weather on

record for this time of the year.

The mean temperature for the month was 64.6 degrees, the normal for June being 65.7 degrees. The highest temperature was 89 on the 4th and 5th; and the lowest 39 on the 22nd, which is an unusually low reading for the last part of June. The precipitation totaled 2.46 inches, compared with the normal of 3.75 inches for June. Rain was recorded every day from the 24th to the 30th. The total precipitation from January 1 to June 30 is now 25.43 inches, while the normal for this period is 21.20 inches. There were 261 hours of sunshine, the normal for June being 257 hours. The total wind movement was 3894 miles, which is 309 miles greater than the normal for June.

Dr. J. K. Shaw of the Pomology Department reports: "While the precipitation during June was below normal, there were no dry periods sufficient to noticeably check the growth of vegetation. There were several rainy periods favorable for apple scab infection, and on trees not well sprayed much scab developed. The strawberry harvesting season was at its peak at the end of the month when frequent showers favored good size of berries but caused some decay. In spite of much unfavorable weather during apple bloom, there was a satisfactory set of fruit and there is promise of a good

crop though less than last year."

GOODELL LIBRARY MASS.STATE COLLEGE AMHERST . MASS . MASSACHUSETTS

10 37

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 619

July 1940

Meteorological Observations

FOR

JULY

1940

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

MONTHLY WEATHER SUMMARY

		Те	mpera	ture					Wind		Precipitation			
Day	Maximum	Time	Minimum	Time	Mean	Humidity	Sunshine	Direction	Movement	Maximum	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 6 17 18 19 20 1 22 23 24 5 26 27 28	69 73 76 68 82 85 82 81 75 82 77 82 87 82 87 88 88 88 88 88 88 88 88 88 88 88 88	2 P 2 P 2 P 6 P 4 P 4 P 2 P 4 P 4 P 2 P 4 P 4 P 2 P 5 P 5 P 7 S P 5 P 7 S P 5 P P P 5 P P 5 P P P 5 P P P 5 P P P	50 48 46 55 50 49 50 59 57 55 49 47 47 57 66 67 63 63 72 70	4 A A 5 A A 5 A A 4 A A A 5 A A 5 A A 5 A A 5 A A 5 A 5	59.5 69.5 61.0 61.5 64.0 65.5 67.5 72.0 74.0 69.5 63.0 64.5 67.0 69.0 74.5 78.5 74.5 75.5 73.5 80.5	67 68 68 89 74 61 61 74 70 82 92 75 62 75 93 99 88 85 85 91 88 91 87 91 88 91 88 91 88 91 88 91 88 91 91 91 91 91 91 91 91 91 91 91 91 91	6.0 8.5 9.8 1.0 9.9 15.1 15.2 15.0 7.5 0.0 14.7 12.5 6.0 14.7 12.5 4.5 10.0 2.1 5.0 6.0 11.5 4.5 10.0 1	nw nw s nw nw sw s s s s s s s s s s s s	1644 911 700 966 844 811 566 933 1055 677 644 1099 999 744 605 737 788 599 744 600 1366 1211 733 544	25 20 9 11 25 13 7 11 28 10 9 12 12 12 12 12 12 17 10 8 5 12 17 10 11 10 11 10 11 11 11 11 11 11 11 11	2:30 P 10 P 0 A 3:30 P 3 P 0 A 8:30 A 6:00 P 5:30 A 3 A 3 P 1:30 A	3 P 12 P 8:30A 4 P 8 P 1 P 9 A 6:30P 6 P 5:30P 3:30A 7 P	.03 T .14 .48 .49 .13 .27 .07 1.18 .29 .54 .32 .28	
29 30 31	89 93 85	2 P 2 P 2 P	67 71 58	5 A 0 A 12 P	$78.0 \\ 82.0 \\ 71.5$	76 81 72	$\begin{array}{c} 5.0 \\ 7.3 \\ 14.5 \end{array}$	s sw nw	82 118 162	$\frac{10}{17}$	3 р 3:55 р	10:30р 4 Р	.33 .12	

^{*}Based on least time required to blow one mile.

July 1940	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum	30.27	30.50, 1892
Minimum	29.59	29.27, 1932
Mean semi-daily	29.96	
Range	0.68	0.97, 1892; 0.47, 1938
Air Temperature, in degrees F.		
Highest	93.9	104, 1911
Lowest	46.4	40, 1890, 1898
Mean	70.8	74.7, 1921; 66.3, 1891
Range	47.7	111, 1121, 0010, 1111
Highest mean daily		
Lowest mean daily		
Mean maximum		
]	
Mean minimum		
Greatest daily range		
Least daily range		
Precipitation, in inches	4.10	14.51 1005 050 1006
Precipitation	4.10	14.51, 1897; 0.70, 1929
Maximum precipitation in 24 hrs. 1.18, 18th		
Number of days with $.01$ or more 15	11	20, 1915; 4, 192-
Wind, in miles		
$\operatorname{Total} \ \operatorname{movement} \ldots 2869$	3422	5097, 1909; 1109, 189-
Greatest daily movement 164, 1st		
Least daily movement 54, 2 th		
Mean hourly velocity	4.6	
Maximum velocity	28.7	44, 1936
Wind, direction		,
Prevailing direction W	SW	
North, days3		
Northeast, days0		
East, days		
Southeast, days0	1	
South, days		
Southwest, days		
West, days		
Northwest, days		
Weather	00.4	
Mean relative humidity, percent76.3		20 1014 91 109
Mean cloudiness, percent 61	50.9	69, 1914; 31, 192
Number of clear days10		22, 1923; 0, 1917
Number of fair days	14	24, '09, '21; 5, '89, '23, '28
		'36
Number of cloudy days		18, 1889; 0, 1910
Number hours bright sunshine 259	268	371, 1910; 180, 1931
Percent of possible hours of bright		1
sunshine	58	
Thunder and lightning 1st, 9th		
11th, 18th, 22nd, 24th, 15th, 26th, 29th		
30th.		1

The weather during the first two weeks in July was cold, while the last half of the month was unusually hot. As a result, the average temperature for the month was practically normal, the mean being 70.6 degrees. The normal mean temperature for July is 70.8 degrees. The highest temperature during the month was 94 degrees on the 26th. The normal maximum for July is 93.9 degrees. weather during the last 10 days was made uncomfortable by the high humidity which accompanied the relatively high temperature. The mean relative humidity for the month was 76.3 percent, compared to a normal of 68.4 percent. A similar spell of hot weather was experienced from the 4th to the 11th day of July in 1937. The temperature was higher during that period than during the hot days in the past month, but the humidity was not as high. rain on 15 days during the month but many of these rains were light showers. A very heavy rain of 1.18 inches fell in half an hour on July 18. The total rainfall during the month was 4.69 inches, the normal for the month being 4.10 inches. The total rainfall since January 1 is now 30.12 inches, the normal for this period being 25.30 inches. There were 259 hours of bright sunshine compared to a normal of 268 hours. The total wind movement during the month was 2869 miles. This is considerably less than normal, as the average wind movement for July is 3422 miles.

GOODELL LIBRARY MASS.STATE COLLEGE AMHERST, MASS. MASSACHUSETTS

10 37

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 620

August 1940

Meteorological Observations

FOR

AUGUST

1940

C. I. GUNNESS

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

OBSERVATORY

MONTHLY WEATHER SUMMARY

	Temperature							Wind			Precipitation			
Day	Maximum	Time	Minimum	Time	Mean	Humidity	Sunshine	Direction	Movement	Maximum*	Began	Ended	Water	t
1 2 3 4 4 5 6 6 7 8 9 0 1 1 2 2 3 3 4 4 5 5 1 5 1 5 1 5 1 5 1 5 1 5 1 5 1 5	77 79 81 83 85 87 88 85 86 89 87 84 82 83 86 82 87 76 63 70 71 72 67 78 84	3 P 3 P 4 P 4 P 3 P 12 M 4 P 4 P 3 P 12 M 7 P 12	54 54 55 56 65 65 66 66 66 66 66 67 53 55 69 59 48 46 40 41 45 59 68	5 A 5 A 5 A 6 A 6 A 6 A 12 P 12 P 6 A 6 A 12 P 12 P 12 P 12 P 12 P 12 P 12 P 12	65.5 66.5 67.0 75.0 77.5 71.5 71.0 73.5 74.0 72.0 69.5 68.0 67.0 75.5 71.5 60.0 64.0 51.5 56.0 66.5 56.5 76.0	66 61 62 775 77 83 74 66 72 66 73 80 71 65 72 80 85 64 71 70 74 65 62 84 84	9.0 14.0 14.4 10.6 2.8 6.6 8.5 14.2 12.0 14.1 10.0 14.0 12.0 10.3 1.0 1.5 6.5 7.5 7.5 6.0 7.0 2.0 8.0 5 7.0 2.0 8.0 7.0 2.0 8.0 7.0 8.0 8.0 7.0 8.0 8.0 8.0 8.0 8.0 8.0 8.0 8.0 8.0 8	n ne se s se n nw nw se se se n n s se se n n w nw w w w w w se nw s	15.5 123 8.5 122 163 176 6.5 51 120 120 10.5 79 99 99 91 11.5 182 59 66 141 161 60 91 74 45 103 103 104 104 105 105 106 106 106 106 106 106 106 106 106 106	18 12 9 12 12 9 13 7 6 6 6 8 10 9 12 8 10 9 11 9 11 9 11 9 11 10 10 10 10 10 10 10 10 10 10 10 10	1 P 2 P 0 A	6:30a	.50 .06 .60 .13	

^{*}Based on least time required to blow one mile.

August 1940	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum	30.32	30.50, 1934
Minimum	29.61	28.87, 1930
Mean semi-daily	30.00	
Range	0.71	
Air Temperature, in degrees F.		
lighest	91.6	100, 1918
Lowest	43.4	37, 1894, 1908
Mean		
Range	48.1	
dighest mean daily		
Lowest mean daily		
Mean minimum		
Greatest daily range		
Least daily range		
Precipitation, in inches	4.08	8.40, 1928; .31, 1894
Maximum precipitation in 24 hrs. 0.60, 19th		5.40, 1926, .31, 169-
Number of days with .01 or more 8	11	16, 1892, 1933; 4, 1899
	11	10, 1002,1000, 4, 100
Wind, in miles	9197	4 971 1010 1 090 190
Total movement		4,271, 1910; 1,920, 189-
Greatest daily movement 182, 10th Least daily movement		
Mean hourly velocity		
Maximum velocity		35, 191
The state of the s	22.1	55, 151
Wind, direction Prevailing direction	$_{ m SW}$	
North, days		
East, days		
Southeast, days		
South, days		
Southwest, days		
West, days		
Northwest, days		
Weather		
Mean relative humidity, percent72.3	70.2	
Mean cloudiness, percent		67, 1901; 27, 1923
Number of clear days		21, 1939; 0, 1915, 1929
Number of fair days		25, 1912; 3, 1939
Number of cloudy days		25, 1912; 3, 1939 18, 1901, '28; 2, '10, '2
Number hours bright sunshine		317, 1921; 152, 1915,1929
Percent of possible hours of bright	İ	
sunshine	55.2	
Thunder and lightning6th, 13th, 23rd		
First Frost	Sept. 21	Aug. 22,'94,'95; Oct 13,'09

The weather during August was colder than usual with a mean temperature of 66.9 degrees. The normal mean temperature for August is 68.6 degrees. The lowest temperature was 34 degrees on the 25th. This is the lowest August temperature recorded at this station. The nearest approach to this minimum was 37 degrees on August 22, 1894 and August 29, 1908. There was frost on August 25 on low land in town, particularly in South Amherst, with damage to tobacco, corn and other crops. The highest temperature was 89 degrees on the 10th. Rainfall was considerably less than normal, being 1.56 inches. The normal rainfall for August is 4.08 inches. There were 263 hours of bright sunshine compared to a normal of 237 hours. The wind movement was 3219 miles which is slightly above normal. The mean relative humidity for August was 70.2 percent.

Dr. J. K. Shaw of the Pomology Department makes the following report: "Apples are gaining size somewhat slowly due to cool weather and perhaps in some cases lack of abundant rains. Some varieties are a week or more later than the average. These conditions, especially cool nights, have favored color development in red varieties so that early and mid-season varieties are showing much better color development than usual. Early peach varieties have matured good crops and later varieties are coming on well though somewhat later than normal. Pears are making a heavier crop than usual. The blueberry crop is about over and the yield was rather light due to winter injury to the plants. The frost of August 25 that caused much injury to tender crops did not harm fruit crops."

GOODELL LIBRARY MASSACHISETTS AMHERST. MASS.

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 621

September 1940

Meteorological Observations

FOR

SEPTEMBER

1940

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

DAILY RECORDS

		Te:	mpera	ture		y.			Wind '			Precipitation				
	Maxi	imum	Min	imum		Iumidit	f Hours ashine		_	×			-			
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Rain	Snow		
$\begin{array}{c} -1234567891011231156789222222222222222222222222222222222222$	82 78 78 75 72 65 66 66 67 75 72 71 73 80 81 84 85 76 76 76 77 77 77 78 78 78 78 78 78 78	2 P 5 P 2 P 3 P 3 P 3 P 4 P 2 P 4 P 2 P 3 P 2 P 4 P 3 P 2 P 4 P 3 P 2 P 4 P 2 P 4 P 3 P 2 P 4 P 3 P 2 P 4 P 4 P 4 P 4 P 4 P 4 P 4 P 5 P 6 P 6 P 6 P 6 P 7 P 8 P 8 P 8 P 8 P 8 P 8 P 8 P 8 P 8 P 8	72 66 60 51 50 45 45 51 44 41 44 46 50 50 47 49 63 24 44 57 49 63 24 49 40 40 40 40 40 40 40 40 40 40 40 40 40	11 P 12 P 12 P 6 A 6 A 6 A 6 A 6 A 12 P 5 A 6 A 5 A 6 A 5 A 6 A 5 A 6 A 7 A 12 P	77.0 72.0 69.0 63.0 58.5 55.0 58.5 54.5 54.5 58.5 61.5 64.0 64.0 67.0 64.0 67.0 55.5 55.5 65.0 64.0 67.0 58.5 74.0 65.5 65.0 65.5 65.5 65.5 65.5 65.5 65	75 79 92 96 75 76 82 82 82	1.3 2.7 6.9 8.9 11.9 10.7 9.3 12.5 0.0 0.5 10.5 7.5 6.4 10.0 10.0 10.0 11.2 12.4 9.5 7.7 3.1 12.2 12.1 0.0 11.9 11.9 11.9 11.9 11.9 11.9 11.	s n nw n nw nw w e n nw w se sw n nw w sw m nw nw w se n nw nw nw nw nw nw nw nw nw nw nw nw n	1822 833 855 1299 666 588 800 744 104 166 120 711 333 433 438 228 655 218 228 844 146 156 157 158 15	20 18 22 22 23 25 20 9 6 10 11 13 5 17 18 22 9 6 8 10 25 20 10 20 20 20 20 20 20 20 20 20 2	1 Р	7:30p 2:30P 1 P	.08 .43			

^{*}Based on least time required to blow one mile.

September 1940	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum	30.45	30.65, 192
Minimum	29.57	28.41, 193
Mean semi-daily	30.06	
Range	.88	1.99, 1938; .5 7, 19 1
Air Temperature, in degrees F.		
fighest	87.7	97, 1895, 192
$Lowest \dots 30, 27th$	33.2	24.5, 191
Mean	61.7	
Range	54.5	
Highest mean daily		
Lowest mean daily		
Mean maximum72.6		
Mean minimum		
Greatest daily range		
Least daily range		
Precipitation, in inches	i	
Precipitation	4.26	14.55, 1938; .52, 191
Maximum precipitation in 24 hrs. 0.79, 25th		
Number of days with .01 or more 5	10	16, 1933; 3, 190
Wind, in miles		,, -, -,
Total movement	3271	4,686, 1896; 1,414, 189
Greatest daily movement 228, 22nd	0	1,000, 1000, 1,111, 100
Least daily movement	i i	
Mean hourly velocity		
Maximum velocity32, 5th, 26th		80, 193
	2.,	00, 130
Wind, direction Prevailing direction	WSW	
	11:011	
North, days		
East, days	!	
Southeast, days	1	
South, days		
Southwest, days4		
Vest, days		
Northwest, days	1	
Weather	* 0.0	
Mean relative humidity, percent79	73.6	
Mean cloudiness, percent	49.9	70, 1934; 27, 1908, 193
Number of clear days16	10	19, 1932; 2, 1907, 193
Number of fair days	10	19, 1908; 3, 188
Number of cloudy days	10	20, 1934; 4, 1908, 19
Number hours bright sunshine 239	201	255, 1916; 106, 195
Percent of possible hours of bright		
sunshine	54.9	
Thunder and lightning 21st		
First frostSept. 27	Sept. 21	[Aug. 22,'94,'95; Oct. 13,'(

The outstanding feature of the weather for September was the small amount of rain. The total for the month was 1.53 inches, as compared to the normal for September of 4.24 inches. Only three times in the past 52 years has the rainfall for September been less. In 1926, only 1.50 inches fell; in 1916, 1.37 inches; and in 1914, .52 inch. The total rainfall since January 1 is now 33.21 inches, compared to a normal of 33.62 inches.

The mean temperature for the month was 60.1 degrees, the normal being 61.7 degrees. The highest temperature was 85 degress on the 21st. The lowest was 30 degrees on the 27th, and this was the first killing frost of the season at the observatory. There have been earlier frosts, however, on low ground in the town of Amherst. There were 239 hours of bright sunshine compared to a normal of 201 hours. The mean humidity was 79 percent, while the normal mean humidity is 73.6 percent. The wind movement was very nearly normal, with no high winds.

Dr. J. K. Shaw of the Pomology Department makes the following report: "The retarded development of apples and other fruits noted last month continued through September, and all varieties are more or less late in coming to maturity. There is much less pre-harvest drop of apples than usual and the fruit has colored well. A general frost on the night of September 27 killed late tender crops but did no harm to fruit crops. Coming later than usual, the frost did no more damage to field and garden crops than usual."

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 622

October 1940

Meteorological Observations

FOR

OCTOBER

1940

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

DAILY RECORDS

		Te	mpera	ture		ŕ			Wind		Pr	ecipitat	ion	
	Maxi	imum	Min	imum		lumidi	Hourshine			x				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	педед	Ended	Rain	
1 2 3 4 5 6 7 8 9 10 11 2 13 14 15 16 17 18 19 22 12 22 3 24	60 69 68 62 69 69 69 76 63 64 65 70 76 72 171 53 52 47 42 49 43 47 61 66	4 P 2 P 4 P 3 P 4 P 5 A A 5 P 5 A P 1 P 1 P 1 P 1 P 2 P 2 P 2 P 2 P 2 P 2	38 46 50 38 30 50 58 53 42 36 36 39 29 22 28 26 36 36 36 36 36 36 36 37 38 38 38 38 38 38 38 38 38 38	7 A 3 A 7 A 12 P 7 A 6 A 12 P 12 P 6 A 7 A 12 P 7 A 12 P 7 A 12 P 7 A 12 P 7 A 12 P	49.0 57.5 59.0 50.0 49.5 67.0 58.0 53.0 51.5 55.0 41.0 37.5 37.5 37.5 38.5 34.5 33.0 43.0	88 87 78 81 86 86 95 79 82 83 81 77 94 65 88 87 49 57 62 63 54	0.4 2.0 6.5 12.5 13.0 6.5 6.5 0.0 11.4 11.3 10.1 11.0 7.5 11.1 2.0 11.1 7.5 10.9 6.3 10.8 6.7 10.7	n n n n n s s s s s n n sw s s nw nw nw nw nw nw nw nw nw nw nw nw nw	107 140 136 148 72 194 201 101 54 69 182 240 94 74 144 208 121 214 96 192 180	10 15 20 14 8 13 17 9 13 13 7 9 20 10 28 18 13 32 15 13 25 13 17 20	8 а	9 р	.15 T	c
25 26 27 28 29 30 31	53 53 47 46 56 40 52	3 P 12 M 12 M 2 P 4 P 8 P 2 P	30 28 22 19 21 32 36	5 A 12 P 4 A 6 A 7 A 0 A 4 A	41.5 40.5 34.5 32.5 38.5 36.0 44.0	73 62 70	$\begin{bmatrix} 0.1 \\ 7.5 \\ 10.5 \\ 6.3 \\ 9.8 \\ 0.0 \\ 3.9 \end{bmatrix}$	n nw n n n n	36 164 148 127 43 36 148	25 17 18 5	7 A		.31	

^{*}Based on least time required to blow one mile.

October 1940	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum	30.53	30.68, 192
Minimum	29.42	29.00, 192
Mean semi-daily $\dots \dots 30.081$	30.06	
Range	1.11	1.47, 1926; .76, 189
Air Temperature, in degrees F.		,,,
Highest	79.4	90.5, 190
owest	23.4	17, 193
Mean	50.5	56.4, 1920; 43.2, 189
Range	56.0	, , , , , , , , , , , , , , , , , , , ,
lighest mean daily 67.0, 7th		
Lowest mean daily		
Jean maximum		
Jean minimum		
Freatest daily range		
Least daily range		
Precipitation, in inches	3.29	0.01 1011. 01 109
Precipitation	3.29	8.81, 1911; .01, 19:
Maximum precipitation in 24 hrs. 0.53, 30th		
51St		17 1019 1 1007 10
Number of days with .01 or more 4	9	15, 1913; 1, 1897, 19
Wind, in miles		
otal movement	4074	[5,467, 1910; 2,540, 189]
Freatest daily movement 240. 15th		
east daily movement 36, 25th, 30th	ļ	
Jean hourly velocity		
Jaximum velocity	29.5	42, 19
Wind, direction		
Prevailing direction NW	II.	
North, days12		
Vortheast, days0	1	
Cast, days0		
outheast, days	!	
outh, days		
outhwest, days	İ	
Vest, days		
Vorthwest, days9	ļ	
Weather	1	
	coo	
Jean relative humidity, percent 76.7		67 1039, 10 10
Mean cloudiness, percent	48	67, 1932; 18, 19
Number of clear days		21, 1938; 1, 19
Number of fair days	10	17, 1924; 3, 193
Number of cloudy days	11	19, 1896; 3, 199
Number hours bright sunshine	175.0	232, '23, '38; 91,19
Percent of possible hours of bright	-10	
	-51.3	
sunshine	01.0	1
sunshine 64.5 Chunder and lightning First frost Sept. 27		

The weather during October was clear, cool and dry. Rain in excess of .01 inch fell on only 4 days, although there were traces of rain on 7 additional days. The total rainfall for the month was 1.04 inches, the normal rainfall for October being 3.29 inches. The total precipitation since January 1 is now 34.25 inches. The normal for this period is 36.91 inches.

The mean temperature for the month was 46.7 degrees, the normal being 50.5 degrees. This is the coldest October since 1925 when it was 43.5 degrees. The highest temperature was 76 degrees on the 7th and 12th. The lowest temperature was 19 degrees on the 22nd and 28th. Only twice during the last 52 years has the temperature been lower than 19 degrees in October. In 1936 it fell to 17 degrees and in 1904, to 18 degrees. The domestic heating load for October was 566 degree-days. There were 183 degree-days in September. The total for September and October is, therefore, 751 degree-days. The normal for these two months is 604 degree-days. There were 220 hours of bright sunshine compared with a normal of 175 hours. There were no unsually high winds.

Dr. J. K. Shaw of the Department of Pomology gives the following report: "Sub-normal temperatures continued through October and there were several sharp freezes during the last half of the month. Some apples remained on the trees during one or more of these freezes and those going through more than one or two of them must have been injured. The total amount of such fruit was small. Late varieties did not size up well. The leaves were frozen prematurely and this was not favorable for the trees and makes them less able to withstand severe freezing if such should occur this winter. Rainfall was also below average and unless abundant rains occur before the ground freezes, the trees will go into the winter under somewhat unfavorable conditions."

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 623

November 1940

Meteorological Observations

FOR

NOVEMBER

1940

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

DAILY RECORDS

		Te	mper	iture		ž	Wind				P	recipita	tion	
	Max	imum	Min	imum	_	lumidi	Hour			ys.				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Numl er of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Rain	Snow
$\frac{1}{2}$ $\frac{3}{4}$	55 48 53 66	3 P 4 P 2 P 1 P	28 39 38 36	7 A 12 P 1 A 6 A	41.5 43.5 45.5 51.0	80 99 70 62	$5.8 \\ 0.0 \\ 2.6 \\ 10.3$	w nw nw sw	52 131 262 148	8 15 32 14	3;30 а	6 Р	2.25 T	
5 6 7 8 9	61 57 43 47 51	4 P 0 A 11 A 2 P 2 P	41 38 33 32 28	10 P 12 P 2 A 12 P	51.0 47.5 38.0 39.5 39.5	78 63 65 70 67	0.0 1.1 1.5 4.6 8.6	re nw nw nw	$\begin{array}{r} 55 \\ 245 \\ 238 \\ 231 \\ 214 \end{array}$	10 25 18 25 18	3:30 A	5 A	.04	
0 1 2 3 4	45 58 60 44 42	3 P 12 P 4 A 0 A 2 P	27 41 44 37 37	3 A 6 A 12 P 12 P 0 A	$ \begin{array}{r} 36.0 \\ 49.5 \\ 52.0 \\ 40.5 \\ 39.5 \end{array} $	80 80 95 96 97	0.4 0.0 0.0 0.0 0.0	nw s s n n	$ \begin{array}{r r} 31 \\ 112 \\ 265 \\ 102 \\ 173 \\ \end{array} $	5 17 28 17 15	0 A 0 A 0 A		T .82 .32	
5 6 7 8	$ \begin{array}{r} 45 \\ 43 \\ 43 \\ 40 \end{array} $	2 P 0 A 2 P 1 P	39 29 28 32	6 A 12 P 6 A 12 P	42.0 36.0 35.5 36.0	$ \begin{array}{c} 92 \\ 77 \\ 76 \\ 65 \end{array} $	$0.0 \\ 0.0 \\ 1.2 \\ 0.2$	n nw sw nw	126 148 158 269	$10 \\ 17 \\ 14 \\ 25$	0 A	8 A	.45	7
$9 \\ 0 \\ 1 \\ 2 \\ 3$	47 54 54 69 49	2 P 3 P 3 P 3 P 11 A	24 36 28 43 39	7 A 0 A 8 A 2 A 12 P	35.5 45.0 41.0 56.0 44.0	74 75 74 76 73	8.0 1.1 4.7 5.3 0.8	s s s	130 250 86 225 170	17			Т	
$\frac{4}{5}$	$\frac{44}{37}$	4 P 0 A	$\frac{37}{20}$	4 A 12 P	$40.5 \\ 28.5 \\ 25.0$		$\begin{array}{c} 0.0 \\ 7.3 \\ 6.5 \end{array}$	se n	119 288 87	$\frac{14}{28}$ $\frac{14}{14}$	5:30 A 10 P	9 A	.04	7 2.5
7 8	$\frac{29}{29}$	2 P 2 P 2 A	$\frac{23}{14}$	4 A 10 P	$\frac{26.0}{21.5}$	$\frac{96}{58}$	$\frac{0.0}{6.6}$	n n nw	$\frac{157}{249}$	$\frac{14}{25}$	0 а	7 A	1.01	8.5
9	$\frac{32}{36}$	12 P 1 P	$\frac{11}{28}$	7 A 11 P	$\frac{21.5}{32.0}$	89 73	$\frac{0.0}{4.4}$	s nw	66 199	$\frac{8}{28}$	11 A 8 P	8 P 12 P	.05	.5 1.5

^{*}Based on least time required to blow one mile.

SOL KRIMINETSKY, Observer

Nov mber 1940	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum		30.87, 1933
Minimum	29.30	28.73, 190-
Mean semi-daily	30.05	
Range	1.29	1.84, 1904; .94, 1905
Air Temperature, in degrees F.	i	, , , ,
Highest	66.0	75, 1938
Lowest	12.9	-4, 1938
Mean	38.9	44.1, 1931; 33.6, 1901
Range	53.0	, , , , ,
Highest mean daily 56, 22nd		
Lowest mean daily 21.5, 28th, 29th		
Mean maximum		
Mean minimum	ì	
Greatest daily range	l	
Least daily range		
Precipitation, in inches	9.11	0.64 1007, 69 1015
Precipitation	3.41	8.64, 1927; .63, 1917
Snow	2.34	13.50, 1938
Maximum precipitation in 24 hrs. 2.25, 2nd		22 1021 2 102
Number of days with .01 or more $\dots 11$	9	22, 1921; 2, 1909
Wind, in miles		
Total movement	4531	5,978, 1906; 2,589, 1889
Greatest daily movement 288–25th		
Least daily movement 31, 10th		
Mean hou ly velocity 6.9	6.3	
Maximum velocity 32, 3rd	30.3	44, 1938
Wind, direction		
Prevailing direction NW	WNW	
North, days7	.,,,,,,	
Northeast, days0		
East, days		
Southeast, days		
South, days		
Southwest, days	1	
West, days		
Northwest, days	l	
Weather	-0.0	
Mean relative humidity, percent77	70.6	
Mean cloudiness, percent67	55.1	72, 1900, 1927; 34, 1917
Number of clear days4	6	15, 1903; 1, 1900, 1911, 1927
Number of fair days	9	16, '12; 4, '89, '30, '32, '35
Number of cloudy days19	15	24, 1927; 9, 1905, 1917
Number hours bright sunshine81	121	182, 1903; 66, 1927
Percent of possible hours of bright		
sunshine	41.3	
First Snow Nov. 16		
	Nov. 6	Oct. 10, '25; Nov. 27, '31

The outstanding feature of November weather was the high rain and snowfall. The total precipitation for the month was 6.31 inches. Of this amount, 2.25 inches fell as rain on the 2nd and 1.30 inches fell as snow on the 26th and 27th. A total of 13 inches of snow fell during the month, the storm on the 26th and 27th giving 11 inches. This heavy snowfall in November has been exceeded only once in the history of the station. A total of 13.50 inches of snow fell in November 1938. On the 24th and 25th of November of that year there was a fall of 11.50 inches. The total precipitation since January 1 is now 40.56 inches, the normal for this period being 40.32 inches.

The mean temperature for the month was 39.4 degrees. The normal mean temperature for November is 38.9 degrees. The highest temperature was 69 degrees on the 22nd and the lowest, 11 degrees on the 29th. The domestic heating load for the month was 769 degree-days, the normal being 783 degree-days. The total heating load for the season is 1520 degree-days, the normal for this period being 1386 degree-days. There were only 81 hours of bright sunshine, whereas the normal for November is 121 hours. The total wind movement for the month was 4986 miles, the normal being 4531 miles. The highest wind velocity was 32 miles per hour on the 3rd.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 624

December 1940

Meteorological Observations

FOR

DECEMBER

1940

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

DAILY RECORDS

		Te	mpera	ture		ty	70		Wind		F	reci	pita	tion	
	Maxi	mum	Min	imum		Iumidi	Hour shine			50					
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	-	Ended	Rain	Snow
$\frac{1}{2}$	44 30	10 A 2 P	22 11	12 P	$\frac{33.0}{20.5}$	81 71	4.0 6.8	nw s	213 139	25 12					Т
3	26	0 A	2	12 P	$\frac{20.5}{14.0}$	60	$\frac{0.3}{9.2}$	nw	259	28					
4	$\frac{1}{23}$	SP	-10	6 A	6.5	87	4.8	nw	97	14		P 12		.18	2.00
4 5	32	4 P	20	2 A 7 A	26.0		0.0	8	57	17	0.4	λ 3	P	.19	2.25
6	26	2 P	7		$\frac{16.5}{30.0}$	67 86	$\frac{3.9}{3.6}$	n	119 181	13 18		ĺ		Т	
7 8	$\frac{46}{42}$	2 P 2 P	$\frac{14}{30}$	7 A 7 A	36.0	82	$\frac{3.0}{2.8}$	s n	112	$\frac{18}{22}$				1	
9	35	2 P	14	12 P	24.5	67	7.8	nw	171	14					
10	43	2 P	13	2 A	-28.0	75	0.4	s	94	22					
11	38	0 A	28	7 A	33.0	64	6.8	\mathbf{n}	201	14	0	10		10	
12	33	12 P	27	7 A	30.0	97	0.0	nw	$\frac{42}{192}$	8 20	$\begin{bmatrix} 6 & 4 \\ 0 & 4 \end{bmatrix}$	$\frac{12}{3}$.16 .08	.75
13 14	$\begin{array}{ c c }\hline 45\\ 34\\ \end{array}$	8 A 2 P	23 12	12 P 12 P	$\frac{34.0}{23.0}$	$\frac{71}{62}$	$\frac{1.8}{7.0}$	nw n	81	6	0 1	1 0	A	.08	
15	32	3 P	7	7 A	$\frac{23.0}{19.5}$	79	6.9	nw	31	$\frac{3}{2}$					
16	33	8 P	18	5 A	25.5	92	0.0	n	48	5	8.	12	P	.87	
17	37	2 P	19	12 P	-28.0	72	0.4	n	153	22					
18	39	3 P	15	8 A	27.0	72	9.0	\mathbf{se}	83						
19	38	2 P	18	7 A	28.0		$\begin{bmatrix} 5.7 \\ 0.0 \end{bmatrix}$	se	58 128	$\begin{array}{ c c } 7 \\ 11 \end{array}$	7	A 8	A	.06	
$\frac{20}{21}$	41	4 P 2 P	31 33	0 A 8 A	$\frac{36.0}{37.0}$	91 87	0.0	se nw	66			A C	A	.00	
$\frac{21}{22}$	38	12 M	$\frac{33}{34}$	5 A	36.0	86	0.0	S	$\frac{60}{62}$					Т	
$\frac{22}{23}$	36	2 A	21	12 P	28.5	60	6.5	n	195						
24	44	3 P	14	7 A	29.0		7.5	\mathbf{s}	69						
25	47	3 P	23	8 A	35.0		9.0	nw	49		_				
26	45	1 P	23	7 A	34.0		3.4	W	44			P 12 A 8		.06	
27	47 39	2 P 2 P	34	0 A 8 A	$\frac{40.5}{37.5}$		$\frac{1.0}{0.0}$	n	69 69			A 8		1.00	
28 29	40	2 P 3 P	$\frac{36}{35}$	8 A 7 A	$\frac{37.5}{37.5}$	96	0.0	n n	57	5		A E		.33	
30	42	11 A	38	8 A	40.0		0.0	SW	104		4			.06	
31	40	2 P	36	7 A	38.0		0.7	nw	307			1			

^{*}Based on least time required to blow one mile.

December 1940	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum	30.66	30.96, 1889
Minimum	29.27	28.85, 1915
Mean semi-daily	30.06	
Range	1.38	1.78, 1895; 1.01, 1892
Air Temperature, in degrees F.	1.00	1.10, 1000, 1.01, 1002
Highest	54.3	65.5, 1908
Lowest	-1.9	-22.5, 1917
	$\frac{-1.5}{27.5}$	
Mean		36.9, 1891; 17.1, 1917
Range	56.2	
Highest mean daily 40.5, 27th	ĺ	
Lowest mean daily 6.5, 4th		
Mean maximum		
Mean minimum	!	
Greatest daily range	ļ	
Least daily range		
Precipitation, in inches	ł	
Precipitation	3.39	7 77 1001 - 97 1906
		7.77, 1901; .87, 1896
Snow	8.50	26.50, '02; t, 1891
Maximum precipitation in 24 hrs. 1.00, 28th	1.0	17 1000 4 1000
Number of days with .01 or more 11	10	17, 1902; 4, 1892
Wind, in miles		
Total movement	4710	6,694, 1925; 3,239, 1918
Greatest daily movement 307, 21st		, ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,
Least daily movement 31, 15th	ļ	
Mean hourly velocity	6.3	
Maximum velocity 28, 3rd, 31st	31.4	48, 1938
	91.4	10, 1300
Wind, direction		
Prevailing direction NW	WNW	
North, days10	i	
Northeast, days0		
East, days0		
Southeast, days		
South, days6		
Southwest, days		
West, days		
Northwest, days		
· ·		
Weather		
Mean relative humidity, percent79.6		
Mean cloudiness, percent62	54.9	71, 1929; 39, 1919
Number of clear days10	8	15, 1890; 2, 1922, 1933
Number of fair days	9	16, '09; 4,'89 '30, '31,'36,'38
Number of cloudy days14	14	23, '33; 7, '09, '23
Number hours bright sunshine109		172, 1896; 63, 1933
Percent of possible hours of bright	1	1
sunshine38.9	45.2	
	10.2	<u> </u>
	1	
	I.	

ANNUAL SUMMARY

Annual 1940	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum	30.81	31.05, 1920
Minimum	28.95	28.41, 1938
Mean semi-daily30.002	30.01	
Range1.57	1.85	2.47, 1938; 1.38, 1933
Air Temperature, in degrees F.		
Highest	95.7	104, 1911
Lowest	-12.2	-26, 1904
Mean	47.4	49.8, 1921; 44, 1904
Range104	107.8	
Highest mean daily 83, July 26		
Lowest mean daily 6.5, Dec. 4		
Mean maximum	57.8	
Mean minimum35.8	36.5	
Greatest daily range		
Least daily range3		
Precipitation, in inches		
Precipitation	43.70	59.00, 1938; 30.68, 1908
Snow54.75	47.78	89.00, 1893; 24.50, 1919
Max. precipitation in 24 hrs 2.58, May 31		
Number of days with .01 or more 122	124	144, 1902; 96, 1924
Wind, in miles		
Total movement	52,223	63,571, 1908; 36,257, 1894
Greatest daily movement572, Feb. 15		
Least daily movement 31, Nov. 10, Dec. 15	- 0	
Mean velocity	5.8	00 1090
Maximum velocity 40, Feb. 14, Apr. 6, May 7	39.5	80, 1938
Wind, direction Prevailing directionWNW	11.	
North, days80	''	
Northeast, days		
East, days		
Southeast, days		
South, days		
Southwest, days		
West, days		
Northwest, days		
Weather		
Mean relative humidity, percent 74.7	67.6	
Mean cloudiness, percent	51.7	60, '98, '01, '02; 41, '08, '24
Number of clear days	116	169, 1936; 59, 1927
Number of fair days	123	182, 1912; 64, 1936
Number of cloudy days121	126	179, '01, '02; 71, 1910
Number hours bright sunshine 2,352	2,353	2838, 1924; 1864, 1902
Percent of possible hours of bright	,	
sunshine 52.8	52.8	
Last snow	April 15	Mar. 14, '10; May 11, '07
First snow Nov. 16	Nov. 6	Oct. 10, '25; Nov. 27, '31
Last snow April 22 First snow Nov 16 Last frost May 13 Limit for Spot 27	May 14	Apr. 23, '04; June 8, '32
First frost	Sept. 21	[Aug. 22, '94, '95; Oct. 13,'09]

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 625

January 1941

Meteorological Observations

FOR

JANUARY

1941

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

DAILY RECORDS

		Te	mpera	ture		Į.	20		Wind			P	recip	oita	tion	•
	Maxi	mum	Min	imum		fumidit	Hours Shine			ø,						
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max, Miles per Hour*	Began		Ended		Rain	Snow
1 2 3 4 5 6 7	39 32 33 35 33 30 26	2 P 1 P 2 P 2 P 12 M 3 P 5 P	27 23 27 25 18 15	12 P 7 A 5 A 10 P 8 A 11 P 8 A	33.0 27.5 30.0 30.0 25.5 22.5 15.5	65 72 95 90 65 63 74	8.6 0.0 0.0 0.0 2.9 9.2 6.3	n nw n nw nw nw	274 137 68 157 210 343 77	22 14 9 14 25 22 9	7 0 3	P A A	12 8 4	P A P	.17 .13 .25	2.00 1.50 4.00
8 9 10 11 12	30 22 37 30 30	3 P 5 P 2 P 3 P 6 P	$\begin{array}{c} 0 \\ 6 \\ 12 \\ 12 \\ 14 \\ \end{array}$	8 A 3 A 7 A 5 A 6 A	$ \begin{array}{r} 15.0 \\ 14.0 \\ 24.5 \\ 21.0 \\ 22.0 \\ \end{array} $	78 87 79 69 77	9.3 5.5 8.2 8.7 7.3	nw nw nw n	28 26 40 203 204	5 3 5 17 25		P P	12	P P	.03	.75 .25
13 14 15 16 17 18	28 15 20 25 35 35	1 P 3 P 2 P 12 P 2 P	5 -2 3 12 23 33	12 P 7 A 2 A 4 A 0 A 8 A	16.5 6.5 11.5 18.5 29.0 34.0	62 64 58 73 95 95	9.4 9.4 6.7 2.8 6.6 0.1	nw nw n n	$\begin{bmatrix} 348 \\ 224 \\ 66 \\ 166 \\ 182 \\ 76 \end{bmatrix}$	25 17 11 12 12 7	3		12 8	P A	.22	1.50 .75
19 20 21 22 23 24	34 22 33 38 42 26	0 A 4 P 3 P 8 P	15 9 4 15 25 13	12 P 7 A 8 A 1 A 12 P	24.5 15.5 18.5 26.5 33.5 19.5	64 63 69 82 74 75	6.5 9.6 9.6 7.2 7.0 3.3	nw nw nw s nw	$ \begin{array}{c c} 209 \\ 326 \\ 101 \\ 60 \\ 216 \\ 170 \end{array} $	10 10 18	3	. 10	12		1.05	T T 12.00
24 25 26 27 28 29	20 31 27 21 32 35	0 A 4 P 3 P 3 P 4 P 3 P	16 16 5 12 11 6	8 A 12 P 7 A 8 A 5 A 12 P	23.5 16.0 16.5 21.5 20.5	83 76 73 63 82	8.3 7.2 4.5 7.2 7.9	n n n n n	228 102 171 105 91	17 17 11 9 28	9	P	12	r	1.03	T
30 31	18 32	5 P 2 P	-9 7	7 A 7 A	4.5 19.5		4.2 7.5	rw nw	129 96		1					Т

^{*}Based on least time required to blow one mile.

January 1941	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum	30.70	31.00, 192
Minimum		28.55, 191
Mean semi-daily	30.07	20.00, 101
Range		2.18, 1913; 0.97, 189
Air Temperature, in degrees F.		100, 1010, 0.01, 100
Highest	51.1	66.0, 193
Lowest9, $30 th$		-26.0, 190
Mean		34.2, 1913; 13.9, 191
Range		,,, 1516, 152
Highest mean daily 34, 18th		
Lowest mean daily		
Mean maximum		
Mean minimum		
Greatest daily range		
Least daily range		
Precipitation, in inches		
Precipitation	3.61	7 15 1909 1 07 190
		7.15, 1898; 1.07, 1890
$egin{array}{lll} ext{Snow} & \dots & 22.75 \ ext{Maximum precipitation in 24 hrs. } 1.05, 24 ext{th} \end{array}$	13.24	33, 1898, 1923; 1.50, '0
		15 202 200 4 100
Number of days with .01 or more 8	111	15, '93, '20; 4, 190
Wind, in miles		
$ \Gamma \text{otal movement} \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots$		7770, 1908; 2896, 189
Greatest daily movement 348, 13th		
Least daily movement 26, 9th		
Mean hourly velocity		
Maximum velocity 28, 19th, 20th, 29th	32.7	47, 193
Wind, direction		
Prevailing direction NNW	WNW	
North, days10		
Northeast, days0		
East, days 0		
Southeast, days0	i i	
South, days \dots 2		
Southwest, days $\dots \dots		
West, days ℓ	1	
Northwest, days	:1	
Weather		
Mean relative humidity, percent 74.6	70.1	
Mean cloudiness, percent		78, 1932; 37, 1903
Number of clear days		17, 1939; 2, 191
Number of fair days		18, 1890, 1926; 3, 1910
Number of cloudy days	13	22, 1923, 1931; 4, 1920, 1939
Number hours bright sunshine 191	137	214, 1920; 74, 193
Percent of possible hours of bright		211, 1020, 11, 100
sunshine	46.7	
	10	
	1	l .

The weather during January was colder than usual and had a heavy fall of snow. The mean temperature for the month was 21.2 degrees which is 3 degrees lower than usual. However, it was 2.7 degrees warmer than January 1940. The lowest temperature was -9 degrees on the 30th and the highest was 42 degrees on the 23rd. The domestic heating load for the month was 1358 degree-days compared to a normal of 1265 degree-days. The total heating load for the season is 3982 degree-days, the normal for this period being 3814 degree-days.

The total precipitation for the month was 2.21 inches, the normal being 3.61 inches. The total snowfall was 22.75 inches of which 12 inches fell on January 24. The normal snowfall for January is 13.32 inches. The total snowfall for the winter is now 40.75 inches, the normal for this period being 24.16 inches. There were 191 hours of bright sunshine compared to a normal of 137 hours. The highest wind velocity was 28 miles per hour.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 626

February 1941

Meteorological Observations

FOR

FEBRUARY

1941

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

DAILY RECORDS

		Te	mpera	ture		Ţ.	70		Wind		1	Precipit	ation	
	Maxi	mum	Min	imum		umidi	Hour			20				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles pcr Hour*	Began	Ended	Rain	Snow
1 2 3 4 5	38 29 30 34 40	3 P 3 P 2 P 2 P 3 P	5 21 18 17 2	8 A 0 A 8 A 12 P 6 A	$21.5 \\ 25.0 \\ 24.0 \\ 25.5 \\ 21.0$	76 93 75 70 74	$9.8 \\ 0.1 \\ 7.8 \\ 7.2 \\ 10.1$	nw nw n n	71 30 207 286 59	$\begin{bmatrix} 8 \\ 4 \\ 20 \\ 25 \\ 7 \end{bmatrix}$	8.	8 Р	.03	1.00
6 7 8 9 10	36 46 44 35 28	2 P 9 P 0 A 1 P 3 P	$\begin{array}{c} 2 \\ 27 \\ 31 \\ 12 \\ 4 \end{array}$	7 A 4 A 7 A 12 P 8 A	19.0 36.5 37.5 23.5 16.0	81 93 61 60 60	8.2 0.0 8.5 8.7 10.3	nw nw w sw n	127 249 287 108	22 11	8 2	11 Р	1.39	
11 12 13 14 15 16	40 38 39 39 38 28	3 P 4 P 3 P 3 P 2 P	$ \begin{array}{r} 9 \\ 15 \\ 19 \\ 32 \\ 24 \\ 15 \end{array} $	7 A 6 A 4 A 6 A 12 P 12 P	24.5 26.5 29.0 35.5 31.0 21.5	75 80 85 94 74 54	10.4 10.4 4.3 0.0 5.3 10.6	w n s ³ n nw	$ \begin{array}{r r} 49 \\ 74 \\ 57 \\ 77 \\ 226 \\ 382 \\ \end{array} $		0 .	6 а	T .11	Т
17 18 19 20 21	30 23 26 31 35	2 P 1 P 4 P 1 P 5 P	12 17 18 19 24	6 A 7 A 7 A 6 A 3 A	$\begin{array}{c} 21.0 \\ 22.5 \\ 22.0 \\ 25.0 \\ 29.5 \end{array}$	72 66 65 67 68	8.4 8.8 7.5 7.3 7.1	nw nw nw nw	99 192 328 367 308	12 25 28 28 28	8 1	12 Р	.02	.25
22 23 24 25 26 27	37 37 32 41 31 39	2 P 2 P 2 P 3 P 4 P	25 25 19 13 17 12	12 P 12 P 7 A 6 A 7 A 6 A	31.0 31.0 25.5 27.0 24.0 25.5	69 65 60 67 58 59	$\begin{array}{c} 6.6 \\ 8.1 \\ 10.9 \\ 8.0 \\ 11.0 \\ 11.1 \end{array}$		178 299 303 69 341 127	22 25 25 28				Т
28	26	4 P 3 P	14	6 A 7 A	20.0 20.0	69	5.5	n n	357		8 1	12 г	.04	1.00

^{*}Based on least time required to blow one mile.

30.63 29.24 30.04 1.38	
$ \begin{array}{c c} 29.24 \\ 30.04 \end{array} $	
$ \begin{array}{c c} 29.24 \\ 30.04 \end{array} $	
30.04	31.05, 192
	28.56, 1898
1.38	1 00 100 100 00 140 10
1.00	1.89, '00, '08; .88, '13, '3
50.5	CT 0 109
50.5 -7.4	65.0, 193
$\frac{-7.4}{23.7}$	-22.5, 191
58	
96	
ŀ	
$_{32.8}$	
14.5	
1.10	
i	
ł	
3.19	8.12, 1900; 0.62, 190
13.99	48.75, 1893; 0.50, 193
	101.0, 1000, 0100, 100
10	15, '93, '20; 4, 190
	, , , ,
4776	6445, 1896; 3438, 1893
	,,,
7.1	
31.7	48, 1934, 1937
VNW	
1	
i	
l	
1	
ccc	
$\frac{66.6}{50.9}$	66 200 227 21 100
10	66, '90, '27; 31, 1905 17, '19, '24; 2, 192'
8	16, 1920; 2, 1930
- 1	16, 1894; 2, 1930
	221, 1924; 110, '27, '3
100	, 10_1, 110, 21, 00
53.4	
1	
5	10 158

The weather during February was marked with less rain and snow than usual, a large amount of sunshine and higher mean temperature than usual. The total precipitation for the month was 1.59 inches compared to a normal of 3.19 inches. The snowfall was 2.25 inches, while normally we get 13.99 inches of snow in February. This is the least snowfall for February on record here at the station with the exception of 1937 when one-half inch fell during the month. There were 212 hours of sunshine, the normal being 158 hours. This has been exceeded only three times at the station, viz., 1919, 1920 and 1924. It will be noted that two of these years were leap years with 29 days in February.

The mean temperature for the month was 25.8 degrees. The normal mean temperature for February is 23.7 degrees. The domestic heating load for the month was 1099 degree-days, the normal being 1156 degree-days. The total heating load for the current season is 5081 degree-days. The normal to this date is 4970 degree-days. The average wind velocity for the month was 7.9 miles per hour, the normal being 7.1 miles per hour. The highest velocity was 40 miles per hour on the 15th.

Meteorological Series

Bulletin No. 627

March 1941

Meteorological Observations

FOR

MARCH

1941

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Te	mpera	ture		ž.			Wind		P	recipitat	ion	
	Maxi	imum	Min	imum		umidi	Hourshine			×.				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Rain	Snow
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 6 17 18 19 20 21 22 23 4 22 5 6 27	38 40 41 32 35 31 37 32 33 41 43 47 49 40 26 34 42 43 44 44 48	3 P 3 P 3 P 3 P 4 P 4 P 2 P 2 P 2 P 4 P 2 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 4	22 27 31 18 17 20 14 25 25 19 17 21 21 21 21 21 21 21 21 21 21 21 21 21	0 A 7 A 3 A 12 P 2 A 7 A 7 A 8 A 7 A 7 A 7 A 7 A 7 A 7 A 7 A 7 A 7 A 7	30.0 33.5 37.5 29.5 24.5 27.5 28.0 28.0 24.5 27.0 26.0 30.0 36.0 27.5 27.5 28.0 30.0 36.0 36.0 36.0 36.0 36.0 36.0 36	66 66 91 71 58 66 61 61 65 61 83 83 53	6.3 11.2 11.0 2.2 11.3 11.3 9.5 0.0 0.0 11.6 0.0 8.5 11.7 11.8 11.5 1.3 6.8 9.9 9.7 11.0 12.2 12.2 2.5 2.5 11.3	n nw s n nw nw n n n n n n n n n n n n n	395 266 134 269 312 203 102 176 159 123 159 67 309 454 549 304 215 69 37 125 38 38 38 38 38 38 38 38	22 11 25 25 20 8 22 10 15 15 18 7 13 9 8 32 48 40 9 9 18 22 22 22 22 10 9 18 22 10 10 10 10 10 10 10 10 10 10 10 10 10	8 A	4 р 12 р 12 р	.02 T .67 .09 .65	.50 T 6.000 2.000 7.000 T
28 29 30 31	51 49 35 48	4 P 1 P 3 P 4 P	25 33 25 24	6 A 3 A 8 P 6 A	38.0 41.0 30.0 36.0	59 54 61 48	10.8 8.4 12.5 12.4	nw n nw nw	124 152 296 257	$\begin{vmatrix} 20 \\ 25 \end{vmatrix}$				

^{*}Based on least time required to blow one mile.

Barometer, in inches (Readings reduced to sea level) Maximum 30.47, 7th Minimum 29.36, 4th Mean semi-daily 29.917 Range 1.11 Air Temperature, in degrees F. Highest 54, 23rd Lowest 6, 18th Mean 30.3 Range 48 Highest mean daily 41, 29th Lowest mean daily 11, 18th Mean maximum 39.0 Mean minimum 21.6 Greatest daily range 34, 14th, 15th Least daily range 6, 8th, 9th, 16th Precipitation, in inches	29.24 30.00 1.34 63.8 6.1 34.4 57.6 43.2 25.4	30.96, 190- 28.47, 191- 2.10, 1914; 0.85, 1913 79.5, 1907, 192- -7.5, 1906
Maximum 30.47, 7th Minimum 29.36, 4th Mean semi-daily 29.917 Range 1.11 Air Temperature, in degrees F. Highest 54, 23rd Lowest 6, 18th Mean 30.3 Range 48 Highest mean daily 41, 29th Lowest mean daily 11, 18th Mean maximum 39.0 Mean minimum 21.6 Greatest daily range 34, 14th, 15th Least daily range 6, 8th, 9th, 16th	29.24 30.00 1.34 63.8 6.1 34.4 57.6 43.2 25.4	28.47, 191- 2.10, 1914; 0.85, 191- 79.5, 1907, 192
Minimum 29.36, 4th Mean semi-daily 29.917 Range 1.11 Air Temperature, in degrees F. 54, 23rd Lighest 54, 23rd Lowest 6, 18th Mean 30.3 Range 48 Highest mean daily 41, 29th Lowest mean daily 11, 18th Mean maximum 39.0 Mean minimum 21.6 Greatest daily range 34, 14th, 15th Least daily range 6, 8th, 9th, 16th	29.24 30.00 1.34 63.8 6.1 34.4 57.6 43.2 25.4	28.47, 191- 2.10, 1914; 0.85, 191- 79.5, 1907, 192
Mean semi-daily 29.917 Range 1.11 Air Temperature, in degrees F. Highest 54, 23rd Lowest 6, 18th Mean 30.3 Range 48 Highest mean daily 41, 29th Lowest mean daily 11, 18th Mean maximum 39.0 Mean minimum 21.6 Greatest daily range 34, 14th, 15th Least daily range 6, 8th, 9th, 16th	30.00 1.34 63.8 6.1 34.4 57.6 43.2 25.4	2.10, 1914; 0.85, 191 79.5, 1907, 192
Range 1.11 Air Temperature, in degrees F. 11ghest 54, 23rd Lighest 6, 18th 30.3 Lowest 6, 18th 41ghest Lighest mean daily 41, 29th 41, 29th Lowest mean daily 11, 18th 41, 18th Lean maximum 39.0 39.0 Mean minimum 21.6 31, 14th, 15th Breatest daily range 31, 14th, 15th 41, 15th Least daily range 6, 8th, 9th, 16th	1.34 63.8 6.1 34.4 57.6 43.2 25.4	79.5, 1907, 192
Air Temperature, in degrees F. lighest 54, 23rd lowest 6, 18th dean 30.3 Range 48 lighest mean daily 41, 29th lowest mean daily 11, 18th dean maximum 39.0 dean minimum 21.6 dreatest daily range 34, 14th, 15th least daily range 6, 8th, 9th, 16th	63.8 6.1 34.4 57.6 43.2 25.4	79.5, 1907, 192
Highest 54, 23rd Lowest 6, 18th Jean 30.3 Range 48 Highest mean daily 41, 29th Lowest mean daily 11, 18th Jean maximum 39.0 Jean minimum 21.6 Greatest daily range 34, 14th, 15th Least daily range 6, 8th, 9th, 16th	6.1 34.4 57.6 43.2 25.4	
.cowest 6, 18th .dean 30.3 .tange 48 lighest mean daily 41, 29th .cowest mean daily 11, 18th .dean maximum 39.0 .dean minimum 21.6 .dreatest daily range 34, 14th, 15th .dreatest daily range 6, 8th, 9th, 16th	6.1 34.4 57.6 43.2 25.4	
Mean 30.3 Range 48 Highest mean daily 41, 29th Jowest mean daily 11, 18th Jean maximum 39.0 Mean minimum 21.6 Greatest daily range 34, 14th, 15th Least daily range 6, 8th, 9th, 16th	34.4 57.6 43.2 25.4	-7.5, 190
Cange 48 Lighest mean daily 41, 29th Lowest mean daily 11, 18th Lean maximum 39.0 Lean minimum 21.6 Greatest daily range 34, 14th, 15th Least daily range 6, 8th, 9th, 16th	57.6 43.2 25.4	
Highest mean daily 41, 29th Lowest mean daily 11, 18th Hean maximum 39.0 Hean minimum 21.6 Hear minimum 34, 14th, 15th Hear daily range 6, 8th, 9th, 16th Hear daily range 6, 8th, 9th, 16th	43.2 25.4	
wwest mean daily 11, 18th Iean maximum 39.0 fean minimum 21.6 freatest daily range 34, 14th, 15th deast daily range 6, 8th, 9th, 16th	43.2 25.4	
Jean maximum 39.0 Jean minimum 21.6 Greatest daily range 34, 14th, 15th Least daily range 6, 8th, 9th, 16th	43.2 25.4	
Mean minimum 21.6 Greatest daily range 34, 14th, 15th Least daily range 6, 8th, 9th, 16th	25.4	
Greatest daily range 34, 14th, 15th Least daily range 6, 8th, 9th, 16th		
east daily range 6, 8th, 9th, 16th	ı.	
Precipitation, in inches	0.50	
	1 0 =0	1
$\operatorname{Precipitation} \ldots 1.63$	3.70	7.13, 1899; 0.12, 191
now	7.47	27, 1899; 0, 192
Iaximum precipitation in 24 hrs		·
Number of days with .01 or more 6	11	17, 1890, 1913, 1936; 3, 191.
Wind, in miles		
otal movement	5944	8182, 1896; 3006, 190
Greatest daily movement 549, 19th		0102, 1777, 77007. 1877
east daily movement		
Ican Lourly velocity 8.7		
Jaximum velocity 48, 18th	31.2	48, 1932, 1939
Wind, direction	1 3	
revailing direction	WNW	
North, days		
Vortheast, days		
last, days		
outheast, days		
outh, days2		
outhwest, days		
Vest, days		
Vorthwest, days		
Weather	610	
Jean relative humidity, percent 68 2		
Jean cloudiness, percent		99 1091, 9 10
Number of clear days		22 1924; 3 19
Number of fair days	10	17, 1900; 4, 1895, 1897, 1936
Number of cloudy days	10	21, 1901; 1, 191;
Number hours bright sunshine256	199	292, 1924; 93, 190
Percent of possible hours of bright	-9.0	
sunshine 69	53 6	

The weather during March was clear, cold and windy. The mean temperature was 30.3 degrees, the normal mean being 34.4 degrees. The highest temperature was 54 degrees on the 23rd and the lowest, 6 degrees on the 18th. The domestic heating load was 1074 degree-days compared to a normal of 949 degree-days. The total heating load for the season is now 6155 degree-days, the normal for this period being 5919 degree-days. There were 256 hours of bright sunshine, the normal being 199 hours. The total precipitation was 1.63 inches where as the normal precipitation for March is 3.70 inches. The snowfall was 15.50 inches. Normal snowfall for March is 7.47 inches. The total snowfall for the winter is now 58.50 inches, the normal for this period being 45.62 inches. The total wind movement for the month was 6449 miles, which is well above the average for March. The highest velocity was 48 miles per hour on the 18th.

Meteorological Series

Bulletin No. 628

April 1941

Meteorological Observations

FOR

APRIL

1941

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Te	mpera	ture		y.			Wind		Pr	ecipitat	ion	
	Maxi	mum	Mini	imum		Iumidit	f Hours nshine		واد	20 10 10				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	per Hour*	Began	Ended	Rain	Snow
1 2 3 4 5 6 7 8 9 10 11 12	47 49 57 60 56 55 62 65 66 65 67	1 P 2 P 4 P 3 P 3 P 3 P 2 P 4 P 2 P 3 P	28 33 31 29 29 34 30 28 32 31 31	5 A 6 A 5 A 5 A 6 A 12 P 6 A 5 A 4 A 6 A 5 A	37.5 41.0 44.0 44.5 39.5 45.0 42.5 46.0 46.5 49.0 49.0	34 39 51 46 48	3.3 6.0 12.8 12.8 4.0 10.7 13.0 13.0 13.1 13.2 13.2	n n n n nw nw ne n n se n se s	275 246 88 75 249 213 221 64 108 113	12 20 22 10 8 25 25 18 9 17 13 12	1 р	12 р	.14	
13 14 15 16 17 18	64 75 86 74 74 76	1 P 4 P 2 P 4 P 3 P 5 P	34 45 53 53 49 48	6 A 6 A 12 P 8 A 4 A	49.0 60.0 69.5 63.5 61.5 62.0	56 76 56 57 83 54	10.4 8.2 12.0 13.4 7.0 12.5	se nw nw nw s	201 172 131 181 198 162	18 18 25 18 25 17	6 A	9 A	.11	
19 20 21 22 23 24 25	83 90 67 56 61 57 62	3 P 2 P 3 P 3 P 2 P 1 P	52 53 50 37 28 38 32	6 A 5 A 11 P 5 A 5 A 12 P 6 A	67.5 71.5 58.5 46.5 44.5 47.5	73 57 52 50 74	11.1 10.3 13.6 13.7 12.1 7.5 12.6	s nw nw s nw	187 305 290 155 175 145	14 25 30 20 17 20 13	1 A 11 P 0 A	12 Р	.04	
26 27 28 29 30	56 60 77 81 82	4 P 4 P 5 P 3 P 2 P	34 37	4 A 1 A 6 A 6 A 2 A	45.0 48.5 64.5 59.0 65.0	56 39 35	11.3 6.7 12.1 14.0 12.4	ne e n se nw	313 156	12 28 25 22 20			Т	

^{*}Based on least time required to blow one mile.

		1
April 1941	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum 30.43, 12th, 23rd	30.48	30.71, 1911
Minimum		28.99, 1907, 1929
Mean semi-daily	29.99	
Range	1.11	1.52, 1930; .72, 1919
Air Temperature, in degrees F.	70.4	00 # 1000
Highest	79.4	88.5, 1896
Mean	$\frac{22.0}{45.7}$	8.5, 1923 52, 1921; 41.2, 1926
Range	$\frac{43.7}{57.4}$	52, 1921; 41.2, 1926
Highest mean daily	37.4	
Lowest mean daily		
Mean maximum	56.8	
Mean minimum	34.8	
Greatest daily range 44, 29th		
Least daily range 16, 2nd		
Precipitation, in inches		
Precipitation	3.35	6.89, 1929; .76, 1892
Snow	2.17	11, 1891; 0, 1910, 1934
Maximum precipitation in 24 hrs.		
.26, 23 rd, 24 th		
Number of days with .01 or more 5	11	18, 1909; 3, 1892
Wind, in miles		
Total movement		8208, 1908; 3853, 1917
Greatest daily movement 313, 28th		
Least daily movement 64, 9th		
Mean hourly velocity	7.5	40 1025 1020 1040
Maximum velocity 30, 21st	31.4	40, 1935, 1938, 1940
Wind, direction	33737337	
Prevailing direction	WNW	
Northeast, days		
East, days		
Southeast, days		
South, days 6		
Southwest, days0		
West, days		
Northwest, days10		
Weather		
Mean relative humidity, percent 57.3	61.6	
Mean cloudiness, percent35	51.8	75, 1901; 34, 1925, 1927
Number of clear days23	11	19, 1911; 3, 1898, 1902
Number of fair days6	9	18, 1915; 2, 1894, 1901
Number of cloudy days	10	22, 1901; 3, 1899, 1924
Number hours bright sunshine 329	220	296, 1911; 103, 1901
Percent of possible hours of bright sunshine	54.7	
sunshine		Mar 14 1910 May 11 1007
2000 Maria M	1.151.10	1

April weather was unusual in many respects, being dry, warm and sunny. The rainfall during the month was only .55 of an inch. This is the least rainfall in Amherst for April since 1836. Other dry Aprils were in 1844 and 1892 when .57 and .76 of an inch respectively were recorded. The normal rainfall for April is 3.35 inches. The total rainfall since January 1 is 5.98 inches while the normal for this period is 13.85 inches. The most serious results from the dry spell have been the disastrous forest fires throughout the state.

The mean temperature for the month was 52.1 degrees compared to a normal of 45.7 degrees. This is the highest mean April temperature in Amherst since 1837 except for the year 1839 when a temperature of 52.2 degrees was recorded. The week beginning April 14 was the most unusual period. A mean temperature of 65 degrees was recorded during that week. This has never been equalled during April since 1889 and has been exceeded only three times during the month of May. The highest temperature recorded was 90 degrees on April 20. The heating load during the month was 399 degree-days, the normal being 579 degree-days. The total heating load during the season is now 6554 degree-days, compared to a normal of 6498 degree-days for this period. There were 329 hours of bright sunshine, which is the highest number of hours

recorded in April. The normal is 220 hours.

Dr. J. K. Shaw, Research Professor of Pomology, makes the following comments: "Most fruit plants came through the winter in good shape except raspberry canes which were badly killed back, perhaps by the severe cold of December 10. Peach buds survived the winter without excessive killing. March was cold with subnormal precipitation but in early April there came an excessively warm period which caused rapid development of foliage and flower Colder weather followed, causing some injury to flower buds of apples, killing many pistils. The extent of this injury is not yet known but apple orchards in low frosty locations have probably suffered severe damage. No serious injury to peach, pear, and plum buds has been observed. Rarely has vegetation been so far advanced at the end of April. There remains a danger period of two or three weeks during which damage from cold is very possible. April was very dry. Fruit plants have not suffered yet but if heavy rains do not come in May, growth will be severely checked."

Meteorological Series

Bulletin No. 629

May 1941

Meteorological Observations

FOR

MAY

1941

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5'' N. Longitude, 72° 31′ 10''W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Te	mpera	ture		y.			Wind		P	recipi	itat	ion
	Maxi	mum	Min	imum		umidi	Hourshine			m				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended		Rain
$ \begin{array}{c} 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \end{array} $	58 56 61 72 66 79	5 P 1 P 5 P 4 P 5 P	44 32 31 30 45 35	2 A 12 P 5 A 5 A 12 P 5 A	51.0 44.0 46.0 51.0 55.5 57.0	90 80 67 49 62 52	5.8 2.5 14.1 12.6 6.9 14.3	ne n n nw nw sw	55 153 204 124 130 68	10 20 18 18 14 9	6 а	11	P	.17 T
7 8 9 10 11 12	79 74 66 65 64 64 68	2 P 5 P 1 P 12 M 4 P 1 P 4 P	56 58 52 42 36 33 32	6 A 6 A 8 P 12 P 5 A 5 A 5 A	67.5 66.0 59.0 53.5 50.0 48.5 50.0	67 53 58 56	10.3 5.8 3.3 9.7 14.3 12.5 12.9	s n n nw sw sw	191 109 110 261 101 68 48	18 9 13 20 15 11 8	3 A 2-30 F 0 A	12	A P A	.31 1.28 .27
14 15 16 17 18 19 20	69 75 82 78 65 75 83	2 P 4 P 12 M 4 P 2 P 5 P	36 40 39 52 45 38 40	5 A 5 A 4 A 12 P 5 A 5 A	52.5 57.5 60.5 65.0 55.0 56.5 61.5	76 48 48 55	12.0 14.6 14.6 8.0 12.1 14.7 14.8	nw nw s s nw nw	86 128 53 131 359 172 70	11 18 7 22 22 25 14 10	1 F	11	P	.05
21 22 23 24 25	86 89 87 64 66	3 P 3 P 2 P 9 A 5 P	48 55 59 40 37	4 A 5 A 12 P 12 P 6 A	67.0 72.0 73.0 52.0 51.5	72 61 79 51	12.5 12.8 12.3 7.5 12.0	sw nw n n	116 90 115 246 246	14 20 25 28	8 F 12 M		P P	.06 .24
26 27 28 29 30 31	79 81 79 72 69 75	3 P 4 P 4 P 3 P 3 P	55 53 50 38	3 A 4 A 12 P 6 A 4 A	66.0 68.0 66.0 61.0 53.5 56.0	73 90 61 54	11.5 7.0 8.5 10.1 15.0 13.0	w nw s n n	155 158 69 251 152 31	$ \begin{array}{c c} 18 \\ 8 \\ 25 \\ 22 \end{array} $	7 4	9	P	T .49

^{*}Based on least time required to blow one mile.

	1	1
May 1941	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum 30.18, 20th, 21st	30.39	30.62, 1936
Minimum	29.51	29.10, 1938
Mean semi-daily	29.96	20.10, 1000
Range	0.88	
Air Temperature, in degrees F.	0.00	
	86.1	04.5 1906 1011
Highest	31.2	94.5, 1896, 1911
Lowest		24.0, 1900
Mean	57.1	
Range	54.9	
Highest mean daily 73, 23rd		
Lowest mean daily	,	
Mean maximum		
Mean minimum		
Greatest daily range 44, 6th		
Least daily range		
Precipitation, in inches		
Precipitation 2.87	3.60	7.44, 1931; .48, 1903
Maximum precipitation in 24 hrs.	0.00	1.11, 1001, .10, 1000
1.55 9th, 10th		
Number of days with .01 or more 8	12	20, 1901; 5, 1903
	12	20, 1301, 5, 1306
Wind, in miles	4.704	7046 1007 0100 1004
Total movement	4504	5946, 1907; 2180, 1894
Greatest daily movement 359 18th		
Least daily movement 31, 31st		
Mean hourly velocity	6.1	
Maximum velocity 28, 25th	28.8	45, 1935
Wind, direction		
Prevailing direction NW	W	
North, days		
Northeast, days1		
East, days0		
Southeast, days0		
South, days		
Southwest, days		
West, days		
Northwest, days		
· •		
Weather	00.5	
Mean relative humidity, percent 63.9	60.7	1001 1000 80 1008
Mean cloudiness, percent46	52.1	70, 1901, 1902; 30, 1923
Number of clear days20	11	20, 1923; 0, 1927
Number of fair days9	11	17, 1907, 1926; 5, 1923
Number of cloudy days	9	20, 1927; 3, 1903, 1922
Number hours bright sunshine338	252	334, 1922; 137, 1927
Percent of possible hours of bright		
sunshine	55.6	
Thunder and lightning		
Last frost	May 14	June 8, 1932; Apr. 23, 1904
	,	, , , , , , , , , , , , , , , , , , , ,

The weather during May continued dry but in other respects was quite normal. The rainfall during the month was 2.87 inches, the normal for May being 3.60 inches. There was only one good rain during the month, on the 8th, 9th and 10th, hence the drouth has continued to the end of the month. The total rainfall since January 1 is now 8.85 inches, while the normal for this period is 17.45 inches.

The mean temperature for the month was 57.9 degrees which is 0.8 of a degree higher than usual. The highest temperature was 89 degrees on the 22nd and the lowest, 30 degrees on the 4th. A temperature of 32 degrees on the 13th brought the last frost of the month near the observatory, although there were later frosts in the low part of the town. The domestic heating load during the month was 247 degree-days. The normal for May is 287 degree-days. The total heating load for the season is now 6801 degree-days, while the normal for the period is 6785 degree-days. There were 338 hours of bright sunshine which is the largest number of hours ever recorded in May. No high winds were recorded.

Dr. J. K. Shaw, research professor of Pomology, makes the following report: "The month of May was characterized by a continued deficiency of rainfall. This has not as vet caused any apparent injury to fruit trees which are still making good growth, but there will be a short hay crop even if there should be ample rain soon. It also sets the stage for more severe damage should rainfall continue to be low. Fruit trees bloomed ten days or more earlier than normal. Frosts killed a few apple blossoms but not enough to interfere with the crop unless it may be in orchards in frosty locations. Considerable cool weather has delayed the progress of vegetation somewhat but it continues to be more advanced than normal. There were few wet periods favorable to apply scab infection, and scab control should be easier than last year. Strawberries are beginning to ripen a week or more earlier than normal. We may expect a light raspberry crop because of cold injury which probably was caused by the cold of December 4 and not on the 10th as stated last month."

Meteorological Series

Bulletin No. 630

June 1941

Meteorological Observations

FOR

JUNE

1941

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		\mathbf{T} e	mpera	ture		S,			Wind		Pi	recipitat	ion	
		mum		imum		Mean Relative Humidity	Number of Hours Bright Sunshine	lling tion	ment	Max. Miles per Hour*		T		
Day	Degrees	Time	Degrees	Time	Mean	Mean Relati	Numl Bright	Prevailing Direction	Total Movement	Max. per H	Began	Ended	Rain	Snow
1 2 3 4 5 6 7 8	62 78 82 68 58 76 84 88	2 P 5 P 3 P 12 M 6 P 5 P 4 P 1 P	55 55 47 52 55 52 47 61	9 A 5 A 5 A 1 A 5 A 5 A 12 P	58.5 66.5 64.5 60.0 56.5 64.0 65.5 74.5	92 70 65 78 89 65 69 67	$\begin{array}{c} 2.3 \\ 11.5 \\ 15.1 \\ 5.7 \\ 0.0 \\ 13.8 \\ 12.2 \\ 12.7 \end{array}$	se n w se n n	106 81 44 65 176 162 71 200	10 8 7 12 17 17 6 28	7 A 5 P 0 A	11:30 _A	.12 .25 .87	
9 10 11 12 13 14 15 16 17 18	72 72 70 72 60 71 78 71 82 78 88	1 P P 2 P P 2 P P 1 P P 1 P P P P P P P	53 50 45 44 53 53 57 58 53 55 53	5 A 4 A 5 A 6 A 0 A 5 A 12 P 5 A 12 P 5 A	62.5 61.0 57.5 58.0 56.5 62.0 67.5 64.5 70.5	57 48 64 65 88 94 80 89	15.2 14.5 15.2 12.0 0.0 2.4 11.0 6.0 11.4 7.8 15.3	nw nw n s s s s s s m w nw	443 312 227 89 90 38 100 52 50 74 69	32 32 25 11 9 7 18 9 8 12 14	6 P 4 A 0 A 0 A 0 A 7 P	12 P 12 P	.01 .40 .37 1.84 .27 .20	
20 21 22 23 24 25 26	93 91 93 67 83 76	3 P 1 P 4 P 6 P 4 P 4 P 3 P	63 62 51 48 55 54	5 A 5 A 4 A 12 P 5 A 6 A	78.0 77.0 77.5 59.0 65.5 65.5 71.0	60 61 58 80 68 50 67	15.3 15.3 15.3 1.4 13.0 15.3 15.3	sw w nw ne w nw	66 60 114 120 95 145 164	11 5 14 13 12 15 17	2 A	12 м	.80	
26 27 28 29 30	88 94 93 91 89	3 P 3 P 2 P 4 P 2 P	65 68 70 65	5 A 4 A 5 A 12 P 5 A	71.0 79.5 80.5 80.5 77.0	61 68 70 71	15.3 14.0 11.3 12.4	se se se n	176 176 195 147 108	13	7 Р	8 P	T 1.00	

^{*}Based on least time required to blow one mile.

June 1941	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum 30.21, 12th, 13th	30.30	30.54, 1889
Minimum	29.55	29.24, 1902
Mean semi-daily	29.95	
Range	0.75	
Air Temperature, in degrees F.		
Highest	91.2	101, 1919
Lowest	40.0	34, 1891
Mean	65.7	
Range	51.3	
Highest mean daily 80.5, 28th, 29th		
Lowest mean daily $\dots 56.5$, 5 th, 13 th		
Mean maximum		
Mean minimum		
Greatest daily range		
Least daily range		
Precipitation, in inches		
Precipitation	3.75	9.68, 1922; 0.76, 1908
Maximum precipitation in 24 hrs.		
2.11, 14th, 15th		15 1000 4 1000
Number of days with .01 or more11	11	17, 1922; 4, 1908
Wind, in miles		
Total movement	3585	4571, 1908; 1409, 1906
Greatest daily movement 443, 9th		
Least daily movement 38, 14th		
Mean hourly velocity	5.0	40, 1000
Maximum velocity 32, 9th, 10th	24.6	48, 1939
Wind, direction	****	
Prevailing direction	WSW	
North, days5		
Northeast, days1		
East, days0		
Southeast, days		
South, days		
Southwest, days		
West, days		
Northwest, days 6		
Weather	44.0	
Mean relative humidity, percent 70.7	66.9	E1 1009 00 1000
Mean cloudiness, percent57	51.1	71, 1903; 28, 1908
Number of clear days	10	22, 1908; 1, 1927
TNHINDER OF 1917 days 3	12	23, 1912; 4, 1895, 1903
Number of clouds do-se	8	22, 1903; 1, 1923 362, 1908; 102, 1903
Number of cloudy days 5	95=	
Number of cloudy days	257	502, 1503, 102, 1505
Number of cloudy days 5 Number hours bright sunshine		502, 1503, 102, 1505
Number of cloudy days 5 Number hours bright sunshine	257 54.0	502, 1908, 102, 1906
Number of cloudy days 5 Number hours bright sunshine		502, 1503, 102, 1505

June weather was warmer than usual, with more precipitation and sunshine than normal.

The mean temperature for the month was 67.2 degrees, compared to the normal for June of 65.7 degrees. The highest temperature was 94 on the 27th and the lowest, 44 on the 12th. Two warm spells occurred during the month, the first from the 19th to the 23rd and the second from the 26th to the end of the month. Most of the daily maximum temperatures during these warm spells ranged in the 90's. While these periods were very warm, they were not as prolonged and as humid as the 8-day heat wave of June, 1938, and the 10-day period in 1933. The domestic heating load for the month was 60 degree-days. The normal for June is 69 degree-days. The total load for the heating season was 6861 degree-days, while the normal for the period is 6854 degree-days.

The total rainfall during the month was 6.13 inches which reduced to some extent the large deficiency in rainfall which has prevailed since the first of the year. The normal for June is 3.75 inches. The total rainfall since January 1 is now 14.98 inches, while the normal for this period is 21.20 inches. There were 328 hours of sunshine. This record has been exceeded only twice in June—360 hours in 1923 and 362 in 1908. The normal for June is 257 hours. The total wind movement for June was slightly higher than normal, although the four-day wind travel from the 8th to the 11th was unusually high and steady.

Dr. J. K. Shaw, research pomologist, gives the following report on the progress of vegetation: "A few timely rains maintained soil moisture and increased the hay crop above the prospects in early June. Yet there was not enough continued wet weather for extensive apple scab development, so that control was much easier than last year. Apples set well and the McIntosh crop promises to be good, but Baldwins are reported a light crop due to light bloom. The strawberry harvest is about over and the crop was average. The progress of vegetation during June was not far from normal."

Meteorological Series

Bulletin No. 631

July, 1941

Meteorological Observations

FOR

JULY

1941

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Te	mpera	iture		Į.	20	,	Wind		Pr	ecipitat	ion	
	Maxi	mum	Min	imum		lumidi	Hours			go.				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Rain	Snow
1 2 3 4 5	91 93 79 69 83	2 P 1 P 2 P 6 P 4 P	69 69 59 61 59	6 A 12 P 5 A 5 A 6 A	80.0 81.0 69.0 65.0 71.0	61 84 69	5.0 14.4 11.0 1.0 11.5	s nw s sw	112 149 103 69 71	18 18 8 6 8	3 P 1 P 10 P 0 A	5 P 2 P 12 P 8 A	.02 T T .04	
6 7 8 9	83 82 85 76 86	12 M 3 P 1 P 3 P 4 P	57 63 66 61 57	6 A 0 A 12 P 5 A 5 A	70.0 72.5 75.5 68.5 71.5	67 82 76 73 52	15.2 4.5 9.6 7.3 15.1	se s s nw sw	146 194 195 135 102	$ \begin{array}{r} 12 \\ 17 \\ 15 \\ 20 \\ 9 \end{array} $	2 A 1 A	3 A 5 A	.01 .32	
11 12 13 14 15	83 83 81 82 82	4 P 1 P 5 P 3 P 4 P	61 69 63 56 52	4 A 4 A 12 P 5 A 6 A	72.0 76.0 72.0 69.0 67.0	80 64	$egin{array}{c} 9.6 \ 4.9 \ 9.3^1 \ 15.0 \ 14.8^1 \ \end{array}$	re n e se	163 117 77 113 95	14 8 12 12 10	3-30 A 4 P	3 P 5 P	.63 .01	
16 17 18 19 20 21	82 73 81 83 76 82	4 P 5 P 4 P 5 P 3 P	53 63 52 65 55 49	6 A 12 P 5 A 12 P 12 P 5 A	67.5 68.0 66.5 74.0 65.5 65.5	93 71 82 63 62	15.0 0.4 13.5 8.7 14.9 14.5	s s s nw w	86 114 95 224 143 69	11 10 15 17 10	7 а	12 м	.50 T T	
22 23 24 25 26 27	84 86 92 91 86 84	2 P 5 P 3 P 2 P 4 P 5 P	54 62 67 64 62 55	5 A 6 A 6 A 12 P 5 A	69.0 74.0 79.5 77.5 74.0 69.5	72 67 76 68	$ \begin{array}{c c} 11.0 \\ 9.4 \\ 10.4 \\ 10.0 \\ 14.6 \\ 12.3 \end{array} $	s w s n n	71 45 70 125 128 59	$\frac{14}{25}$ 13	7 Р	9 Р	Т .44	
28 29 30 31	73 82 72 78	4 P 3 P 6 P 4 P	61 59 61 67	12 P 6 A 1 A 4 A	67.0 71.5 66.5 72.5	91	0.0 14.6 0.4 2.1	nw n se €	81 109 106 81	20 6	8 A 6-30A	9 P 8 P	1.90	

^{*}Based on least time required to blow one mile.

July 1941	Normal	Extremes
Barometer, in inches		,
(Readings reduced to sea level)		
Maximum	30.27	30.50, 1892
Minimum	29.59	29.27, 1932
$Mean semi-daily \dots 29.936$	29.96	
Range0.62	0.68	0.97, 1892; 0.47, 1938
Air Temperature, in degrees F.		
Highest	93.9	104, 1911
Lowest	46.4	40, 1890, 1898
Mean	70.8	74.7, 1921; 66.3, 1891
Range	47.7	
Highest mean daily 81, 2nd		
Lowest mean daily 65, 4th		
Mean maximum		
Mean minimum		
Greatest daily range		
Least daily range		
Precipitation, in inches		
Precipitation	4.10	14.51, 1897; 0.70, 1929
Maximum precipitation in 24 hrs. 1.90, 28th	4.10	14.01, 1001, 0.10, 1020
	11	20, 1915; 4, 1924
Number of days with .01 or more 10	11	20, 1915, 4, 1929
Wind, in miles	0.400	5005 1000 1100 100 t
Total movement	3422	5097, 1909; 1109, 1894
Greatest daily movement 224, 19th		
Least daily movement 45, 23rd		
Mean hourly velocity 4.6		
Maximum velocity $\dots 25, 25$ th	28.7	44, 1936
Wind, direction		
Prevailing direction	SW	
North, days3		
Northeast, days0		
East, days		
Southeast, days		
South, days		
Southwest, days		
West, days		
Northwest, days		
	ļ	
Weather Many relative hymidity, persont 72.0	68.4	
Mean relative humidity, percent 72.9		en 1011-21 1024
Mean cloudiness, percent	50.9	[69, 1914; 31, 1924 22, 1923; 0, 1915
Number of clear days		22, 1925, 0, 1916
Number of fair days	14	24, '09, '21; 5, '89, '23, '28, '36
No	_	
Number of cloudy days		18, 1889; 0, 1910
Number hours bright sunshine300	268	371, 1910; 180, 1931
Percent of possible hours of bright	-0	
sunshine	58	
Thunder and lightning . 1st, 2nd, 8th, 12th		
13th, 25th, 28th	1	

The weather during July was very near normal in most respects. The mean temperature for the month was 71.2 degrees, the normal being 70.8 degrees. The highest temperature was 93 degrees on the 2nd and the lowest, 49 degrees on the 21st. The rainfall during the month was 4.04 inches which is .06 less than normal. The heaviest rainfall of the month was 1.90 inches on the 28th. The total rainfall since January 1 is now 19.02 inches, while the normal for this period is 25.30 inches. There were 300 hours of bright sunshine compared to a normal of 268 hours. The mean relative humidity was 72.9 per cent while the normal for July is 68.4 per cent.

Dr. J. K. Shaw reports as follows: "The weather during July continued to be favorable for the development of fruit plants. Timely showers supplied ample soil moisture for growth of both trees and fruits. A heavy shower accompanied with high winds on July 28th damaged a few trees and blew over some corn and tobacco. The dry spring was followed by sufficient precipitation and seems to have given conditions very favorable to growth. Apple trees have grown much better than last year. High temperatures in early spring which brought trees in bloom very early still show their effect in unusually early maturity. The blueberry crop ripened nearly three weeks earlier than last year."

Meteorological Series

Bulletin No. 632

August 1941

Meteorological Observations

FOR

AUGUST

1941

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Те	mpera	ture		ty.			Wind		Pr	ecipitatio	n
	Maxi	imum	Min	imum		lumidi	Hour			y.			
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Numl er of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water
$ \begin{array}{c} 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 8 \\ 9 \\ 10 \\ 11 \\ 12 \\ \end{array} $	\$8 \$6 \$3 \$4 \$6 \$5 \$6 91 \$5 78 \$0 \$3	5 P 4 P 2 P 6 P 3 P 4 P 2 P 4 P 2 P 4 P 2 P	65 67 58 55 57 57 53 53 57 52 46 53	3 A 5 A 3 A 5 A 5 A 6 A 6 A 6 A 3 A 12 P 6 A 12 P	76.5 76.5 70.5 69.5 71.5 71.0 69.5 72.0 71.0 63.0 68.0	61 62 68 68 65 62 62	9.0 13.5 13.8 14.2 14.3 14.3 14.2 6.8 14.2 10.6 11.1	nw n n n n n n n w n w n w n w n w n w	68 112 135 50 99 73 77 63 54 205 92 267	14 18 4 18 10 10 9 8	11-30 p 0 a	5 д 12 р 12 30 д	.08 T T T .01
13 14 15 16 17 18 19 20 21	68 74 75 80 71 80 70 78 79	2 P 2 P 11 A 3 P 3 P 3 P 3 P 3 P	45 46 59 60 49 41 60 52 47	5 A 5 A 3 A 12 P 12 P 6 A 4 A 12 P 6 A	56.5 60.0 67.0 70.0 60.0 60.5 65.0 63.0	51 57 77 74 55 64 93	13.2 10.7 1.9 8.0 13.9 12.0 0 0 13.7 13.2	nw sw s nw nw w s nw	254 123 177 160 243 78 151 145 62	32 12 15 17 22 8 11 14	12 m 2-30 a 0 a 3-30 a	12 p 3 a 2 a 10 a	.06 .17 .17
22 23 24 25 26 27 28 29	79 81 81 70 89 75 70 75	3 P 1 P 3 P 12 M 4 P 2 P 3 P 5 P	57 56 47 59 67 53 45 42	3 A 12 P 6 A 5 A 5 A 12 P 6 A 6 A	68.0 70.0 64.0 64.5 78.0 64.0 57.5 58.5	71 68 63 91 72 53 54 63	7.2 10.4 12.4 0.0 11.8 13.5 12.9 13.5	s sw sw s w mw mw w	147 117 82 127 156 244 155 64	$ \begin{array}{c} 13 \\ 13 \\ 10 \\ 10 \\ 17 \\ 25 \\ 22 \\ 9 \end{array} $	9 1	6-Р	.34 T
30 31	88 80	3 р 11 а	60 61	0 A 12 P	$74.0 \\ 70.5$	68 86	$\frac{11.0}{2.5}$	sw se	144 106		1-30a	12 г	.45

^{*}Based on least time required to blow one mile.

August 1941	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum	30.32	30.50, 1934
Minimum	-29.61	28.87, 1930
Mean semi-daily	30.00	
Range	0.71	
Air Temperature, in degrees F.		
Highest	91.6	100, 1918
Lowest 41, 18th	43.4	37, 1894, 1908
Mean	68.6	34, 1940
Range	48.1	
Highest mean daily 78, 26th		
Lowest mean daily 56.5, 13th	l	
Mean maximum		
Mean minimum		
Greatest daily range		
Least daily range 10, 19th		
Precipitation, in inches		
Precipitation		8.40, 1928; .31, 189-
Maximum precipitation in 24 hrs. 0.47, 19th		
Number of days with .01 or more 9	11	16, 1892, 1933; 4, 1899
Wind, in miles		
Total movement	3127	4,271, 1910; 1,920, 189
Greatest daily movement 267, 12th		
Least daily movement		
Mean hourly velocity 5.4	4.2	
Maximum velocity 40, 12th	22.7	35, 1917
Wind, direction		
Prevailing direction	sw	
North, days3	, ,,	
Northeast, days		
East, days0		
Southeast, days		
South, days		
Southwest, days		
West, days4		
Northwest, days12		
Weather		
Mean relative humidity, percent66.3	70.2	
Mean cloudiness, percent	49.7	67, 1901; 27, 1923
Number of clear days23	9	21, 1939; 0, 1915, 1929
Number of fair days4	13	25, 1912; 3, 1939
Number of cloudy days4	19	18, 1901, '28; 2, 1910, '28
Number hours bright sunshine	237	317, 1921; 152, 1915, 1929
Percent of possible hours of bright		31., 302., 102.
sunshine	55.2	
Thunder and lightning 10th, 17th, 23rd,	0.7.2	
31st		
First frost	Sept. 21	Aug. 22, '94, '95; Oct. 13, '09
	, clic. 21.	11 (1g. 22, 01, 00) () (1. 10, 0)

August was slightly cooler than normal, had less rain than usual and had an unusually large amount of sunshine. The mean temperature for the month was 67.1 degrees which is 1.5 degrees cooler than normal. The highest temperature during the month was 91 degrees on the 8th and the lowest 41 degrees on the 18th. The mean temperature for the three summer months. Iune, Iuly and August, was 68.5 degrees, while the normal mean temperature for these months is 68.7 degrees. The highest temperature during the summer was 94 degrees on June 27. The total rainfall during August was 1.79 inches, the normal being 4.08 inches. The total rainfall since January 1st is 20.81 inches while usually we get 29.38 inches during this period. There were 332 hours of bright sunshine compared to a normal of 237 hours. This is the largest amount of sunshine ever recorded in August at this station. The nearest approach to this high amount was 317 hours in 1921. The mean relative humidity was 66.3 per cent, the normal being 70.2 per cent. The total wind movement was 4030 miles. The normal wind movement for August is 3127 miles. The highest wind velocity was 40 miles per hour on the 12th.

Dr. J. K. Shaw makes the following report: "The progress of vegetation during August has been about normal. The effects of the early start in the spring are still seen in the early ripening of most varieties. There are still some signs of a low water supply but not more than in most seasons. Apple scab is less prevalent than usual, partly because of little weather favorable to infection last spring. Vegetative growth of most fruit plants through the season has been better than usual. This suggests that a dry spring followed by adequate summer rainfall is favorable to fruit plants, if not for the hay crop."

Meteorological Series

Bulletin No. 633

September 1941

Meteorological Observations

FOR

SEPTEMBER

1941

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Ter	mpera	ture		>.			Wind		Precipitation			
	Maximum		Minimum			fumidit	Hours Shine			x.				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Numl er of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour	Began	Ended	Water	
1 2 3 4	89 74 78 76	1 P 2 P 3 P 2 P	60 59 44 46	3 A 12 P 5 A 4 A	74.5 62.0 61.0 61.0	74 62 64 82	10.4 13.1 13.0 3.8	nw nw sw s	196 187 68 76	28 11 7 10	0 а	8 a	2 04	
5 6 7 8 9	83 86 79 72 75	3 P 3 P 1 P 3 P	64 61 55 51 48	6 A 12 P 12 P 6 A 5 A	73.5 75.0 67.0 61.5 61.5	86 62 49 64 80	3.4 13.0 12.9 12.9 8.0	nw nw nw nw	135 202 252 153 83	$ \begin{array}{r} 9 \\ 22 \\ 28 \\ 14 \\ 9 \end{array} $	0 а	8 a	.01	
$\frac{0}{1}$ $\frac{2}{3}$	91 70 68 71	3 P 1 P 3 P 3 P	66 52 42 38	4 A 12 P 6 A 6 A	$\begin{array}{c} 78.5 \\ 61.0 \\ 55.0 \\ 54.5 \\ 60.5 \end{array}$	79 57 63 69	5.5 11.2 12.7 12.6 12.6	nw nw nw	141 128 144 58 118	$ \begin{array}{r} 14 \\ 28 \\ 20 \\ 6 \\ 14 \end{array} $	8 P 0 A	12 P 1 A	.78 .01	
4 5 6 7 8	79 86 91 84 78	2 P 2 P 1 P 3 P 2 P	42 55 65 64 54	6 A 3 A 6 A 12 P	70.5 78.0 74.0 66.0	67 67 55 52	$\begin{array}{c} 12.5 \\ 11.2 \\ 11.4 \\ 12.4 \end{array}$	sw se nw n	162 1, 6 224 173 105	12 10 25 15 10				
$\frac{9}{0}$ $\frac{1}{2}$ $\frac{2}{3}$	70 71 80 86 86	4 P 4 P 4 P 3 P 2 P	42 34 34 49 46	7 A 7 A 7 A 6 A 7 A	56.0 52.5 57.0 67.5 66.0	62 68 73 70	12.3 12.3 9.7 8.8 10.0	nw s s s	64 85 84 101	8 7 7 13				
$\frac{4}{5}$ $\frac{6}{6}$ $\frac{7}{8}$	78 74 71 71 75	3 P 4 P 3 P 4 P	50 42 48 37 38	12 P 5 A 12 P 7 A 5 A	64.0 58 0 59.5 54 0 56.5	70 59	$ \begin{array}{c c} 12.0 \\ 7.8 \\ 11.9 \\ 12.0 \\ 10.7 \end{array} $	n s nw s	148 73 203 54 109	11 10 20 4 13			Т	
.0 !9 :0 :1	59 60	5 P 12 P	33 28	12 P 5 A	46.0 44.0	60	3.9 3.0	n s	240 73	$\begin{array}{c} 25 \\ 17 \end{array}$	Й:30 г	12 P	.04	

^{*}Based on least time required to blow one mile.

September, 1941	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum	-30.45	30.65, 192
Minimum	29.57	28.41, 193
Mean semi-daily	-30.06	
Range	.88	1.99, 1938; .57, 1910
Air Temperature, in degrees F.		
Highest 91, 10th, 16th	87.7	97, 1895, 1929
Lowest 28, 30th	33.2	24.5, 191-
Mean	61.7	
Range	54.5	
dighest mean daily 78 5, 10th		
Lowest mean daily		
Mean maximum		
Mean minimum		
Greatest daily range 40, 23rd		
Least daily range 18, 11th		
Precipitation, in inches		
Precipitation	4.26	14.55, 1938; .52, 191
Maximum precipitation in 24 hrs. 2.04, 1st		
Number of days with .01 or more 5	10	16, 1933; 3, 190
Wind, in miles		
Fotal movement	3271	4,686, 1896; 1,414, 189
Greatest daily movement 252, 7th		
Least daily movement		
Mean hourly velocity 5.5	4.5	
Maximum velocity 28, 1st, 7(h, 11th	25.9	80, 1938
Wind, direction		
Prevailing direction	WSW	
North, days4		
Northeast, days0		
East, days0		
Southeast, days2		
South, days		
Southwest, days		
West, days 0		
Northwest, days		
Weather		
Mean relative humidity, percent 67.0	73.6	
Mean cloudiness, percent	49.9	70, 1934; 27, 1908, 1914
Number of clear days23	10	19, 1932; 2, 1907, 1928
Number of fair days4	10	19, 1908; 3, 1889
Number of cloudy days	10	20, 1934; 4, 1908, 1916
Number hours bright sunshine 307	201	255, 1916; 106, 193-
ercent of possible hours of bright		
sunshine	54.9	
Thunder and lightning 1st, 10th		
First frostSept. 30	ept. 21	Aug. 22, '94, '95; Oct., 13, '69
	chira mi	

The weather during September was dry, warm and clear. The total rainfall during the month was 2.88 inches compared to a normal of 4.26 inches. The total rainfall since January 1 has been 23.69 inches, while the normal rainfall for this period is 33.62 inches. This deficiency of almost 10 inches has resulted in acute water shortage in wells and reservoirs. The rainfall since April 1 has been 18.26 inches or 4.85 inches less than normal for that period with some damage to crops.

The mean temperature for the month was 62.5 degrees, the normal being 61.7 degrees. There was a light frost on the 20th on nearby lowland and a heavy frost on the 30th with a temperature of 28 degrees at the observatory. The normal date for the first frost is Sept. 21st. The domestic heating load for the month was 149 degree-days, which is normal for September. There were a total of 307 hours of bright sunshine, which is the largest amount of sunshine ever recorded at this station in September.

Dr. J. K. Shaw of the Pomology Department makes the following report: "The weather during September has been very dry. Leaves are dropping from some apple trees prematurely. This may result in poor foliation in the spring. Unless there are abundant rains soon, fruit plants will go into the winter in poor condition. The lack of rainfall has not interfered with the crop of this year and plenty of sunshine has favored good color development. The apple crop is of very good quality, but not large. There seems to be good fruit bud formation for next year's crop."

Meteorological Series

Bulletin No. 634

October 1941

Meteorological Observations

FOR

OCTOBER

1941

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 48.5″ N. Longitude, 72° 31′ 10″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Те	mpera	ıture		Λ.			Wind		Pr	ecipitatio	n
	Maximum		Minimum			umidit	umidit; Hours shine			T.			
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Numl er of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour	Бедап	Ended	Water
1 2 3 4 4 5 6 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 22 5 26 27 28 29	70 73 67 82 88 77 60 68 65 65 66 67 70 67 67 67 67 67 67 67 67 68 68 68 68 68 68 68 68 68 68 68 68 68	2 P P 3 P P 12 P	51 41 49 629 53 60 44 46 38 37 41 41 48 42 43 43 43 43 43 43 44 43 43 44 43 43 44 44	12 P 6 A 3 A 12 P 2 A 12 P 4 A 7 A 6 A 12 P 11 A 6 A 4 A 11 P 12 P 6 A 2 A 12 P 7 A 7 A 12 P 12 P 12 P 12 P 12 P 13 P 14 A 15 P 16 A 17 A 18 P 18 P 18 P 18 P 18 P 18 P 18 P 18 P	60.5 57.0 58.0 72.0 73.5 64.0 54.5 53.0 54.0 54.5 55.0 46.5 49.5 52.5 52.0 44.0 54.5 52.5 52.0 44.0 53.5 53.0 54.0 53.5 53.0 54.0 54.0 54.0 54.0 54.0 54.0 54.0 54	69 72 96 77 86 77 89 54 63 63 66 63 57 66 63 67 69 63 748 57 789 80	2.2 9.4 0.0 10.6 5.5 9.6 0.0 7.6 9.6 0.0 3.3 11.2 0.3 5.4 10.1 11.0 0.0 10.9 2.7 9.7 9.7 9.1 1.8 1.8 1.8	nw sw sw w sw w sw mw se nw se nw w sw mw w se nw w nw w s nw mw nw nw nw s nw mw nw s s nw mw s s nw mw nw s s	2088 566 366 1599 1890 1890 1890 1890 1991 1992 1992 1992 1992 1992 1992 19	18 4 8 4 8 5 25 18 9 32 20 22 32 15 7 10 25 5 10 10 32 15 20 10 32 15 20 10 32 15 20 10 32 15 20 10 32 15 20 10 32 15 20 10 32 15 20 10 32 15 20 10 32 16 20 10 32 17 20 10 32 18 20 32 18 32 18 32	0 A 5 A 0 A 5 A 4 A 4 P	2:30 A 12 P 5 A 8 A 8 P	.42 .42 .04 .17 T .72 T .08
30 31	41 54	4 P 3 P	29 38	6 A 0 A	35.0 46 0	77 85	$0.0 \\ 0.5$	n n	55 64	9			T

^{*}Based on least time required to blow one mile.

October, 1941	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum 30.60, 29th, 30th	30.53	30.68, 192
Minimum	29.42	29.00, 192
Mean semi-daily	30.06	20.00, 102
Range	1.11	1.47, 1926; .76, 189
	1.11	1.47, 1520, .70, 165
Air Temperature, in degrees F.	70.1	00 - 100
Highest	79.4	90.5, 190
Lowest	23.4	17, 1930
Mean	50.5	56.4, 1920; 43.2, 189
Range	56.0	
Highest mean daily		
Lowest mean daily	1	
Mean maximum		
Mean minimum		
Chantout dully manage 97 17th		
Greatest daily range		
Least daily range		
Precipitation, in inches		
Precipitation	3.29	8.81, 1911; .01, 192-
Maximum precipitation in 24 hrs72, 10th		, , , , , , , , , , , , , , , , , , , ,
Number of days with .01 or more 8	9	15, 1913; 1, 1897, 192-
		10, 1510, 1, 1001, 102
Wind, in miles	40-4	F 10 F 101 0 F10
Total movement	4074	5,467, 1910; 2,540, 189-
Greatest daily movement 295, 8th		
Least daily movement		
Mean hourly velocity 6.0		
Maximum velocity	29.5	42, 1937
	20.0	12, 100
Wind, direction		
Prevailing direction WNW	11.	
North, days6		
Northeast, days0		
East, days0		
Southeast, days2		
South, days		
Southwest, days4		
West, days		
Northwest, days10		
Weather		
Mean relative humidity, percent70.7	69.0	
Mean cloudiness, percent	48	67, 1932; 18, 192-
Number of clear days	10	21, 1938; 1, 1911
Number of fair days4	10	17, 1924; 3, 1938
	11	
Number of cloudy days	1	19, 1896; 3, 1924
Number hours bright sunshine165	175.0	232, '23, '38; 91, 1913
Percent of possible hours of bright		
sunshine	51.3	
Thunder and lightning		
	Samt 91	Aug. 22 '04 '05: Oat 15 '20
First frostSept. 30	cept. an	八、は鳥に高る。 ひまこ ひひょくがしこしひこ ひに

The drouth continued through October with a rainfall of only 2.13 inches, the normal for October being 3.29 inches. The total precipitation since January 1 is now 25.82 inches whereas the normal for this period is 36.91 inches. Reports have come in of additional wells going dry during the month.

The mean temperature of the month was 52.5 degrees which is 2 degrees higher than normal. The highest temperature was 88 degrees on the 5th and the lowest 24 degrees on the 29th. The domestic heating load was 407 degree-days, the normal for October being 456 degree-days. There were 165 hours of bright sunshine compared to a normal of 175 hours.

Meteorological Series

Bulletin No. 635

November 1941

Meteorological Observations

FOR

NOVEMBER

1941

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Te	mpera	ture		8	10	Wind Precipitation							
	Maximum		um Minimum			fumidíí	f Hours shine			T.					
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Numl er of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow	
1 2 3 4	46 56 58 60	8 P 2 P 2 P 3 P	42 44 38 29	8 A 7 A 12 P 7 A	44.0 50.0 48.0 44.5	65 69	0.0 0.6 6.6 8.8	nw w w	77 84 169 82	15 15 17 13	0 a 0 a	12 P 4 A	2.06 .03		
5 6 7 8 9	63 51 57 55 51	3 P 7 P 2 P 2 P 2 P	40 42 43 36 30	7:30 _A 6 A 12 P 7 A 6 A	51.5 46.5 50.0 45.5 40.5	$\frac{74}{76}$	$\begin{array}{c} 3.3 \\ 0.0 \\ 0.6 \\ 2.0 \\ 2.5 \end{array}$	n n s s	68 81 127 116 81	6 9 18 10 9	7 A 0 A 5 P	12 р 3 р 12 р	.49 T		
10 11 12 13 14	51 45 44 40 58	1 P 11 A 2 P 2 P 3 P	32 29 26 24 27	12 P 6 A 12 P 2 A 6 A	$\begin{array}{c} 41.5 \\ 37.0 \\ 35.0 \\ 32.0 \\ 42.5 \end{array}$	60 69 74	4.2 0.5 7.6 0.0 1.6	nw w nw nw sw	72 80 246 49 66	10 13 32 5 11				Т	
15 16 17 18 19 20	59 57 48 62 74 72	2 P 12 M 4 P 2 P 3 P 1 P	35 34 25 28 37 45	7 A 12 P 7 A 0 A 7 A 7 A	$\begin{array}{r} 47.0 \\ 45.5 \\ 36.5 \\ 45.0 \\ 55.5 \\ 58.5 \end{array}$		4.2 5.0 9.7 5.7 6.9 0.0	nw n se	$ \begin{array}{ c c c } 75 \\ 214 \\ 168 \\ 150 \\ 89 \\ 176 \end{array} $	9	1 Р	1:30 р	.04		
21 22 23 24 25	48 47 46 42 44	12 M 4 P 7:30 P 1 P 2 P	35 27 30 26 19	9 P 7 A 0 A 12 P 7 A	41.5 37.0 38.0 34.0 31.5	54 68 88 57 70	7.8 8.2 0.0 7.8 9.1	HW SW HW HW	340 101 42 30.1 118	15 18 32 11	5 а	12 м	.32		
26 27 28 29 30	$\begin{bmatrix} 49 \\ 52 \\ 51 \\ 64 \\ 51 \end{bmatrix}$	1 P 1 P 2 P 2 P 0 A	32 32 26 25 28	0 A 12 P 5 A 3 A 12 P	40.5 42.0 38 5 44.5 39 5	58 63 64	2.8 4.3 9.4 8.8 7.0	s nw s n	201 183 131 93 214	20				T	

^{*}Based on least time required to blow one mile.

Sol Kriminetsky, Observer

November, 1941	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)	00.70	00.07 1000
Maximum		30.87, 1932
Minimum		28.73, 1904
Mean semi-daily	$\frac{30.05}{1.29}$	1.04 1001- 04 1005
Air Temperature, in degrees F.	1.29	1.84, 1904; .94, 1905
Highest	66.0	75, 1924, 1938
Lowest	12.9	-4, 1938
Mean	38.9	44.1, 1931; 33.6, 1901
Range		1111, 1001, 0010, 1001
Highest mean daily 58 5, 20th		
Lowest mean daily 31.5, 25th		
Mean maximum		
Mean minimum		
Greatest daily range 39, 29th		
Least daily range 4, 1st		
Precipitation, in inches		
Precipitation 4.29		8.64, 1927; .63, 1917
Snow Trace	2.34	13.50, 1938
Maximum precipitation in 24 hrs. 2.06, 1st		
Number of days with .01 or more	9	22, 1921; 2, 1904
Wind, in miles		
Total movement		5,978, 1906; 2,589, 1889
Greatest daily movement 340, 21st		
Least daily movement 42, 23rd		
Mean hourly velocity 5.6	6.3	11 1096
Maximum velocity	30.3	44, 1938
Wind, direction	****	
Prevailing direction W	WNW	
North, days4		
Northeast, days 0		
East, days		
South, days		
Southwest, days		
West, days		
Northwest, days		
Weather		
Mean relative humidity, percent70.7	70.6	
Mean cloudiness, percent		72, 1900, 1927; 34, 1917
Number of clear days		15, 1903; 1, 1900, 1911, 1927
Number of fair days		16, '12; 4, '89, '30, '32, '35
Number of cloudy days		24, 1927; 9, 1905, 1917
Number hours bright sunshine135	121	182, 1903; 66, 1927
Percent of possible hours of bright		
sunshine	41.3	
Thunder and lightning		

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1940.

The weather during November was warmer than usual, more sunshine than normal, and heavy rainfall during the first part of the month. The mean temperature for the month was 42.8 degrees, the normal being 38.9 degrees. This is the warmest November at Amherst since 1889, except for four years. It was equalled in 1934 and exceeded in 1927, 1931, and 1935 with mean November temperatures of 43.9, 44.1 and 43.4 degrees. The highest temperature was 74 degrees on the 19th, which was the highest temperature ever recorded so late in the season at this station. temperature was 19 degrees on the 25th. The domestic heating load for the month was 667 degree-days, the normal for November being 784 degree-days. The total degree-days for the season is now 1223 degree-days compared to a normal of 1389 for that period. There were 135 hours of bright sunshine, the normal for November being 121 hours. The total rainfall for the month was 4.29 inches. whereas the normal for November is 3.41 inches. Of this amount, over 2 inches fell on the first day of the month and $1\frac{\pi}{2}$ inches on the 6th and 7th. The total rainfall since January 1 is now 30.11. The normal being 40.32 inches. Snow flurries were seen on the 12th and 26th.

Meteorological Series

Bulletin No. 636

December 1941

Meteorological Observations

FOR

DECEMBER

1941

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Ter	mpera	ture		2.			Wind			Precipit	ation	
	Max	imum	Min	imum		Inmidit	Hourshine			7.				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Pright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4	36 47 50 42	3 r 4 p 3 r 2 p	17 32 33 33	7 A 0 A 11 P 0 A	26.5 39.5 41.5 37.5	100 100 100	8.8 0.0 2.7 0.2	n se nw sw	125 86 26 25	9 9 4 4	4 A		0.20	
5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 22 1 22 23 24 5 26 27 28 29 31	50 51 34 39 41 42 26 37 41 36 45 45 44 46 40 25 30 31 56 45 45 45 47 49 40 20 20 20 20 20 40 40 40 40 40 40 40 40 40 4	S P	39 31 21 17 29 25 21 8 20 33 29 28 25 32 28 28 32 30 32 40 30 30 40 40 40 40 40 40 40 40 40 40 40 40 40	0 A 12 P 12 P 6 A 12 P 7 A A 12 P 7 A A 7 A 12 P 7 A A 7 A 12 P 7 A A 7 A 12 P 7 A A 7 A 12 P 7 A A 7 A 12 P 7 A A 7 A 12 P 7 A A 7 A 12 P 7 A A 7 A 12 P 7 A A 7 A 12 P 7 A A 7 A 12 P 7 A A 4 A 7 A 12 P 7 A A 7 A 12 P 7 A A 7 A 12 P 7 A A 7 A 12 P 7 A A 1 A 12 P 7 A A 1 A 12 P 7 A A 1 A 12 P 7 A A 1 A 12 P 7 A A 1 A 1 A 1 A 1 A 1 A 1 A 1 A 1 A 1	44.5.41.0 27.5.28.0 33.5.0 33.5.19.0 32.5.3 32.5.0 32.0 32.0 32.0 32.0 32.0 32.0 32.0 32	100 721 61 63 63 54 63 78 62 56 63 63 63 63 63 63 63 63 63 63 63 63 63	0.2 7.3 5.7 5.4 4.7 9.1 4.5 6.2 0.0 8.2 3.7 8.2 5.6 9.0 0.0 8.1 4.7 8.2 9.1 9.1	nw nw nw nw nw nw nw nw nw nw nw nw nw n	31 212 417 418 239 258 267 126 6107 334 4186 6184 186 285 71 139 196 134 148 111	4 32 28 22 25 15 44 28 22 20 18 25 32 20 9 3 8 20 11 25 22 15 15 2	11 A O A O A S P O A	9 а	0.92 1.20 0.15 1.00 T T	1.00

^{*}Based on least time required to blow one mile.

December, 1941	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum	30.66	30.96, 1889
Minimum	29.27	28.85, 1915
Mean semi-daily	30.06	20.00, 101.
	1.38	1 79 1905, 1 01 1909
Range	1.00	1.78, 1895; 1.01, 1892
Air Temperature, in degrees F.	510	# 5 1 100C
Highest	54.3	65.5, 1908
Lowest 5, 21st	-1.9	-22.5, 1917
Mean31.1	27.5	36.9, 1891; 17.1, 1917
Range	£6.2	
Highest mean daily 45, 24th		
Lowest mean daily 15, 21st		
Mean maximum		
Mean minimum		
Greatest daily range 30, 20th		
Least daily range		
Precipitation, in inches	0 - 0	
Precipitation	3.39	7.77, 1901; .87, 1896
Snow	8.50	[26.50, '02; t, 1891
Maximum precipitation in 24		
hrs 2.12, 13th, 14th		
Number of days with .01 or more 8	10	17, 1902; 4, 1892
Wind, in miles		<i>' ' '</i>
	4710	6,694, 1925; 3,239, 1918
Total movement	4710	0,094, 1529, 5,255, 1916
Greatest daily movement		
Least daily movement		
Mean hourly velocity 6.8	6.3	
Maximum velocity 32, 6th, 20th	31.4	48, 1938
Wind, direction		
Prevailing direction NW	WNW	
North, days6	11 111	
Northeast, days		
East, days		
Southeast, days		
South, days		
Southwest, days1		
West, days0		
Northwest, days		
Weather		
Mean relative humidity, percent 70.6	69.4	
Mean cloudiness, percent	54.9	71, 1929; 39, 1919
Number of clear days9	8	15, 1890; 2, 1922, 1933
		16, 1000, 2, 100, 1000
Number of fair days	9	16, '09; 4,'89 '30, '31, '36. '38 23, '33; 7, '09. '23
Number of cloudy days9	1.4	25, 35; 7, 79, 25
Number hours bright sunshine145	128	172, 1896; 63, 1933
Percent of possible hours of bright		
II.	45.2	
sunshine	40.4	

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1940.

ANNUAL SUMMARY

Annual 1941	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum	30.81	31.05, 1920
Minimum	28.95	28.41, 1938
Mean semi-daily	-30.01	
Range	1.85	2.47, 1938; 1.38, 1933
Air Temperature, in degrees F.		·
Highest	95.7	104, 1911
Lowest9, Jan. 30	-12.2	-26, 1904
Mean	47.4	49.8, 1921; 44, 1904
Range103	107.8	
Highest mean daily81, July 2		
Lowest mean daily 4.5, Jan. 30		
Mean maximum	57.8	
Mean minimum	36.5	
Greatest daily range 44, April 29, May 6		
Least daily range		
Precipitation, in inches		
Precipitation	43.70	59.00, 1938; 30.68, 1908
Snow	47.78	89.00, 1893; 24.50, 1919
Max. precipitation in 24 hrs.2.12, Dec. 13,14		
Number of days with .01 or more90	124	144, 1902; 96, 1924
Wind, in miles		111, 1552, 65, 151
Total movement	52,223	63,571, 1908; 36,257, 1894
Greatest daily movement 549, Mar. 19		,,,,
Least daily movement 11, Dec. 31		
Mean velocity	5.8	
Maximum velocity	39.5	80, 1938
Wind, direction	150,00	00, 1000
Prevailing direction WNW	W	
North, days	,,,	
Northeast, days		
East, days 4		
Southeast, days		
South, days		
Southwest, days		}
West, days		
Northwest, days141		
Weather	0-0	
Mean relative humidity, percent68.7	67.6	00 200 201 200 11 200 29
Mean cloudiness, percent	51.7	60, '98, '01, '02; 41, '08, '24
Number of clear days	116	169, 1936; 59, 1927
Number of fair days	123	182, 1912; 64, 1936
Number of cloudy days	126	179, '01, '02; 71, 1910
Number hours bright sunshine 3,038	2,353	2838, 1924; 1864, 1902
Percent of possible hours of bright	~ O . O	
sunshine	52.8	N. 11 110 N. 11 107
Last snowApril 17	April 15	Mar. 14, 10; May 11, 07
First snowDec. 13	Nov. 6	Oct. 10, '25; Nov. 27, '31
r a		
Last snow April 17 First snow Dec. 13 Last frost May 13 First frost Sept. 30	May 14	Apr. 23, '04; June 8, '32

Meteorological Series

Bulletin No. 637

January 1942

Meteorological Observations

FOR

JANUARY

1942

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Те	mpera	ture		>.			Wind			Precipits	ıtion	
	Maxi	mum	Min	imum		umidit	Hours			ν:				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Numl er of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3	38 56	12 P 9 A	22 34	0 A 12 P	$\frac{30.0}{45.0}$	67	0.0	n nw	13 327	8 32	6:30 p 0 a	12 P 8 P	.13 .52	.25
3 4 5 6	$ \begin{array}{r} 37 \\ 29 \\ 34 \end{array} $	3 P 1 A 2 P	27 23 13	10 P 11 P 10 P	32.0 26.0 23.5	53 90 86	8.2 0.8 4.7	nw n w	274 148 42	32 15 8	2 A	5 г	.36	5.50
6 7 8 9 10 11 12 13 14	16 14 15 19 20 17 32 32 40	3 P 2 P 2 P 12 M 4 P 12 P 4 A 4 P	$ \begin{array}{c} 0 \\ 0 \\ -13 \\ -10 \\ 15 \\ -10 \\ -1 \\ 8 \\ 3 \end{array} $	10 P 7 A 7 A 5 A 8 A 4 A 8 A 11 P 4 A	8.0 7.0 1.0 4.5 17.5 3.5 15.5 20.0 21.5	55	9.2 8.0 9.3 4.9 7.1 0.0 9.4 7.2	nw sw s nw se s nw	128 234 49 111 168 151 170 263 174	14 22 7 13 22 15 18 28 18	5 а	8 a	.03	.50
15 16 17 18 19 20 21 22 23	40 29 32 47 44 40 36 39 45	2 A 4 P 3 P 1 A P 2 P 4 P 3 P 3 P	24 3 -4 24 35 33 23 17 22	7 P 12 P 7 A 5 A 6 A 5 A 12 P 7 A 6 A	32.0 16.0 14.0 35.5 39.5 29.5 28.0 33.5 39.5	65 52 65 66 92 76 60 74 73	7.1 9.5 9.5 5.8 0.0 4.3 7.0 5.3 4.2 5.6	nw nw s se nw nw nw w sw	252 348 121 165 100 190 221 58 132 139	32 35 12 11 9 25 20 8 15	8:30a	9 Р	1.41	т
$\frac{24}{25}$	47 34 32	2 P 2 P 1 A		7 A 8 A 12 P	$\begin{array}{c} -39.5 \\ -30.5 \\ -29.5 \end{array}$	67	$\begin{bmatrix} 3.0 \\ 2.7 \\ 0.0 \end{bmatrix}$	n ne	136 154					Т
$\frac{27}{28} \\ 29$	$\begin{vmatrix} 44 \\ 36 \\ 25 \end{vmatrix}$	3 P 1 P 3 P	$\begin{array}{c} 25 \\ 22 \\ 11 \end{array}$	3 A 12 P 12 P	$ \begin{array}{r} 34.5 \\ 29.0 \\ 18.0 \end{array} $	82 74 47	$7.1 \\ 0.0 \\ 9.9$	s n n	84 163 475	$\frac{25}{32}$	2 A	S A	.02	.25
30 31	33 34	4 P	$\frac{6}{22}$	7 A 2 A	$19.5 \\ 28.0$		9.1	n -e	194 105		12 м	12 Р	1.07	

^{*}Based on least time required to blow one mile.

January, 1942	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum	30.70	31.00, 192
Minimum 29.28, 2nd, 31st	29.20	28.55, 1913
Mean semi-daily	30.07	
Range	1.50	2.18, 1913; 0.97, 1896
Air Temperature, in degrees F.		
Highest	51.1	66.0, 1932
Lowest	-7.1	-26.0, 190
Mean	24.2	34.2, 1913; 13.9, 1918
Range	58.2	, , , , , , , , , , , , , , , , , , , ,
Highest mean daily 45, 2nd		
Lowest mean daily 1, 8th		
Mean maximum		
Mean minimum		
Greatest daily range 37, 14th		
Least daily range 5, 10th, 26th		
Precipitation, in inches		
Precipitation	3.61	7 15 1909 1 07 1906
Snow	13.32	7.15, 1898; 1.07, 1896
Maximum precipitation in 24	10.02	33, 1898, 1923; 1.50, '08
hrs 1.41, 19th	1.1	15 209 200 1 1001
Number of days with .01 or more	11	15, '93. '20; 4, 1901
Wind, in miles		
Total movement	5055	7770, 1908; 2896, 1895
Greatest daily movement 475, 29th		
cast daily movement 13, 1st		
Ican hourly velocity	6.8	
Maximum velocity 35, 16th	32.7	47, 1938
Wind, direction		
Prevailing direction NW	WNW	
North, days		
Northeast, days		
East, days0		
Southeast, days3		
South, days		
Southwest, days		
Vest, days		
Northwest, days		
Weather		
Jean relative humidity, percent68.4	70.1	
Mean cloudiness, percent	55	78, 1932; 37, 1908
Number of clear days	9	18, 1941; 2, 1914
	9	18, 1890, 1926; 3, 1916
Number of fair days		
Number of fair days		[22] [1928] [98] (4. 1920) [030
Number of fair days	13	22, 1923, 1931; 4, 1920, 1939 214 1920 · 74 1932
Number of fair days		
Number of fair days	13 137	22, 1923, 1931; 4, 1920, 1939 214, 1920; 74, 1932
Number of fair days	13	

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1941.

The weather during January was very near normal except that there was less snow than usual. Only 6.5 inches of snow fell during the month while we get an average of 13.32 inches of snow in January. The snowfall for the current season thus far is only 8.5 inches, while the normal for this period is 24.16 inches. The total precipitation during the month was 3.54 inches, the normal for January being 3.61 inches. The mean temperature during the month was 24.1 degrees compared to a normal of 24.2 degrees. The lowest temperature was –13 degrees on the 8th. The domestic heating load during the month was 1267 degree-days, the normal for January being 1265 degree-days. The total heating load for the season to the end of January is 3539 degree-days, while the normal for this period is 3817 degree-days. There were 159 hours of bright sunshine which is well above the average of 137 hours. The mean relative humidity was 68.4 per cent, the normal being 70.1 per cent.

Meteorological Series

Bulletin No. 638

February 1942

Meteorological Observations

FOR

FEBRUARY

1942

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

-		Те	mpera	iture		iy.	70		Wind			Precipita	tion	
	Max	imum	Min	imum		Iumidii	f Hours ashine		-	7C				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 23 34 45 66 78 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28	36 22 17 24 27 34 38 25 34 34 41 41 43 26 20 8 33 34 43 44 43 43 44 43 43 44 43 44 43 44 43 44 43 44 44	12 M 0 A A 3 P 12 P 2 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P	22 7 0 0 14 27 28 17 10 15 9 9 13 23 26 16 7 8 13 16 16 16 17 8 18 19 19 19 19 19 19 19 19 19 19	12 P 12 P 7 A 6 A 0 A 0 A 7 A 7 A 7 A 7 A 1 A 7 A 1 A 12 P 12 P 7 A 4 A 12 P 7 A 12 P 7 A 7 A 12 P 7 A 13 P 12 P 7 A 14 A 15 P 16 A 17 A 17 A 18 A 18 A 18 A 18 A 18 A 18 A 18 A 18	29.0 14.5 8.5 12.0 20.5 30.5 27.5 22.0 24.5 16.5 23.5 32.0 27.5 21.0 25.0 25.0 25.0 26.0 27.5 29.5 29.5	72 59 59 88 88 76 61 57 61 53 56 68 82 56 64 62 64 61	0.3 10.00 10.00 7.1 0.00 1.2 0.00 3.4 10.3 7.5 6.8 10.4 8.5 10.2 3.0 1.2 10.6 8.9 8.7 7.7 6.8 3.3 10.9 9.5 6.0	w nw nw nw nw n n n n n n n n n n n n n	204 364 264 70 80 2009 202 203 360 120 220 387 361 145 216 279 401 368 271 153 224 311 260 187	22 25 22 7 8 13 25 25 25 15 14 28 25 25 17 28 25 25 25 25 25 25 25 25 25 25 25 25 25	0 A 10 A 10 A 0 A 5 A	12 P 2 A 5 P 8 A	.56 .04 .32 .05	6.5 0.5 T 1.00 T T

^{*}Based on least time required to blow one mile.

February, 1942	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)	•	
Maximum :	30.63	31.05, 1920
Minimum	29.24	28.56, 1895
Mean semi-daily	30.04	20.00, 1000
Range	1.38	1.89, '00, '08; .88, '13 '31
Air Temperature, in degrees F.	1100	1.00, 00, 00, 100, 10 01
Highest	50.5	65.0, 1930
Lowest 0, 3rd, 4th	-7.4	-22.5, 1918
Mean	23.7	,
Range	58	
Highest mean daily38.5, 17th		
Lowest mean daily 8.5, 3rd		
Mean maximum	32.8	
Mean minimum	14.5	
Greatest daily range, 24, 4th, 10th, 24th		
Least daily range 7, 6th	1	
Precipitation, in inches		
Precipitation	3.19	8.12, 1900; 0.62, 1901
Snow 8.00	13.99	48.75, 1893; 0.50, 1937
Maximum precipitation in 24	li	
hrs 0.56, 5th		
Number of days with .01 or more	10	15, '93, '20; 4, 1901
Wind, in miles		
Total movement	4776	6445, 1896; 3438, 1892
Greatest daily movement 401, 21st		
Least daily movement 70, 4th		
Mean hourly velocity 9.7	7.1	
Maximum velocity32, 13th, 21st	31.7	48, 1934, 1937
Wind, direction		
Prevailing direction NW	WNW	
North, days		
Northeast, days0		
East, days0	l i	
Southeast, days	[
South, days0		
Southwest, days0		
West, days		
Northwest, days21		
Weather		
Mean relative humidity, percent67.1	66.6	
Mean cloudiness, percent 43	50.9	66, '90, '27; 31, 1905
Number of clear days17	10	19, 1941; 2, 1927
	8	16, 1920; 2, 1936
Number of fair days 4	10	16, 1894; 2 1920
Number of cloudy days7	- 1	
Number of cloudy days	158	221, 1924; 110, '27, '38
Number of cloudy days	158	221, 1924; 110, '27, '38
Number of cloudy days	- 1	221, 1924; 110, '27, '38

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1941.

The weather during February was marked by the small amount of rain and snow, the high wind, and the large amount of sunshine. The temperature was very near normal, with a mean temperature of 24 degrees, the normal being 23.7 degrees. The highest temperature was 44 degrees on the 17th and the lowest 0 on the 3rd and 4th. The normal low temperature for February is –7.4 degrees. It may be interesting to recall that the lowest temperature ever recorded at the station was –22.5 degrees in February, 1918. The domestic heating load for the month was 1149 degree-days, the normal for February being 1165 degree-days. The total heating load for the season is 4688 degree-days. The normal heating load to the end of February is 4982 degree-days. The normal heating load for February is based on the 50-year record including 11 leap years and hence is high compared with years which are not leap years.

The total precipitation for the month was 1.66 inches, while the normal for February is 3.19 inches. It is interesting to note that the precipitation in February, 1941 was 1.59 inches. The total precipitation for January and February of this year is 5.20 inches whereas for the same period in 1941 it was only 3.80 inches. Only 8 inches of snow fell in February compared to a normal of 13.99 inches. The total snowfall for the winter is 16.50 inches, while the normal to the end of February is 38.15 inches. There were 190 hours of bright sunshine, the normal being 158 hours. The total wind movement during the month was 6531 miles which is the highest ever recorded in February. The nearest to this record was 6445 miles in 1896. The normal for February is 4776 miles. The highest wind velocity was 32 miles per hour on the 13th and 21st. The wind blew from the northwest on 21 days.

Meteorological Series

Bulletin No. 639

March 1942

Meteorological Observations

FOR

MARCH

1942

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Те	mpera	ture		S.	_		Wind			Precipita	ation	
	Maxi	mum	Min	imum		lumidit	f Hours shine			<i>x</i> :				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5	45 46 43 43 55	3 P 4 P 2 P 3 P 3 P	31 32 32 32 32 27	7 A 7 A 6 A 12 P 4 A	38.0 39.0 37.5 37.5 41.0	63 63 91 69 58	4.9 6.3 0.0 4.1 9.0	n sw ne nw	65 68 271 253 136	8 10 48 25 17	2 A	12 Р	1.82	2.00
6 7 8 9	39 50 63 62 43	5 P 5 P 3 P 9 A	34 28 31 37 32	12 P 6 A 7 A 12 P	36.5 39.0 47.0 49.5 37.5	83 56 73 81	$\begin{array}{c} 3.0 \\ 0.0 \\ 11.5 \\ 10.7 \\ 3.0 \\ 5.5 \end{array}$	nw s s sw	151 184 145 358 436	20 15 17 35 32	6 A 8 P 0 A	3 р 12 р 4 р	.03	Т
10 11 12 13 14	54 48 52 36	3 P 4 P 2 P 5 P 2 P	29 34 28 31	5 A 7 A 12 P 7 A 7 A	41.5 41.0 40.0 33.5	43 47 80 52 73	11.6 1.8 11.7 0.0	w s nw nw	202 170 138 77	12 32 13 6	3 A 2 A	11 a 12 p	.13	3.25
15 16 17 18	38 47 38 53	1 P 4 P 2 P 3 P	30 31 33 35	6 A 5 A 3 A 5 A	34.0 39.0 35.5 44.0	89 87 99 80 60	4.6 7.3 0.0 5.6 6.3	n w nw nw	146 61 89 92 332	10 10 13 18 40	6:30P 0 A	12 р 10 30 р	.67 1.17	
19 20 21 22 23 24	42 50 52 47 40 43	5 P 4 P 2 P 4 P 2 P 3 P	37 28 27 33 30 28	8 A 6 A 3 A 12 P 4 A 5 A	39.5 39.0 39.5 40.0 35.0 35.5	69 74 66 62	12.1 7.4 6.3 5.6 6.3	w nw se w nw	283 128 280 280 252 207	25 15 40 25 25	5 P 0 A	12 р 6 30а	.87 .82	T T
25 26 27 28 29	46 52 59 59 51	5 P 2 P 4 P 3 P 2 P	28 32 26 31 26	6 A 6 A 6 A 6 A	37.0 42.0 42.5 45.0 38.5	59 60 61 60 52	11.7 10.0 12.4 9.2 9.0	nw n w e	235 226 83 169 180	32 25 13 18 28	10:30	P 12 P	.01	
30 31	43 53	4 P 1 P	30 30	6 A 6 A	36.5 41.5	83 66	3.1 8.0	n se	215 142	27 15	0 A 5 P		.14	2.00

^{*}Based on least time required to blow one mile.

SOL KRIMINETSKY, Observer

March, 1942	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)	20.57	20.00.100
Maximum	$30.57 \\ 29.24$	30.96, 1904
Mean semi-daily	30.00	28.47, 1914
Range	1.34	2.10, 1914; 0.85, 1915
Air Temperature, in degrees F.	*.01	2.10, 1011, 0.00, 1010
Highest 63, 8th	63.8	79.5, 1907, 1921
Lowest	6.1	-7.5, 1906
Mean	34.4	
Range	57.6	
Highest mean daily		
Lowest mean daily	40.0	
Mean maximum	43.2	
Mean minimum	25.4	
Least daily range 5, 6th, 14th, 17th		
Precipitation, in inches		
Precipitation	3.70	7.13, 1899; 0.12, 1915
Snow	7.47	27, 1899; 0, 1921
Maximum precipitation in 24		21, 1000, 0, 1021
hrs 1.82, 3rd		
Number of days with .01 or more 13	11	17, 1890, 1913, 1936; 3, 1915
Wind, in miles		
Total movement	5944	8182, 1896; 3006, 1905
Greatest daily movement 436, 10th		
Least daily movement 61, 16th		
Mean hourly velocity	7.7	10 1000 1000 1011
Maximum velocity48, 3 rd	31.2	48, 1932, 1939, 1941
Wind, direction	****	
Prevailing direction NW	WNW	
North, days		
Northeast, days		
Southeast, days		
South, days4		
Southwest, days2		
West, days		
Northwest, days		
Weather		
Mean relative humidity, percent68.0	64.2	
Mean cloudiness, percent	51.8	
Number of clear days11	11	22, 1924; 3, 1901
Number of fair days	10	17, 1900; 3, 1941
Number of cloudy days	10	21, 1901; 1, 1915
Number hours bright sunshine205	199	292, 1924; 93, 1901
Percent of possible hours of bright		
Percent of possible hours of bright	53.6	
Percent of possible hours of bright sunshine	53.6	

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1941.

The weather during March was warmer than usual, with a large amount of rain. The mean temperature during the month was 39.4 degrees which is 5 degrees warmer than usual. The highest temperature was 63 degrees on the 8th and the lowest 26 degrees on the 27th and 29th. This is the highest minimum for March ever recorded at this station. The nearest was 21 degrees in March, 1889 and the next 19 degrees in 1903. The College Pond was clear of ice on March 20. This is about 10 days earlier than usual. The domestic heating load for the month was 792 degree-days. The normal is 950 degree-days. The total heating load for the season to the end of March is 5480 degree-days, while the normal for this period is 5932 degree-days.

The total precipitation during the month was 7.89 inches. This is the greatest precipitation ever recorded at this station during March. The nearest to this record was 7.13 inches in 1899. The normal for March is 3.70 inches. The total for the three-month period beginning at New Year's is 13.09 inches while the normal for this period is 10.50 inches. The total snowfall during the month was 7.25 inches which is slightly less than normal. The total snowfall to the end of March was 23.75 inches while the normal for that period is 45.62 inches. During the winter of 1936-1937 the total snowfall was 23.25 inches.

There were 205 hours of bright sunshine which exceeds the normal value by 6 hours.

Meteorological Series

Bulletin No. 640

April 1942

Meteorological Observations

FOR

APRIL

1942

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Т	empe:	rature		ity	gs.		Wind		P	recipitat	ion	
		imum		imum		Mean Relative Humidity	Number of Hours Bright Sunshine	iling	nent	Max. Miles per Hour*				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relati	Numbe	Prevailing Direction	Total Movement	Max. per H	Began	Ended	Water	Snow
1 2 3 3 4 5 6 7 8 9 10 11 12 13 14 4 15 16 17 18 19 20 21 22 23 24 5 27 28 29 3 31	453 655 44 84 146 150 767 670 22 33 4 52 8 8 7 7 8 2 5 7 8 4 8 8 1 8 1 8 1 8 1 8 1 8 1 8 1 8 1 8	6 P 4 P 3 P 4 P 3 P 12 M 4 P 5 P 0 A 2 P 5 P 5 P 3 P 11 A 3 P 4 P 4 P 4 P 4 P 4 P 4 P 6 P 6 P 6 P 6	32 30 47 40 31 36 44 40 35 30 26 44 44 39 38 35 38 39 36 38 49 36 36 49 49 49 49 49 49 49 49 49 49 49 49 49	6 A 5 A 6 A 12 P 6 A 12 P 11 P 3 A 12 P 10 P 6 A 6 A 12 P 1 A 12 P 1 A 12 P 6 A 6 A 6 A 12 P 6 A 5 A 4 A 6 A 6 A	38.5 46.5 56.0 47.0 49.0 57.5 34.5 34.5 35.0 38.5 40.5 43.0 57.0 44.5 48.5 56.5 60.0 63.5 53.0 52.0 65.5	6339773585444588834688632871484345555944	1.4 12.4 9.3 6.6 5.2 12.9 4.8 8.9 2.0 1.0 5.4 3.9 9.3 6.6 2.8 6.9 6.0 4.1 5.4 13.5 13.5 13.8 12.5 10.3 12.9	w sw n n s s nw nw nw sw se nw n n n se se nw nw nw sw se se se s s nw	237 163 132 135 115 142 113 223 132 105 111 274 162 106 181 149 191 273 261 163 56 87 85 119 123 161 123 113	22 20 8 13 13 17 13 20 10 15 11 22 20 15 17 22 20 15 15 17 22 20 14 12 21 15 15 16 17 17 18 19 19 10 10 10 10 10 10 10 10 10 10	2 A 8 P 3 A 0 A 6 P 12:30 P 8 P 0 A	5 A 12 P 8 P 8 A 6:30 P 1 P 12 P 2 A	.05 .15 .43 .11 .01 .01 .08 .12 T	4.00 1.00 T T

^{*}Based on least time required to blow one mile.

April 1942	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum 30.36 9th	30.48	30,71, 1911
Minimum 29.55, 11th, 12th	29.38	28.99, 1907, 1929
Mean semi-daily 30.013	29.99	
Maximum 30,36, 9th Minimum 29.55, 11th, 12th Mean semi-daily 30.013 Range 0.81	1.11	1.52, 1930; .72, 1919
Highest85, 25th	79.4	90, 1941
Lowest 26, 14th	22.0	8.5, 1923
Mean 49.7	45.7	52, 1921; 41.2, 1926
Range59	57.4	
Highest mean daily 65.5, 30th		
Mean maximum 62.1	56.8	
Mean minimum 37.1	34.8	
Greatest daily range 44, 24th, 25th	01.0	
Air Temperature, in degrees F. Highest 85, 25th Lowest 26, 14th Mean 49.7 Range .59 Highest mean daily 65.5, 30th Lowest mean daily 34, 10th Mean maximum 62.4 Mean minimum 37.1 Greatest daily range 44, 24th, 25th Least daily range 8, 10th	1	
Precipitation, in inches		
Precipitation 0.96	3.35	6.89, 1929; .55, 1941
Precipitation 0.96 Snow 5.00 Maximum precip. in 24 hrs. 0.43, 10th	2.17	11, 1891; 0, 1910, 1934
Maximum precip. in 24 hrs 0.43, 10th		
Number of days with .01 or more	11	18, 1909; 3, 1892
Wind, in miles		
Total movement	5404	8208, 1908; 3853, 1917
Greatest daily movement 274, 12th		
Moon howly volocity 50, 25rd	7.5	
Least daily movement 56, 23rd Mean hourly velocity 6.4 Maximum velocity 25, 19th	31.4	40, 1935, 1938, 1940
Wind, direction	31.4	10, 1703, 1700, 1740
Wind, direction North, days 9 Northeast, days 0 East, days 0 Southeast, days 6 South, days 4 Southwest, days 3	WNW	
North days	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	
Northeast, days 0		
East, days 0		
Southeast, days		
South, days +		
Southwest, days 3 West, days 1		
West, days		
Weather	61.6	
Mean relative humidity, percent	51.8	75,1901; 34, 1925, 1927
Number of clear days	11	23, 1941; 3, 1898, 1902
Number of fair days 11	9	18, 1915; 2, 1894, 1901
Number of cloudy days 6	10	22, 1901; 1, 194;
Number hours bright sunshine 245	220	329, 1941; 103, 1901
Percent of possible hours of bright		
Number of cloudy days 6 Number hours bright sunshine 245 Percent of possible hours of bright sunshine 60.9 Last snow April 11	54.7	N 11 1010 N 11 11 100
Last snow April H	Apr. 15	Mar. 14, 1910 ; May 11, 1907

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1941.

The weather during April was dry and warm, quite similar to April 1941 although not quite as extreme. The mean temperature for the month was 49.7 degrees, the normal being 45.7 degrees. Last year, April had a mean temperature of 52.1 degrees. The domestic heating load of the month was 458 degree-days, compared to a normal of 579 degree-days. The total heating load for the season to the end of April is 5938 degree-days whereas the normal for this period is 6511 degree-days. The total precipitation for the month was .96 of an inch. Last year only .55 of an inch of rain fell in April. Five inches of snow fell during the month. The normal rainfall for April is 3.35 inches. The total precipitation since January 1st is now 14.05 inches, the normal for these four months being 13.85 inches. There were 245 hours of bright sunshine. The normal sunshine for April is 220 hours.

Dr. J. K. Shaw, Research Professor of Pomology, makes the following comments: "The past winter was not severe, yet there was some injury to fruit plants. The hardier varieties of peaches came through with plenty of live fruit buds, but those of tender varieties, such as Elberta, were all killed. The amount of killing will vary with different orchards; therefore, the peach crop will be small and spotty depending on the site and varieties. Raspberry canes show considerable winter injury, especially among the more tender varieties. As last year, this season is now far advanced and with little rainfall. Vegetation is at just about the same stage as it was a year ago. Last year was, on the whole, very favorable to fruit plants. If, as last year, we escape a too prolonged drought and spring frosts, fruit crops, except peaches and raspberries, should be good. Apple trees are beginning to bloom plentifully. Frost injury to blossoms and young apples is unusual in this area, but the earlier the bloom, the more danger of frost injury. McIntosh trees are especially full of bloom."

Meteorological Series

Bulletin No. 641

May 1942

Meteorological Observations

FOR

MAY

1942

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Т	empe:	rature		ity	S		Wind			recipitat	ion	
	Maximum		Minimum			Humid	f Houn	, bo		es				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 22 23 24 25 27 28 29 30 31 31 31 31 31 31 31 31 31 31 31 31 31	78 77 78 86 81 64 68 50 61 56 63 64 70 85 77 77 77 77 77 63 66 72 69 62 75 75 74 74 77 77 77 77 77 77 77 77 77 77 77	1 P 2 P 4 P 12 M 3 P 5 P 4 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P 1	51 48 55 55 58 43 35 44 45 49 42 31 43 55 57 56 55 57 56 57 57 56 57 57 57 57 57 57 57 57 57 57 57 57 57	12 P 5 A 6 A 12 P 11 A 6 A 6 A 12 P 5 A 6 A 5 A 5 A 5 A 5 A 7 A 5 A 6 A 12 P 4 A 5 A 8 A	64.5 62.5 70.5 69.5 53.5 51.5 52.5 52.5 56.5 71.0 65.0 67.0 65.5 63.5 59.0 66.0 64.5 61.5 62.5 63.0 64.5 64.5 64.5 64.5	49 48 73 65 54 56 94 79 84 63 50 87 78 87 79 80 85 89 82 70 66 68 77	10.2 11.3 7.6 9.3 14.2 10.8 0.0 7.3 1.3 9.0 13.8 6.3 11.0 8.9 7.4 4.9 0.3 4.0 4.2 1.8 1.2 3.6 10.6 12.0 10.8 11.0 10.0 10.0 10.0 10.0 10.0 10	sw s s s s nw sw n se se se se se se se se se se se se se	72 169 119 169 265 82 103 64 96 180 82 171 123 106 154 195 140 86 77 108 135 119 72 69 48 149 48 149 48 149 98 128 195 90	13 13 12 15 22 8 7 9 8 17 11 25 8 13 13 15 17 17 15 9 7 9 9 15 11 11 10 10 10 10 10 10 10 10 10 10 10	12-30 p 8 p 0 A 6 p 0 A 2-30 A 1 p 8-30 A	1-30 P 12 P 6 P 13 P 1 A 11 A 9 P 2 P 9 A	.02 .03 .72 T .43 .11 1.19 T T .27 .01 .01 T	

^{*}Based on least time required to blow one mile.

Sol Kriminetsky, Observer

May 1942	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum 30.31, 6th	30.39	30,62, 1936
Minimum 29.77, 23rd	29.51	29.10, 1938
Minimum 29.77, 23rd Mean semi-daily 30.050	29.96	,
dange 0.54	0.88	
Air Temperature, in degrees F.		
Highest 87, 30th Lowest 31, 11th Mean 61.4 Range 56 Highest mean daily 74, 30th 157, 74	86.1	94.5, 1896, 1911
Lowest 31, 11th	31.2	24.0, 1900
dean61.4	57.1	
Range 50	54.9	
lighest mean daily	İ	
Lowest mean daily		
Mean maximum		
Mean minimum		
Mean minimum 50.9 Greatest daily range 37, 29th Least daily range 6, 7th, 21st		
Precipitation, in inches		
Precipitation 2.98	3.60	7.44, 1931; .48, 1903
	0.50	7.44, 1931, .40, 1900
Snow	1	
Number of days with .01 or more 10	12	20, 1901; 5, 1903
W/:1 ::1		
Total movement 3799 Greatest daily movement 265, 5th Least daily movement 48, 26th	4504	5946, 1907; 2180, 189
Greatest daily movement 265, 5th		
Least daily movement 48, 26th		
Mean hour velocity 5.1	6.1	
Maximum velocity25, 12th	28.8	45, 1933
Wind, direction		
Prevailing directionSSW_	W	
North, days/		
Northeast, days 0		
Cast, days		
South days 7		
Southwest days		
West days		
1		
Weather		
	60.7	
Mean relative humidity, percent	52.1	70, 1901, 1902; 30, 192
Number of clear days	11	20, 1923, 1941; 0, 192
Number of fair days 10	11	17, 1907, 1926; 5, 192
Number of clear days	9	20, 1927; 2, 194
Number hours bright sunshine 258	252	338, 1941; 137, 192
Percent of possible hours of bright		
sunshine 57	55.6	
I nunder and lightning		
4th, 12th, 13th, 31st	31 11	June 8 1032, Apr 22 100
Last frost	May 14	June 8, 1932; Apr. 23, 190

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1941.

The weather during May was normal in most respects except that the temperature was higher than usual. The mean temperature for the month was 61.4 degrees, while the normal is 57.1 degrees. This mean temperature has been exceeded only twice in May since 1889, in 1911 and 1918. In 1911 the mean was 62.5 degrees and in 1918, 62.2 degrees. The highest temperature was 87 degrees on the 30th and the lowest, 31 degrees on the 11th. These are normal ranges of temperature for May. The last frost of the month occurred on May 11. The domestic heating load for the month was 144 degree-days as compared to a normal of 287 degree-days. The total heating load for the season is now 6082 degree-days compared to a normal of 6798 degree-days. In other words, the heating load during the current year has been only 90 percent of the normal value.

The total precipitation for the month was 2.98 inches, the normal being 3.60 inches. This brings the total precipitation since January 1 to 17.03 inches which is 0.42 less than normal for that period. There were 258 hours of bright sunshine which is 6 hours higher than the average. There were no high winds during the month.

Dr. J. K. Shaw, research pomologist, makes the following statement regarding the progress of vegetation: "Weather during May was seasonable and fruit plants have made normal progress. The bloom of apple trees was very early, about the same date as last year. Bloom was very heavy. Many trees, especially McIntosh, that produced a good crop last year bloomed heavily. Conditions for pollination were good and set of young apples seems to be heavy. Some frost injury was reported from low-lying orchards but indications now point to one of the heaviest apple crops in recent years. The peach bloom was better than anticipated but varieties tender in bud show little or no bloom. Strawberries are beginning to ripen the first of June. The prospects are for a good blueberry crop but raspberries may be a short crop owing to winter injury to the canes."

Meteorological Series

Bulletin No. 642

June 1942

Meteorological Observations

FOR

JUNE

1942

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Т	emper	ature		ity	S		Wind		Pı	ecipitat	ion	
	Maximum		Minimum			Humid	of Hou	bo	ıt.	les.				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max, Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 6 7 8 9 100 111 12 13 114 15 116 17 18 19 200 21 22 23 244 25 5 26 27 28 29 30	68 62 61 64 77 82 77 83 85 86 88 88 89 75 68 78 84 83 79 74 74 74 74 78 83 85 85 86 88 88 88 88 88 88 88 88 88 88 88 88	12 M 12 M 15 P 4 P 6 P 5 P 6 P 3 P 2 P 3 P 4 P 4 P 6 P 7 P 6 P 7 P 6 P 7 P 6 P 7 P 6 P 7 P 6 P 7 P 6 P 7 P 6 P 7 P 6 P 7 P 6 P 7 P 6 P 7 P 6 P 7 P 6 P 7 P 6 P 7 P 6 P 7 P 6 P 7 P 6 P 7 P 7 P 8 P 8 P 8 P 8 P 8 P 8 P 8 P 8 P 8 P 8	50 49 48 56 57 50 62 64 53 54 67 71 68 61 59 57 50 59 59 51 56 55 56 56 57 68 61 57 68 61 57 69 69 69 69 69 69 69 69 69 69 69 69 69	12 P 2 A A 4 A A 12 P 5 A A 7 A A 2 A A 4 A A 12 P 5 A A 5 A A 2 A A 5 A A 2 A A 5 A A 5 A A 5 A A 6 A A 5 A A 6 A A 5 A A 6 A A 7 A A A 7 A A 7 A A 7 A A 7 A A 7 A A 7 A A 7 A A 7 A A 7 A A 7 A A 7 A A A 7 A A 7 A A A 7 A A A 7 A A A 7 A A A 7 A A A A	59.0 55.5 54.5 60.0 67.0 69.5 73.5 69.0 71.5 66.0 61.5 65.0 67.0 72.0 68.0 69.0 63.5 62.0 69.5 62.0 66.0 67.0 75.0 75.0	74 69 91 91 65 63 86 77 65 63 87 72 75 84 68 63 92 72 74 68 63 59 72 72 74 65 66 68 66 72 74 68 66 72 74 66 66 72 74 66 74 74 74 74 74 74 74 74 74 74 74 74 74	4.0 3.4 0.0 0.0 12.8 15.2 15.2 15.2 15.2 10.0 9.2 7.8 8.4 11.0 13.6 8.1 15.3 0.0 13.6 12.7 10.0 7.0 10.0 11.7	sw se nw s nw s s n s s s s s s s s e n n e nw w nw w	64 84 53 64 172 83 85 84 59 113 135 1147 68 80 337 201 1101 58 76 108 38 63 63 141 139 64 67 57 62 177	7 11 6 11 25 10 12 11 9 10 12 11 15 11 20 28 18 12 20 15 15 18 12 20 15 18 20 17 18 20 19 19 10 10 10 10 10 10 10 10 10 10 10 10 10	3 P 0 A 2 A 6 P 9 A 0 A 5 P 4 P 5:30 P 7 A 2 A	12 P 6 A 6 P 12 P 12 P 4 A 7 P 11 P 2 P 6 :30 P	1.40 T .05	

^{*}Based on least time required to blow one mile.

J. F. Hanson, Observer

June 1942	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum30.32, 2nd	30.30	30.54, 1889
Minimum29.58, 14th	29.55	29.24, 190.
Mean semi-daily29.963	29.95	27.24, 1702
Range0.74	0.75	
Air Temperature, in degrees F.		
Highest90, 13th	91.2	101, 1919
Lowest48, 3rd	40.0	34, 189
Mean67.5	65.7	0,, 10,
Range42.	51.3	
Highest mean daily79.5, 12th		
Lowest mean daily54.5, 3rd		
Mean maximum78.0		
Mean minimum57.0		
Greatest daily range34, 19th		
Least daily range8, 4th		
Precipitation, in inches		
Precipitation3.63	3.75	9.68, 1922; 0.76, 1908
Snow		
Maximum precipitation in 24 hrs1.40, 14th		
Number of days with .01 or more12	11	17, 1922; 4, 1908
Wind, in miles		
Total movement3043	3585	4571, 1908; 1409, 1900
Greatest daily movement337, 15th	0.00	1011, 1200, 1102, 120
Least daily movement38, 21st		
Mean velocity4.2	5.0	
Maximum velocity28, 15th	24.6	48, 1939
Wind, direction		
Prevailing directionSW	WSW	
North, days3		
Northeast, days0		
East, daysl	İ	
Southeast, days4		
South, days11		
Southwest, days2		
West, days3		
Northwest, days6		
Weather		
Mean relative humidity, percent73.6	66.9	
Mean cloudiness, percent	51.1	71, 1903; 28, 190
Number of clear days13	10	22, 1908, 1941; 1, 192
Number of fair days11	12	23, 1912; 3, 194
Number of cloudy days	8	22, 1903; 1, 192.
Number hours bright sunshine285	257	362, 1908; 102, 190,
Percent of possible hours of bright		
sunshine	54.0	
Thunder and lightning7th, 12th, 13th,		
14th, 18th, 19th, 23rd		

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1941.

June was a good growing month in Amherst. Rainfall was very near normal. Temperature was higher than usual and there was more sunshine than normal. The total precipitation for the month was 3.63 inches while the normal is 3.75 inches. This brings the total precipitation since January 1 to 20.66 inches, the normal for this period being 21.20 inches. The mean temperature for the month was 67.5 degrees compared to a normal of 65.7 degrees. The highest temperature was 90 degrees on the 13th. The domestic heating load for June was 42 degree-days the normal being 68 degree-days. The heating load for the season is 6124 degree-days while the normal is 6866 degree-days. Amherst residents should therefore have burned 11 percent less fuel than normal during the past winter.

There were 285 hours of bright sunshine. The normal for June is 257 hours. There was slightly less wind than normal with a total movement of 3043 miles. There were no high winds, the highest being 28 miles per hour on the 15th,

Dr. J. K. Shaw, research pomologist, makes the following comment regarding vegetation: "Progress of vegetation during June has been normal. There has been sufficient rain to keep plants growing well. Nights have been rather cool for best growth of warm weather crops. The hay crop is good and grass has started well towards a second crop. There has been a heavy June drop of Mc-Intosh, but in most orchards enough apples remain to make a good crop. Considerable scab appears in some orchards, but those sprayed thoroughly and at the right time are quite free from it. The strawberry crop was generally disappointing. There was some rot, and berries failed to set and develop well. Peaches and raspberries will be short due to winter cold. Blueberries promise a good crop."

Meteorological Series

Bulletin No. 643

July, 1942

Meteorological Observations

FOR

JULY

1942

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Т	`empe	rature		ity	s.		Wind		Pr	ecipitati	on	
	Maximum		Minimum			Humid	of Hou	5.c		les				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 4 5 6 7 8 9 10 11 1 13 14 15 16 17 18 19 20 1 22 2 24 2 5 2 6 2 7 8 2 9 3 3 1	80 72 84 87 88 88 88 88 74 77 78 68 82 88 89 91 77 78 89 89 89 87 67 87 67 88 87 77 77 77 77 78 88 89 89 89 89 89 89 89 89 8	5 P 4 P 4 P 1 P 4 P 2 P 2 P 11 A 3 P 2 P 2 P 2 P 3 P 2 P 2 P 3 P 4 P 2 P 3 P 4 P 3 P 3 P 4 P 1 P 4 P 3 P 3 P 1 P 4 P 3 P 1 P 4 P 3 P 1 P 4 P 3 P 1 P 4 P 3 P 1 P 4 P 3 P 1 P 4 P 3 P 1 P 4 P 3 P 1 P 4 P 3 P 1 P 4 P 3 P 1 P 4 P 3 P 1 P 4 P 3 P 1 P 4 P 3 P 3 P 1 P 4 P 1 P 4 P 1 P 1 P 1 P 1 P 1 P 1	65 64 65 66 65 66 66 66 66 66 66 66 66 66 66	3 A 7 A A 7 A A 12 P A 6 A A 12 P A 1	72.5 68.5 74.0 75.5 74.0 63.5 63.0 62.0 64.0 60.5 71.5 74.5 67.0 69.0 73.0 72.5 68.0 72.5 68.0 72.5 65.5 75.5 75.5 75.0 75.0	82 90 78 72 72 74 72 53 64 66 90 67 68 83 57 66 88 77 65 57 66 71 71 75 99 86 80 88 87 86	3.5 0.3 9.8 9.2 11.5 9.6 15.2 9.7 15.1 12.0 0.4 15.1 15.0 14.9 10.7 11.5 6.4 7.5 8.0 11.2 12.5 14.9 12.0 12.8 9.6 9.9 0.0 9.9 12.0 9.6 12.0 12.0 12.0 12.0 12.0 12.0 12.0 12.0	se s nw se w nw se se nw nw nw nw nw me nw w w w w w se se nw nw nw se se nw nw nw sw nw sw nw sw nw sw se nw nw sw se nw nw sw se nw nw sw se nw nw sw se nw w sw n s	148 45 96 125 70 120 139 113 83 87 85 146 74 167 255 58 64 90 119 88 68 47 45 94 49 52 111 57	13 8 10 13 11 10 25 14 18 17 10 10 22 8 7 10 13 11 10 10 22 11 11 10 10 10 10 10 10 10 10	7 P 0 A 0 A 4 P 12 M 2 A 4:30 P	12 P 12 P 3 A 1 A 7:30 P 1 P 1 P 5:30 P	1.13 .45 .26 T 8 .54 .05 .43	

^{*}Based on least time required to blow one mile.

July 1942	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum30.31, 20th	30.27	30.50, 1892
Minimum29.65, 6th	29.59	29.27, 193,
Mean semi-daily29.970	29.96	29.27, 193.
Range0.66	0.68	0.97, 1892; 0.47, 1938
Air Temperature, in degrees F.	00	1 2, 11, 1500
Highest	93.9	104, 1911
owest47, 9th	46.4	40, 1890, 1898
Iean	70.8	74.7, 1921; 66.3, 1893
Range40	47.7	
Highest mean daily80, 19th		
Lowest mean daily60.5, 11th		
lean maximum81.3		
Iean minimum59.6		
reatest daily range33, 13th, 14th		
east daily range3, 27th		
Precipitation, in inches		
recipitation4.95	4.10	14.51, 1897; 0.70, 1929
now		
laximum precipitation in 24 hrs1.32, 27th		
Tumber of days with .01 or more11	11	20, 1915; 4, 192-
Wind, in miles		
otal movement2950	3422	5097, 1909; 1109, 189-
Greatest daily movement255, 15th	- ·- <u>-</u>	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
east daily movement45, 2nd, 25th		
lean hour velocity4.0	4.6	
laximum velocity25, 6th, 14th, 15th	28.7	44, 193
Wind, direction		
revailing directionW	SW	
Corth, days1	5,11	
Vortheast, days1		
Cast, days0		
Southest, days7		
South, days3		
bouthwest, days2		
Vest, days7		
Forthwest, days10		
Weather		
Jean relative humidity, percent73.6	68.4	
lean cloudiness, percent59	50.9	69, 1914; 31, 192-
Sumber of clear days14	10	22. 1923 : 0. 1913
Number of fair days11	14	24, '09, '21; 5, '89, '23, '28
·		36
Sumber of cloudy days6	7	18, 1889; 0, 1910
Sumber hours bright sunshine291	268	371, 1910; 180, 1931
Percent of possible hours of bright		
sunshine63	58	
hunder and lightning1st, 4th, 6th, 19th		

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1941.

The weather during July was very near normal in most respects. The mean temperature for the month was 70.5 degrees, the normal being 70.8 degrees. The highest temperature was 93 degrees on the 19th and the lowest 47 on the 9th. The rainfall during the month was 4.95 inches which is .85 more than normal. The greater part of this rain fell in two storms: on the 1st, 2nd and 3rd, 1.84 inches was recorded and on the 27th, 1.32 inches. The total rainfall since January 1 is now 25.61 inches while the normal for this period is 25.30 inches. There were 291 hours of sunshine compared to a normal of 268 hours. The mean relative humidity was 73.6 percent while the normal for July is 68.4 percent.

Dr. J. K. Shaw reports as follows: "The progress of vegetation during July continued to be normal. Timely rains have favored growth and practically all fruit plants grew well. Ripening dates are yet a week or more earlier than usual, reflecting the early bloom. The peach crop will be rather light especially that of varieties having fruit buds tender to winter cold. Apples, especially McIntosh, continue to promise crops above normal. The blueberry crop is larger than usual and pears and plums seem about normal."

Meteorological Series

Bulletin No. 644

August, 1942

Meteorological Observations

FOR

AUGUST

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Т	emper	ature		ity	S	,	Wind		Pre	cipitatio	on	
	Maxi	cimum Minimum			Humid	f Hou:	bn	t	les					
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 22 23 24 25 27 28 29 30 30 30 31 31 31 31 31 31 31 31 31 31 31 31 31	85 79 82 75 74 81 67 75 85 83 86 82 85 84 86 90 91 84 71 71 73 78 79 76 80 84	4 P 2 P 3 P 2 P 3 P 4 P 2 P 3 P 4 P 2 P 3 P 2 P 3 P 2 P 10 A 1 P 4 P 6 P 3 P 4 P 2 P 3 P 4 P 4 P 6 P 3 P 4 P 2 P 3 P 4 P 4 P 4 P 4 P 4 P 4 P 4 P 4 P 4	64 55 59 52 52 48 47 50 61 62 64 68 71 72 70 65 54 42 41 47 47 47 47 47 47 47 47 47 47 47 47 47	5 A 12 P 15 A A 5 A A 6 A A 12 P 12 P 12 P 12 P 12 P 12 P 5 A A 6 A A 12 P 15 A A 6 A A 12 P 5 A A 6 A A A 12 P 5 A A 12 P 5 A A 6 A A 12 P 5 A A 6 A A 12 P 5 A A 6 A A 12 P 5 A A 6 A A 12 P 5 A A 6 A A 12 P 5 A A 6 A A 12 P 5 A A 6 A A 12 P 5 A A 6 A A 12 P 5 A A 6 A A 12 P 5 A A 6 A A A 12 P 5 A A 6 A A A 12 P 5 A A 6 A A A 12 P 5 A A 6 A A A 12 P 5 A A 6 A A A A 12 P 5 A A 6 A A A A 12 P 5 A A 6 A A A A A A A A A A A A A A A A	73.5 67.0 70.5 63.5 63.5 61.0 64.0 65.5 64.0 68.5 71.0 69.0 77.5 78.0 77.5 78.0 72.0 78.5 80.5 74.5 59.5 56.5 57.0 63.5 67.5 67.5 67.5 67.5	70 75 48 58 63 62 65 73 92 88 87 78 81 90 92 84 85 76 76 79 70 63 64 70 70 70 79 80 71	10.8 6.5 14.4 11.9 14.3 10.7 13.6 11.3 0.0 5.0 7.0 5.8 8.4 13.5 13.5 9.5 13.5 10.5 10.5 13.5 13.5 13.5 13.5 13.5 13.5 13.5 13	w se w nw sw se se w w nw s sw se se w w nw nw nw nw se se w w sw sw se se w sw sw se w sw sw se w sw sw se w sw sw se se nw se se ne e s	124 666 189 175 104 94 92 47 72 465 38 40 104 158 101 50 43 117 83 75 144 143 92 40 91 104 108 78	13 7 22 22 10 14 6 8 7 10 10 10 12 8 6 6 7 7 8 9 9 14 15 17 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9	6:00 A 12 M 0 A 7:30 P 0 A	10 p 12 p 8 p 12 p 1 :30 A	.50 T .31 1.82 .02 .03 T T	

^{*}Based on least time required to blow one mile.

August 1942	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum30.45, 20th	30.32	30.50, 1934
Minimum29.71, 2nd, 3rd	29.61	28.87, 1930
Mean semi-daily30.048	30.00	
Range0.74	0.71	
Air Temperature, in degrees F.		
Highest	91.6	100, 1918
Lowest41, 26th	43.4	37, 1894, 1908
Mean	68.6	34, 1940
Range50	48.1	
Highest mean daily80.5, 22nd		
Lowest mean daily 56.5, 25th Mean maximum 80.2		
Mean minimum 57.6		
Greatest daily range		
Least daily range		
Precipitation, in inches		
Precipitation2.93	4.08	8.40, 1928; .31, 1894
Maximum precipitation in 24 hrs2.05,		, , , , , , , , , , , , , , , , , , , ,
13th, 14th		16 1003 1033 4 1000
Number of days with .01 or more6	11	16, 1892, 1933; 4, 1899
Wind, in miles		1 271 1010 1020 1004
Total movement2790	3127	4,271, 1910; 1.920, 1894
Greatest daily movement189, 3rd		
Least daily movement32, 7th	4.2	
Mean velocity3.8 Maximum velocity2, 3rd, 4th	4.2 22.7	40, 1941
	22.7	10, 1511
Wind, direction Prevailing directionWSW	SW	
North, days	311	
Northeast, days1		
East, days1		
Southeast, days6		
South, days4		
Southwest, days4		
West, days7		
Northwest, days7		
Weather		
Mean relative humidity, percent73.9	70.2	
Mean cloudiness, percent52	49.7	67, 1901; 27, 1923
Number of clear days16	9	23, 1941; 0, 1915, 1929
Number of fair days12	13	25, 1912; 3, 1939
Number of cloudy days3	227	18, 1901, '28; 2, 1910, '23
Number hours bright sunshine288	237	332, 1941; 152, 1915, 1929
Percent of possible hours of bright sunshine	55.2	
Thunder and lightning14th, 16th, 17th, 22nd		

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1941.

The weather during August was very near normal in most respects. There was more sunshine and less rain than usual. The mean temperature for the month was 68.9 degrees which is 0.3 of a degree higher than normal. The highest temperature was 91 degrees on the 22nd and the lowest 41 degrees on the 26th. The rainful during the month was 2.93 inches, the normal for August being 4.08 inches. The total rainfall since January 1st is 28.54 inches compared to a normal of 29.38 inches for this period. There were 288 hours of bright sunshine while the normal for August is 237 hours. The mean relative humidity was 73.9 percent. The normal for August is 70.2 percent. Total wind movement for the month was 2790 miles which is 337 miles less than normal.

Dr. J. K. Shaw, research pomologist, makes the following report on the progress of vegetation: "The season continues to be nearly normal in all respects. There has been sufficient rain to permit vegetation to progress well. Towards the end of the month there were cool nights that noticeably improved the color of McIntosh and other red varieties of the same season, though a clear atmosphere that is likely to accompany cool nights may be the real reason. Rainfall was deficient towards the last of the month, but at this season there is little danger of injury to crops from dry weather. The promise of good crops of all fall-ripening fruit continues."

Meteorological Series

Bulletin No. 645

September 1942

Meteorological Observations

FOR

SEPTEMBER

1942

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Т	emper	ature		ity	S.		Wind		Pre	ecipitati	o n	
	Maxi	mum	Min	imum		Humidity	r of Hours Sunshine	ър	t	es				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative	Number of Bright Su	Prevailing Direction	Total Movement	Max, Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30	87 90 87 69 77 70 77 72 61 76 65 67 78 79 88 89 88 89 88 88	3 P 2 P 3 P 2 P 4 P 2 P 4 P 1 P 4 P P 1 P 4 P P 1 P 4 P P 1 P 4 P P 1 P 4 P P 1 P 4 P P 1 P 4 P P 1 P	51 54 57 57 57 57 57 57 57 57 57 50 43 42 56 60 60 55 55 66 63 63 63 63 43 44 40 41 62 42 43 44 44 46 47 48 48 48 48 48 48 48 48 48 48 48 48 48	5 A 5 A 6 A 12 P A 6 A A 12 P A 12 P A 12 P A 12 P A 12 P A 12 P A 12 P A 12 P A 12 P A 5 A A 12 P A 5 A A 12 P A 5 A A 12 P A 6 A A 5 A A 7 A 12 P A 6 A A 5 A A 7 A 12 P A 6 A A 6 A A 7 A 12 P A 6 A A 6 A A 7 A 12 P A 6 A A 6 A A 7 A 12 P A 6 A A 7 A 12 P A 6 A A 7 A 12 P A 6 A A 6 A A 7 A 12 P A 6 A A 6 A A 7 A 12 P A 6 A A 6 A A 7 A 12 P A 6 A A A 6 A A A 6 A A A 6 A A 6 A A 6 A A 6 A A 6 A A 6 A A 6 A A 6 A A 6 A A 6 A A 6 A A 6 A A 6 A A 6 A A 6 A A A 6 A A A 6 A A A 6 A A A 6 A A A 6 A A A A A 6 A	69.0 72.0 63.0 63.5 56.5 59.5 64.0 65.5 62.5 63.5 64.0 71.5 77.5 73.5 62.0 58.0 62.0 57.0 58.0 62.0 57.0 58.0 64.5 44.5 46.5	70 78 80 91 64 74 85 88 87 85 69 74 80 86 84 71 72 75 77 74 70 89 64 68 67	13.2 13.1 8.3 0.0 12.3 13.0 11.5 0.0 0.5 12.6 10.5 4.5 9.6 0.0 10.5 3.2 12.0 8.7 0.0 11.5	sw sw nw s nw nw se nw se n s w se s w se s w s s s s m s s s n w s s n w s s n w s s n w s s s n w s s s s	52 68 73 47 89 64 82 51 55 56 98 31 39 79 228 213 86 62 23 38 38 38 38 38 251 58 43	5 5 9 13 3 14 19 10 9 10 7 7 9 6 13 11 1 7 18 4 11 1 8 20 13 25 8 13 25 5 13 6	1 A 11 A 0 A 8 P 2:30 P 5 A 3:30 P 1:30 P 5 A	3 P 12 P 9 P 9:30 P 10 P 7:30 P 3:30 P 11 A	.08 1.45 .04 .03 .21 .03 .18 T	

^{*}Based on least time required to blow one mile.

J. F. Hanson, Observer

September, 1942	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum30.56, 29th	30.45	30.65, 1924
Minimum 29.64, 27th	29.57	28.41, 1938
Mean semi-daily30.059	30.06	20.11, 1500
Range0.81	.88	1.99, 1938; .57, 1910
Air Temperature, in degrees F.		
Highest90, 2nd	87.7	97, 1895, 1929
Lowest28, 30th	33.2	24.5, 1914
Mean	61.7	
Range62	54.5	
Highest mean daily77.5, 16th		
Lowest mean daily43.0, 29th		
Mean maximum74.4		
Mean minimum51.0		
Greatest daily range37, 30th		
Least daily range		
Precipitation, in inches		
Precipitation3.94	4.26	14.55, 1938; .52, 1914
Maximum precipitation in 24 hrs1.69, 27th		
Snow		
Number of days with .01 or more11	10	16, 1933; 3, 1903
Wind, in miles		
Total movement2707	3271	4,686, 1896; 1,414, 1894
Greatest daily movement251, 28th		
Least daily movement31, 17th		
Mean velocity3.8	4.5	
Maximum velocity28, 28th	25.9	80, 1938
Wind, direction		
Prevailing directionSW	WSW	
North, days2		
Northeast, days0		
East, days0		
Southeast, days5		
South, days		
Southwest, days5		
West, days5		
Northwest, days7		
Weather		
	737	
Mean relative humidity, percent77.8	73.6	70 1031 25 1000 101
Mean cloudiness, percent53	49.9	70, 1934; 27, 1908, 1914
Number of clear days16	10	70, 1934; 27, 1908, 1914 19, 1932; 2, 1907, 1928
Number of fair days5	10	19, 1908; 3, 1889
Number of cloudy days9	10	20, 1934; 3, 1941
Number hours bright sunshine224	201	255, 1916; 106, 193-
Percent of possible hours of bright		
Percent of possible hours of bright sunshine60.1	54.9	
Percent of possible hours of bright	54.9	

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1941.

The weather during September was nearly normal in most respects. The mean temperature was only one degree higher than normal for the month but was unusually high during the third week. The highest temperature was 90 degrees on the 2nd and the lowest, 28 degrees on the 30th. The first frost was on the 29th while the normal date for the first frost is September 21.

The domestic heating load for the month was 138 degree-days. The normal for September is 149 degree-days.

The rainfall during the month was 3.94 inches, the normal for September being 4.26 inches. The total precipitation since January 1 is 32.48, while the normal is 33.62 inches. There were 224 hours of bright sunshine compared to a normal of 201 hours. The total wind movement was only 2707 miles while the normal is 3271 miles. The highest wind velocity was 28 miles per hour on the 28th.

Dr. J. K. Shaw, research pomologist, comments as follows: "Alternate cool and warm periods prevailed during September. A warm, moist period came in the midst of the McIntosh harvest and was probably the principal cause of an unusually heavy pre-harvest drop. The McIntosh crop was even larger than expected; the fruit was of good size and fairly well colored; it matured a little earlier than usual. The below freezing temperature of the 29th and 30th may have injured the keeping quality of apples still on the trees, but no serious harm is expected. All tender vegetation was completely killed."

Meteorological Series

Bulletin No. 646

October 1942

Meteorological Observations

FOR

OCTOBER

1942

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23' 31'' N. Longitude, 72° 31' 48'' W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Т	empei	rature		ity	S		Wind		Pre	cipitati	on	
	Махі	inum	Min	imum		Humidity	of Hour	ρū	ı t	*es				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative I	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 22 23 24 25 26 27 28 29 30 31 31 31 31 31 31 31 31 31 31 31 31 31	68 63 64 66 72 63 71 73 72 66 68 62 72 64 65 55 73 68 63 62 54 94 95 96 97 97 97 97 97 97 97 97 97 97 97 97 97	2 P 3 P 2 P 1 P 2 P 1 P 3 P 3 P 4 P 4 P 4 P 1 P 2 P 2 P 1 P 2 P 2 P 1 P 2 P 2 P 2 P 1 P 2 P 2 P 1 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2	40 38 32 55 45 40 36 46 46 46 46 37 40 38 35 55 47 40 39 40 30 55 55 47 40 30 30 55 55 55 47 40 40 40 40 40 40 40 40 40 40 40 40 40	4 A A 12 P 6 A A 4 A A 12 P 7 A A 12 P 6 A A 12 P 3 A A 12 P 3 A A 12 P 12 P 3 A A 12 P 12 P 3 A A 12 P 12 P 3 A A 12 P 13 A A 15 A A 6 A A 6 A A 6 A A 6 A 6 A 6 A 5 A	54.0 50.5 49.0 63.5 54.0 55.5 54.5 51.0 61.0 57.5 56.0 43.0 64.0 64.0 64.0 64.0 64.0 64.0 64.0 65.5 65.5 65.5 65.5 66.0	74 53 71 77 82 73 67 64 81 71 64 67 77 84 80 72 70 83 94 86 81 71 63 71 70 70 81 70 70 70 70 70 70 70 70 70 70 70 70 70	8.9 11.7 11.7 5.6 2.0 2.9 81.5 4.8 11.3 11.3 11.3 11.3 0.9 3.1 5.6 2.2 0.0 5.3 0.5 5.2 5.2 0.0 8.5 10.5	se n nw s se se w nw w w w sw nw nw nw w se se n nw sw sw sw se se se se se se se se se se se se se	102 145 94 73 151 171 123 85 74 79 58 143 47 22 48 73 18 182 61 40 191 149 43 98 73 196 105 66 148 173	9 18 11 11 10 20 7 8 9 7 8 9 10 6 6 6 6 6 12 13 15 28 8 11 17 17 5 10 10 10 10 10 10 10 10 10 10 10 10 10	9:30 P 0 A 5:30 P 0 A 3 P	12 P 4 A 12 P 6 A 12 P	T .04 .62 T T T .78 .44 T T T .95	

^{*}Based on least time required to blow one mile.

October, 1942	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)	1	
Maximum30.66, 13th	30.53	30.68, 1929
Minimum29.42, 26th	29.42	29.00, 1926
Mean semi-daily30.085	30.06	
Range1.24	1.11	1.47, 1926; .76, 1899
Air Temperature, in degrees F.		
Highest	79.4	90.5, 1908
Lowest26, 29th	23.4	17, 1936
Mean	50.5	56.4, 1920; 43.2, 1890
Range	56.0	
Highest mean daily		
Lowest mean daily 41, 27th Mean maximum 64.9		
Mean minimum		
Greatest daily range		
Least daily range13, 16th		
Precipitation, in inches	1	
Precipitation3.27	3.29	8.81, 1911; .01, 1924
Snow	3.49	0.01, 1911, .01, 1924
Maximum precipitation in 24 hrs1.22,	i	
17th, 18th		
Number of days with .01 or more6	9	15, 1913; 1, 1897, 1924
Wind, in miles		
Total movement 3209	4074	5,467, 1910; 2,540, 1894
Greatest daily movement196, 27th		
Least daily movement22, 14th	1	
Mean velocity 4.3		
Maximum velocity28, 19th	29.5	42, 1937
Wind, direction		
Prevailing directionW) W	
North, days3		
Northeast, days0		
East, days		
Southeast, days		
Southwest, days		
West, days		
Northwest, days8		
Weather		
Mean relative humidity, percent73.7	69.0	
Mean cloudiness, percent51	48	67, 1932; 18, 1924
Number of clear days14	10	21, 1938; 1, 1911
Number of fair days8	10	17, 1924; 3, 1938
Number of cloudy days9	11	19, 1896; 3, 1924
Number hours bright sunshine195	175.0	232, '23, '38; 91, 1913
Percent of possible hours of bright	51.3	
sunshine57.2 Thunder and lightning	51.3	
First frost Sept. 29	Sept. 21	Aug. 22, '94, '95; Oct. 15, '39
11150 11050	l Septime	

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1941.

The weather during October was unusually pleasant, being warm and clear. The mean temperature for the month was 52.7 degrees as compared to a normal of 50.5 degrees. The highest temperature was 76 degrees on the 9th and the lowest, 26 degrees on the 29th. The domestic heating load for the month was 382 degree-days as compared to a normal of 456 degree-days. The total for September and October is 520 degree-days, the normal being 605 degree-days. The heating load for September and October represents 7.6% of the total heating load for the season.

The rainfall during the month was 3.27 inches which is .02 less than normal. The total rainfall since January 1st is now 35.75 inches while the normal for this period is 36.91 inches. There were 195 hours of bright sunshine during the month. The normal for October is 175 hours. There were no high winds during the month and the total wind movement was 3209 miles compared to a normal of 4074 miles.

Meteorological Series

Bulletin No. 647

November 1942

Meteorological Observations

FOR

NOVEMBER

1942

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23' 31'' N. Longitude, 72° 31' 48'' W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be αddressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Tempe	rature		ity	s.		Wind		Pre	cipitatio	on	
	Maximur	n Mir	imum		fumid	of Hours unshine	.	_	es				
Day	Degrees Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hour Bright Sunshine	Prevailing Direction	Total Movement	Max, Miles per Hour	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8	58 1 47 2 46 3 52 3	39 P 32 P 32 P 26 P 21	12 P 5 A 11 P 12 P 7 A	52.0 45.0 39.5 36.0 36.5	82 72 80 51 67	1.0 2.9 2.9 10.2 8.4	w s n n s	141 137 188 214 74	32 14 22 20 10	1 A 0 A	8 a 10 a	1.12	
6 7 8 9 10 11 12 13 14 15	56 2 43 4 54 2 63 4	P 41 P 33 P 32 P 30 P 35 A 29 P 27 M 28 P 21 P 17	12 P 12 P 6 A 6 A 1 A 12 P 1 A 12 P 12 P 7 A	49.0 44.5 37.5 42.0 49.0 36.5 33.5 24.5 28.0	79 77 81 79 57 65 65 47	0.0 7.5 0.0 5.5 0.0 2.5 2.5 4.0 4.5 7.0	n nw sw	127 59 32 72 290 271 103 260 362 166	12 7 9 8 40 18 20 32 25 11	4:30 r	8 r 8 a	.24 T .09	T
16 17 18 19 20	58 3 60 3 58 4 68 2 43 1 37 2	P 27 P 25 P 36 P 36 P 36 P 27 P 20	6 A 7 A 12 P 7 A 0 A 12 P 12 P	41.0 41.5 48.0 47.0 52.0 35.0 28.5	61 76 87 87 68 61	3.0 7.7 5.3 9.3 2.5 0.0 9.5	se w nw nw w n	123 52 101 67 131 137 157	9 15 14 10 14 15	0 A 4 A	11 A	.41 .52	
21 22 23 24 25 26 27 28 29 30	37 12 46 12 49 2	P 15 P 28 P 37 P 43 A 30 P 28 P 24 P 27	12 P	28.0 32.5 41.5 46.0 36.5 31.0 29.0 35.0	73 93 92 84 75 59 73	9.5 0.0 0.0 0.0 3.0 9.3 3.0 9.0	W HW SC HW W HW	33 99 277 128 199 320 89 138	5 17 22 11 22 22 9 10	2 A 0 A 10 F 0 A	12 r 12 r 12 r 12 r 10 x	.83 1.45 .01 .16	Т

^{*}Based on least time required to blow one mile.

November, 1942	Normal	Extremes
Barometer, in inches		-
(Readings reduced to sea level)	20 50	20.97 10.23
Maximum	30,59 29,30	30.87, 1932 28.73, 1904
Minimum 29.39, 13th Mean semi-daily 30.048	30.05	20.70, 1504
Range 1.22	1.29	1.84, 1904; .94, 1905
Air Temperature, in degrees F.		
Highest 68, 20th	66.0	75 , 1924 , 1938
.owest 15, 23rd	12.9	-4, 19:8
Jean 38.8		44.1, 1931; 33.6, 1901
Range 53	53.0	
Tighest mean daily 52, 1st, 20th		
cowest mean daily 24.5, 14th	1	
Mean maximum 48.5		
Mean minimum = 29.4	1 1	
Range 53 Highest mean daily 52, 1st, 20th Lowest mean daily 24.5, 14th Mean maximum 48.3 Mean minimum 29.4 Greatest daily range 33, 17th Least daily range 6, 26th, 28th		
Precipitation, in inches		
Precipitation, in inches Precipitation 6.07 Snow T	3.41	8.64, 1927; .63, 1917
Snow	2.34	13.50, 1938
Maximum precipitation in 24 hrs. 1.45, 25th	2.04	10.30, 1200
Number of days with .01 or more 11	9	22, 1921; 2, 1904
Wind, in miles		
Fotal movement4547Greatest daily movement362, 14thLeast daily movement32, 8thMean hourly velocity6.3Maximum velocity40, 10th	4531	5,978, 1906; 2,589, 1889
Greatest daily movement 362, 14th		
Least daily movement 32, 8th	1	
Mean hourly velocity 6.3	6.3	
Maximum velocity 40, 10th	30.3	44,1978
Wind, direction		
Prevailing direction WNW	MXM	
North, days		
Northeast, days		
Prevailing direction WNW North, days		
Southeast, days		
Southwest days		
West, days	3	
Southeast, days 5 South, days 5 Southwest, days 1 West, days 8 Northwest, days 16		
Weather		
Mean relative humidity, percent Mean cloudiness, percent Number of clear days Number of fair days Number of fair days	70.6	
Mean cloudiness, percent	55.1	72, 1900, 1927; 34, 1917
Number of clear days 10	6	15.1903 : 1, 1900, 1911, 1927
Number of fair days	9	16, '12; 4, '89, '30, '32, '35
Number of cloudy days 13	15	24, 1927; 9, 1905, 1917
	121	182, 1903; 66, 1927
Percent of possible hours of bright	1 (1)	
sunshine 44. Thunder and lightning	41.3	
First snow Nov 2	Nov 6	Oct. 10, 25 · Nov. 27, 31
Thurder and lightning First snow Nov. 22	Nov. 6	Oct. 10, '25; Nov. 27, '

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1941

We had a large rainfall during November, the total for the month being 6.07 inches. The normal for the month is 3.41 inches. The total precipitation since January 1 is 41.82 inches while the normal for this period is 40.32 inches. Only traces of snow fell during the month. There were traces of snow in the air on the 12th and 13th but a heavier trace on the 27th is reported as the first snow of the season. The normal date for the first snowfall is November 6 while the latest date for the first snowfall previously observed is November 27.

In spite of 11 days of rain the total hours of bright sunshine was 130 which is well above the average of 121 hours.

The mean temperature during the month was 38.8 degrees, the normal being 38.9 degrees. The lowest temperature was 15 degrees on the 23rd. The domestic heating load was 784 degreedays which is the normal for November. The total heating load for the season to the end of November is now 1304 degree-days or 19 percent of the normal heating load of the whole season. The normal heating load to the end of November is 1389 degreedays. The mean relative humidity was 72.2 percent. The highest wind velocity was 40 miles on the 10th.

Meteorogical Series

Bulletin No. 648

December 1942

Meteorological Observations

FOR

DECEMBER

1942

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23' 31'' N. Longitude, 72° 31' 48'' W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Т	emper	ature		ity	S	,	Wind		Pre	cipitati	n	
	Max	imum	Min	imum		Humidity	of Hour	to.	+	ies				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 22 33 34 25 26 27 28 29 30 30 30 30 30 30 30 30 30 30 30 30 30	48 53 31 34 36 38 37 27 40 43 40 27 25 35 27 23 34 45 35 36 47 27 28 27 33 33 32 33 33 32 33 33 34 35 36 37 37 37 37 37 37 37 37 37 37 37 37 37	11 P P 6 A A 2 P P 1 P 2 P P 2 P P 1 P 1 P 1 P P 1 P P 1 P P 1 P P 1 P P 1 P P 1 P P 1 P P 1 P P 1 P P 1 P P 1 P P 1 P P 1 P	23 22 20 21 29 28 26 19 17 25 29 27 7 7 7 24 18 -7 7 7 -8 -15 -15 -8 30 32 25 11 27 28 30 27	8 A 12 P 1 A A 7 A A A A 11 P 12 P 4 A A 1 A 12 P 7 A A 5 A A 12 P 12 P 8 A A 7 A A 6 A A 12 P 12 P A 12 P 12 P A A 12 P 12 P	35.5 37.5 27.5 32.5 32.5 32.5 32.0 22.0 32.5 36.0 29.5 22.5 8.0 0.0 16.5 32.5 34.0 32.5 34.0 32.5 34.0 32.5 34.0 32.5 34.0 32.5 34.0 32.5 34.0 32.5 34.0 32.5 34.0 32.5 34.0 32.5 34.0 32.5 34.0 32.5 34.0 32.5 34.0 32.5 34.0 32.5 34.0 32.5 34.0 32.5 36.0 36.0 36.0 36.0 36.0 36.0 36.0 36.0	90 63 61 62 61 68 78 72 88 83 85 62 70 54 58 62 70 65 73 90 82 81 65 72 95 95 95	0.3 3.1 2.3 7.7 9.2 9.2 9.2 9.1 1.0 0.0 0.5 1.0 0.5 7.6 9.0 7.6 9.0 3.4 1.4 0.0 8.2 9.0 0.0 0.0 0.0 7.2	nw w w w w w nw nw nw nw nw se nw sw nw nw nw nw nw nw nw nw nw nw nw nw nw	54 179 340 264 257 245 205 104 125 92 105 77 134 244 158 245 245 116 240 222 57 173 95 81 46 177 94 97 101 135 360	28 48 28 25 22 25 18 10 20 9 9 11 28 32 8 40 10 15 8 6 6 22 8 10 7 7 12 10 10 10 10 10 10 10 10 10 10 10 10 10	7 A 8 P 0 A 5 P 10 P 0 A 0 A	12 P 7 A 12 P 4 A 12 P 12 P 12 P 12 P 12 P	.31 .04 .08 .14	1.50 T T 14.00 T .50 1.00

^{*}Based on least time required to blow one mile.

J. F. HANSON, Observer

December, 1942	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.90, 27th Minimum 28.57, 2nd Mean semi-daily 30.027	30.66 29.27 30.06	30.96, 1889 28.85, 1915
Range 2.33	1.38	1.78, 1895; 1.01, 1892
Air Temperature, in degrees F.		25 5 Aug 2
Highest 53, 2nd Lowest -15, 20th, 21st Mean 24,8 Range 68 Highest mean daily -5, 20th Lowest mean daily -5, 20th Mean maximum 32.5 Mean minimum 17.1 Greatest daily range 42, 22nd Least daily range 4, 24th, 29th	54.3 -1.9 27.5 56.2	65.5, 1908 -22.5, 1917 36.9, 1891; 17.1, 1917
Precipitation, in inches		
Precipitation 6.03 Snow 7.00	3.39 8.50	7.77, 1901; .87, 1896 26.50, '02; t, 1891
Maximum precipitation in 24 hrs. 2.33, 30th Number of days with .01 or more	10	17, 1902; 4, 1892
Wind, in miles	1540	
Total movement 4967 Greatest daily movement 360, 31st Least daily movement 46, 25th	4710	6,694, 1925; 3,239, 1918
Mean hourly velocity 6.7 Maximum velocity 48, 2nd	6.3 31.4	48, 1938
Wind, direction		
Prevailing direction WNW North, days 3 Northeast, days 0 East, days 4 Southeast, days 1 Southwest, days 1 West, days 9 Northwest, days 13	WNW	
Weather	60.1	
Mean relative humidity, percent 71.7 Mean cloudiness, percent 58 Number of clear days 14 Number of fair days 2 Number of cloudy days 15 Number hours bright sunshine 135 Percent of possible hours of bright	69.4 54.9 8 9 14 128	71, 1929; '39, 1919 15, 1890; 2, 1922, 1933 16, '09; 4, '89' 30, '31, '36, '38 23, '33; 7, '09, '23 172, 1896; 63, 1933
sunshine 47.7 Thunder and lightning 2nd	45.2	

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1941.

ANNUAL SUMMARY

Annual 1942	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum	30.81	31.05, 1920
Minimum 28.57, Dec. 2	28.95	28.41, 1938
Mean semi-daily	30.01	,
Range 2.33	1.85	2.47, 1938; 1.38, 1933
Air Temperature, in degrees F.		
Highest 93, July 19 Lowest 95, Dec. 20, 21	95.7	104, 1911
Lowest15, Dec. 20, 21	-12.2	-26, 1904
Mean48.8	47.4	49.8, 1921; 44, 1904
Range 108 Highest mean daily 80.5, Aug. 22 Lowest mean daily -5, Dec. 20 Mean maximum 59.1	107.8	
Highest mean daily 80.5, Aug. 22		
Lowest mean daily —5, Dec. 20	57.8	
Mean minimum 38.6	37.6 36.5	
Mean minimum 38.6 Greatest daily range44, Apr. 24, 25	1.0.5	
Least daily range3, July 27		
Precipitation, in inches		
Precipitation 47.85	43.70	59.00, 1938; 30.68, 1908
Precipitation 47.85 Snow 33.75	47.78	89.00, 1893; 24.50, 1919
Maximum precipitation in 24 hrs. 233, Dec. 30		
Number of days with .01 or more 112	124	144, 1902; 96, 1924
Wind, in miles		
Total movement	52,223	63,571, 1908; 36,257, 1894
Greatest daily movement 475, Jan. 29		
Least daily movement 13, Jan. 1		
Mean hourly velocity	5.8	
Maximum velocity 48, Mar. 3, Dec. 2	39.5	80, 1938
Wind, direction		
Prevailing direction W North, days 50	W	
North, days 50		
Northeast, days 4		
East, days 4		
Southeast, days		
Southwest, days 28		
West, days 56		
Northwest, days		
Weather		
Mean relative humidity, percent 70.9	67.6	
Mean cloudiness, percent 55	51.7	60, '98, '01, '02; 41, '08, '24
Mean cloudiness, percent55Number of clear days165Number of fair days102	116	217, 1941; 59, 1927
Number of tair days102	123	182, 1912; 64, 1936
Number of cloudy days 98	126 2,353	179, '01, '02; 71, 1910, 1941
Number hours bright sunshine 2605 Percent of possible hours of bright	4,000	3038, 1941; 1864, 1902
sunshine 58.5	52.8	
Last snow Apr. 11	April 15	Mar. 14, '10; May 11, '07
First snow Nov. 27	Nov. 6	Oct. 10, '25; Dec. 13, '41
First snow Nov. 27 Last frost May 11	May 14	Oct. 10, '25; Dec. 13, '41 Apr. 23, '04; June 8, '32
First frost Sept. 29	Sept 21	Aug. 22, '94, '95 ; Oct. 13, '09

Meteorological Series

Bulletin No. 649

January 1943

Meteorological Observations

FOR

JANUARY

1943

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23' 31'' N. Longitude, 72° 31' 48'' W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

-		Т	emper	ature	<u>-</u>	ty	ys.		Wind		Pre	cipitati	on	
	Max	imum	Min	imum		Humidity	f Hour nshine	bo		es				,
Day	Degrees	Тіте	Degrees	Time	Mean	Mean Relative 1	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 4 5 6 6 7 8 9 10 11 12 13 14 15 16 6 17 18 19 20 21 22 23 24 25 26 27	33 29 32 22 18 22 29 28 28 30 34 35 32 41 29 29 31 35 16 12 20 32 32 32 32 41 29 29 29 29 29 29 29 20 20 20 20 20 20 20 20 20 20 20 20 20	11 P P 3 P P 1 P P P P P P P P P P P P P	27 14 20 10 7 2 4 4 -4 -3 4 0 21 11 14 9 24 19 19 19 19 10 8 8 25 31 8 18 19 19 19 19 19 19 19 19 19 19 19 19 19	7 A A 8 A A 12 P 12 P 0 A 12 P 11 P 12 P 12 P 12 P 0 A 12 P 11 P 12 P 0 A 12 P 11 P 12 P 12 P 12 P 12 P 12 P 12	30.0 21.5 26.0 16.0 12.5 12.0 16.5 12.0 17.0 28.0 17.0 28.0 29.5 24.5 24.0 24.0 29.5 12.5 6.5 12.5 6.0 30.0 30.0 35.0 32.0 32.0 32.0 32.0 32.0 32.0 32.0 32	82 50 59 57 61 67 74 75 80 75 80 95 87 88 68 61 75 80 95 96 96 96 96 96 96 96 96 96 96 96 96 96	0.8 9.1 8.87 6.3 9.2 9.3 9.3 5.0 7.7 4.8 2.0 0.0 0.8 8.4 6.6 4.5 8.3 4.1 0.0	s w w nw nw w w w nw nw s nw nw nw nw nw nw nw nw nw nw nw nw nw	151 211 158 144 268 209 121 76 46 41 76 202 2246 203 83 31 244 233 344 233 140 175 344 233 140 116 83 63 181 181	25 25 17 28 25 22 17 7 25 3 17 22 22 22 17 10 10 32 28 21 17 10 32 11 11 11 11 11 11 11 11 11 11 11 11 11	0 A 11 P 0 A 6 A 5 A 4 P	12 P 4 A 12 M 8 P 4 P	T .54 T .02 .07 .17 .70 .10 T .04 .01	T 4.50 T 0.25 1.50 T .50 .25
28 29 30 31	24 35 23 42	12 M 4 P 3 P 2 P	17 19 19 21	4 A 0 A 1 A 0 A	20.5 27.0 21.0 31.5	83 66 84 86	0.0 8.4 2.9 7.1	nw nw nw nw	173 224 158 147	17 15 10 12	11 A 2 P	10 Р 12 Р	.93 .17	9.00 3.25

^{*}Based on least time required to blow one mile.

J. F. HANSON, Observer

January, 1943	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.65, 29th, 30th Minimum 29.36, 15th Mean semi-daily 30.066 Range 1.29	30.70 29.20 30.07 1.50	31.00, 1927 28.55, 1913 2.18, 1913; 0.97, 1896
Air Temperature, in degrees F.	1.50	2.10, 1910, 0.97, 1090
Highest 42, 31st Lowest -4, 8th Mean 21.4 Range 46 Highest mean daily 35, 25th Lowest mean daily 6.5, 21st Mean maximum 29.1 Mean minimum 13.6 Greatest daily range 34, 11th Least daily range 3, 18th	51.1 -7.1 24.2 58.2	66.0, 1932 -26.0, 1904 34.2, 1913; 13.9, 1918
Precipitation, in inches		
Precipitation 2.92 Snow 19.25 Maximum precipitation in 24 hrs. 0.93, 28th	3.61 13.32	7.15, 1898; 1.07, 1896 33, 1898, 1923; 1.50, '08
Number of days with .01 or more 11	11	15, '93, '20; 4, 1901
Wind, in miles		
Total movement 4874 Greatest daily movement 344, 20th Least daily movement 41, 10th Mean hourly velocity 6.6 Maximum velocity 32, 19th	5055 6.8 32.7	7770, 1908; 2896, 1895 47, 1938
Wind, direction	34.7	77, 1930
Prevailing direction WNW North, days 0 Northeast days 0	WNW	
East, days 0 Southeast, days 2 South, days 4 Southwest, days 0 West, days 6 Northwest, days 19		
Weather		
Mean relative humidity, percent 73.3 Mean cloudiness, percent 63 Number of clear days 15 Number of fair days 9 Number of cloudy days 7 Number hours bright sunshine 174 Percent of possible hours of bright sunshine 59.2 Thunder and lightning 59.2	70.1 55 9 9 13 137 46.7	78, 1932; 37, 1908 18, 1941; 2, 1914 18, 1890, 1926; 3, 1916 22, 1923, 1931; 4, 1920, 1939 214, 1920; 74, 1932

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1942.

The weather during January was more severe than usual. The mean temperature during the month was 21.4 degrees, while the normal mean temperature for January is 24.2 degrees. It should be noted, however, that the mean temperature during January, 1940 was 18.5 degrees and in 1941 it was 21.2 degrees. The lowest temperature was —4 degrees on the 8th and the highest 42 degrees on the 31st. The domestic heating load was 1352 degreedays compared to a normal of 1265 degree-days. The total heating load for the season to the end of January is 3902 degree-days which is slightly more than 2 percent higher than the normal of 3817 degree-days for this period. The heating load for the current year to the end of January is 57 percent of the normal annual load.

There was a total snowfall of 19.25 inches during the month. The normal snowfall during January is 13.32 inches. The total precipitation during January is 3.61 inches. The mean humidity during the month was 73.3 percent compared to a normal of 70.1 percent. The total wind movement during the month was 4874 miles, the normal being 5055 miles. The highest wind velocity was 32 miles per hour on the 19th.

Meteorological Series

Bulletin No. 650

February 1943

Meteorological Observations

FOR

FEBRUARY

1943

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23' 31'' N. Longitude, 72° 31' 48'' W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be αddressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Т	`empei	ature		Humidity	S		Wind		Pre	ecipitati	o n	
	Maximum		Minimum		mum		f Hournshine	bo		es				
Day	Degrees	Time	Degrees	Time	Меап	Mean Relative I	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8 9 10	40 30 36 39 45 35 38 31 31 26 36	2 P 2 P 2 P 8 P 4 P 3 P 11 A 1 P 4 P 12 P	26 16 1 28 27 24 30 25 17 10 23	0 A 12 P 7 A 0 A 12 P 4 A 12 P 12 P 12 P 7 A 12 P	33.0 23.0 18.5 33.5 36.0 29.5 34.0 28.0 24.0 18.0 29.5	74 61 70 88 71 95 88 58 65 83 86	4.8 10.0 9.5 0.0 10.1 0.0 5.0 6.0 10.3 5.4 0.0	W W Se Se W NW SW NW NW NW	166 361 87 204 129 62 189 356 259 67 92	28 28 11 15 18 11 18 32 18 12 32	5 A 8 A 7 A 0 A	11 A 8 P 12 P 1 A	.14 .03 .37 .03 T	.50
11 12 13 14 15 16 17 18 19 20 21 22 23 24 25	28 25 25 -8 16 27 23 37 55 52 55 58	1 P 10 P 3 A 2 P 4 P 5 P 4 P 3 P 4 P 3 P 4 P 3 P	15 8 0 -19 -23 -7 7 18 23 32 25 28	12 P 6 A 12 P 8 A 7 A 4 A 8 A 6 A 12 P 7 A	21.5 16.5 12.5 -13.5 -3.5 10.0 15.0 27.5 39.0 42.0 40.0 43.0	66 88 64 59 61 75 58 61 71 64 68 73	9.0 0.3 10.5 10.5 10.6 5.5 10.0 6.5 9.2 10.8 9.0 10.9	W NW W NW NW S NW SE NW S SE	275 72 247 352 69 114 190 84 127 103 48 82	20 14 28 35 8 18 17 8 13 10 7	10 A 0 A	12 P 3 A	.32 .04	4.00
24 25 26 27 28	47 35 28 34 45	12 M 11 A 4 P 1 P 12 P	33 17 11 19 23	12 P 12 P 7 A 7 A 0 A	40.0 26.0 19.5 26.5 34.0	83 60 72 72 63	0.5 11.4 5.4 11.2 6.6	se nw nw w s	147 265 98 158 269	20 20 10 18 28	12 м	3 P	.07 T	Т

^{*}Based on least time required to blow one mile.

February, 1943	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
	30.63	31.05, 1920
Maximum 30.50, 18th Minimum 29.27, 7th	29.24	28.56, 1895
Mean semi-daily 29.947 Range 1.23	30.04	
	1.38	1.89, '00, '08; .88, '13 '31
Air Temperature, in degrees F.		
Highest 58, 23rd Lowest -23, 16th Mean 25,1	50.5	65.0, 1930
Lowest	-7.4	-22.5, 1918
Range 81	23.7 58	
Highest mean daily 43, 23rd	30	
Highest mean daily 43, 23rd Lowest mean daily -13.5, 15th		
Mean maximum	32.8	
Mean minimum15.6	14.5	
Greatest daily range 39, 16th Least daily range 6, 8th		
Precipitation, in inches		0.12 1000 0.63 1001
Precipitation 1.63 Snow 5.00	3.19	8.12, 1900; 0.62, 1901
Maximum precipitation in 24 hrs. 0.63, 11th	13.99	48.75, 1893; 0.50, 1937
Number of days with .01 or more 8	10	15, '93, '20; 4, 1901
Wind, in miles		
Total movement 4672 Greatest daily movement 361, 2nd Least daily movement 48, 22nd Ween beardy reducitive 77	4776	6445, 1896; 3438, 1892
Greatest daily movement 301, 2nd		
Mean hourly velocity 7	7.1	
Mean hourly velocity 7. As 221d Maximum velocity 35, 15th	31.7	48, 1934, 1937
Wind, direction		,
Prevailing direction W North, days 0 Northeast, days 0	WNW	
North, days		
Northeast, days0		
East, days 0		
Southeast, days 6 South, days 3		
Southwest days]	
West, days		
Southwest, days		
Weather		,
Mean relative humidity, percent	66.6	•
Mean cloudiness, percent 58	50.9	66, '90, '27; 31, 1905
Number of clear days 15	10	19, 1941; 2, 1927
Number of clear days 15 Number of fair days 8 Number of cloudy days 5	8	16, 1920; 2, 1936 16 1891; 2, 1936
Number of cloudy days Number hours bright sunshine 199		16,1894; 2, 1920 221, 1924; 110, '27, '38
Percent of possible hours of bright	1.00	221, 1927, 110, 27, 30
sunshine 67.2	53.4	
Thunder and lightning		

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1942.

The weather during February was extreme in many respects. There was less snow than usual but alternate periods of warm and cold weather produced more icy walks than normal. The snowfall during the month was 5 inches, while the normal snowfall during February is 13.99 inches. Only three years since 1889 has there been less snow in February. The total precipitation during the month was 1.63 inches, the normal being 3.19 inches. The mean temperature for the month was 25.1 degrees compared to a normal of 23.7 degrees. The highest temperature was 58 degrees on the 23rd. This has been exceeded only three times since 1889. The lowest temperature was -23 degrees on the 16th. This is the lowest temperature ever recorded at this station in February. Only twice have lower temperatures been recorded since 1889, -26 degrees in January 1904 and -23.5 in January 1907. The mean temperature on February 16 was -13.5 degrees which is the lowest mean daily temperature recorded at this station. The domestic heating load for the month was 1117 degree-days, the normal for February being 1165 degree-days. The total heating load for the season to the end of February is 5019 degree-days. The normal for this period is 4982 degree-days. The heating load for the season is therefore less than one percent higher than normal. The total wind movement was 4672 miles, slightly less than normal, and the highest wind velocity was 35 miles per hour on the 15th. There were 199 hours of bright sunshine, the normal for February being 158 hours. The mean relative humidity was 71.3 percent compared to a normal of 66.6 percent.

Meteorological Series

Bulletin No. 651

March 1943

Meteorological Observations

FOR

MARCH

1943

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23' 31'' N. Longitude, 72° 31' 48'' W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

-		Т	emper	ature		ity	r.s		Wind		Pre	cipitatio	n									
	Maximum		Minimum		Minimum		Minimum		Minimum		Minimun			Humid	f Houn	bo		es				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow								
1	37 29	1 Р 1 Р	26 21	8 A 8 A	31.5 25.0	58 55	9.2 8.0	w	136 198	22 20				Т								
3	29 21	3 P	7 1	12 P 4 A	18.0 11.0	81 61	$\frac{0.0}{11.3}$	nw	259 232	25 25 25 25	2 л	6 Р	.17	2.25								
1 2 3 4 5 6 7 8	40 31 28 24	1 P 8 P 3 A 2 P	20 18 15 8	0 A 8 A 12 P 7 A	30.0 24.5 21.5 16.0	54 95 57 54	11.3 ().0 11.5 11.5	se nw w nw	315 177 284 180	25 15 28 22	6 а	10 р	.84	5.25								
$\frac{9}{10}$	26 44	4 Р 9 Р	8 -7 18	7 A 3 A	9.5 31.0	68 86	11.6 1.0	W NW	55 67	6 14				Т								
11 12 13	49 50 38	4 P 3 P 1 P	40 32 28	2 A 12 P 3 A	44.5 41.0 33.0	92 84 83	2.5 2.0 0.0	se nw nw	124 144 69	10 17 5	6 р 0 а	12 P 8 A	.12 .26	Т								
13 14 15	50 56	4 P 4 P	25 27	5 A 7 A	37.5 41.5	67 68	11.8 11.8	w se	87 119	14 17				1								
16 17	40 37 60	0 A 7 P 3 P	31 30 32	8 P	35.5 33.5 46.0	95 95 66	0.0 0.3 12.0	nw nw	114 77 218	8 11 22	5 A 4 A	8 P 5 P	.62 .19 T									
18 19 20	45 48	3 P 2 P 5 P	33 32	5 A 4 A 2 A	39.0 40.0	60 84	0.5 3.8	se nw	105 106	14 25	5 P 0 A	12 P	.83									
21 22	40 41	3 P 2 P	29 30	2 A 7 A 7 A	34.5 35.5	68 38	9.6 10.6	11W 11W	299 222	20 25												
23 24 25	40 50 54	4 P 5 P 3 P	26 24 41	5 A 6 A 3 A	33.0 37.0 47.5	38 45 50	11.1 9.7 11.8	nw s w	384 119 277	28 14 25												
21 22 23 24 25 26 27 28	68 53	4 P 1 P	30 38	6 A 8 A	49.0 45.5	47 41	11.7 9.8	SW DW	137 221	18 25												
28 29 30	44 40 34	2 P 3 P 9 P	26 20 24	12 P 6 A 0 A	35.0 30.0 29.0	40 39 72	12.5 12.5 0.0	nw nw se	360 155 78	25 22 8				Т								
31	38	4 P	31	2 A	34.5	85	0.6	se	62	9	0 а	12 м	.03	1								

^{*}Based on least time required to blow one mile.

Joseph T. Jodka, Observer

March, 1943	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 31.05, 9th Minimum 29.43, 6th Mean semi-daily 30.164	30.57 29.24 30.00	30.96, 1904 28.47, 1914
Range 1.62	1.34	2.10, 1914; 0.85, 1915
Air Temperature, in degrees F.	62.0	#10 # 400 # 400 A
Highest 68, 26th Lowest -7, 9th Mean 32.9 Range 75 Highest mean daily 49, 26th	63.8 6.1 34.4 57.6	79.5, 1907, 1921 -7.5, 1906
Lowest mean daily9.5, 9thMean maximum41.4Mean minimum24.4Greatest daily range38, 26thLeast daily range7, 17th, 31st	43.2 25.4	
Precipitation, in inches		
Precipitation 3.07 Snow 7.50 Maximum precipitation in 24 hrs. 84, 19th,	3.70 7.47	7.89, 1942; 0.12, 1915 27, 1899; 0, 1921
Number of days with .01 or more9	11	17, 1890, 1913, 1936; 3, 1915
Wind, in miles		
Total movement 5380 Greatest daily movement 384, 23rd Least daily movement 55, 9th Many hours religious 7, 2	5944	8182, 1896; 3006, 1905
Mean hourly velocity 7.2 Maximum velocity 28, 7th, 23rd	7.7 31.2	48, 1932, 1939, 1941, 1942
Wind, direction		
Prevailing direction WNW North, days 0 Northeast, days 0 East, days 0 Southeast, days 6 South, days 1 Southwest, days 1 West, days 5 Northwest, days 18	WNW	
Weather Mean relative humidity, percent 65.4	64.2	
Mean cloudiness, percent 51 Number of clear days 19 Number of fair days 11 Number of cloudy days 11 Number hours bright sunshine 220 Percent of possible hours of bright	51.8 11 10 10 199	22, 1924; 3, 1901 17, 1900; 3, 1941 21, 1901; 1, 1915 292, 1924; 93, 1901
sunshine 59 Thunder and lightning 17th	53.6	

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1942.

The weather during March was somewhat colder than usual but near normal in other respects. The mean temperature for the month was 32.9 degrees while the normal for March is 34.4 degrees. The lowest temperature for the month was –7 degrees on March 9. Only once has there been a lower temperature in March since the station was established in 1889. A minimum of –7.5 degrees was recorded on March 24 in 1906. The low temperature of –7 on March 9 this year was accompanied with the highest barometer ever recorded at this station, a reading of 31.05 inches, corrected to sea level. The domestic heating load for the month was 995 degree-days, the normal being 950 degree-days. The total heating load of the season to the end of March is 6014 degree-days compared to a normal of 5932 degree-days. The heating load to the end of March represents 88 percent of a normal year's total heating load.

The snowfall during the month was 7.50 inches which is very near the normal snowfall for March. The total snowfall for the season is 38.75 inches while the normal snowfall to the end of

March is 45.62 inches.

The total precipitation for the month was 3.07 inches, the normal being 3.70 inches. The total precipitation since January 1 is 8.31 inches. The normal precipitation for that period is 10.50 inches. There were 220 hours of bright sunshine compared to a normal of 199 hours.

Dr. J. K. Shaw, research pomologist, makes the following statement regarding vegetation: "The minimum temperature of the past winter was unusually low. As a result we may look for more winter injury to fruit trees than usual. The peach fruit buds in Massachusetts were practically all destroyed. More or less browning of the wood indicates wood injury. We shall have to wait until growth starts to tell how severe this will be. The more tender varieties of apples also show wood injury, while the hardier varieties show little or none. How severe this may be, also remains to be seen. The covering of snow at the time of the severe cold should have prevented injury to roots and low-growing plants. A few warm days in the latter half of the month have caused some bud swelling in early leafing trees and shrubs, but the month closed with colder weather, checking the development of vegetation."

Meteorological Series

Bulletin No. 652

April 1943

Meteorological Observations

FOR

APRIL

1943

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23' 31'' N. Longitude, 72° 31' 48'' W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Тє	mpera	ture		ty	x.		Wind		1	Precipit	ation	
	Maximum		Minimum			Iumidi	r of Hours Sunshine			60 du				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Bright Su	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5	50 52 34 47	5 P 2 P 1 P	31 27 25 19	6 A 6 A 7 A 6 A	40.5 39.5 29.5 33.0	84 75 48 35	2.6 5.5 11.0 10.2	sw w nw	62 93 450 203	6 25 32 18	11 A	2 P	T .05	
5 6 7	49 30 43	11 .v 3 P	30 19 19	12 p 7 a 6 a	39.5 24.5 31.0	61 47 27	8.5 12.9 11.5	nw nw	330 481 283	32 32 28	8 A	9 .1	.01	
8 9 10	39 47 46	2 P 2 A 1 P 2 P	28 33 28	5 A 6 A 1 A	33.5 40.0 37.0	84 55 57	1.5 7.8 8.7	11//	64 172 219	18 28 32	2 A	2 1	.09	1.25 T
11 12 13 14 15	48 50 58 41 40	12 M 12 M 2 P 4 P	27 23 37 30 23	1 A A A A A A A A A A A A A A A A A A A	37.5 36.5 47.5 35.5 31.5	44 66 48 66 54	13.2 6.7 9.3 11.8 13.3	w w w w	316 165 245 305 223	25 32 22 25	2 1 9 P 0 A	10 r 12 r 4 s	.44 .37 .23	.50
16 17 18	52 58	1 P 2 P 5 P	22 39	5 A 1 A 5 A	33.0 45.5 43.0	56 63	8.8 0.3 13.5	se se W	173 242 114	22 18 17 17	3 A	4 A	.01	Т
20 21 22 23 24 25 26 27 28 29 30	47 49	11 A 4 P 11 A 5 P 4 P	28 35 35 36 35 29	12 P 1 A 12 P 3 A 5 A	41.0 42.0 39.5 46.0 49.0	42 72 86 82 76 54	0.5 4.7 0.0 6.6 13.7	n n ne nw w	171 157 82 72 80	22 11 11 12 11	12 M 0 A 12 M 3 A	12 r 9 x 10 r 4 x	.74 .20 .26 .01	Т
24 25 26	70 72 63	3 P 4 P 2 P	39 39 48	6 A 5 A 6 A	54.5 55.5 55.5	46 58 47	11.5 8.5 13.8	se sw w	189 147 331	17 22 32			Т	
27 28 29	62 60 59	2 P 2 P 11 A 2 P	32 39 34	5 A 12 P 5 A 5 A	47.0 49.5 46.5	54 73 39	9.5 7.3 13.8	se W	149 256 247	28 39 35	4 P	12 P 3 A	.28 .52	
36	52	6 р	38	5 .	45.0	83	0.0	s	179	32	11 A	8 р	.45	

^{*}Based on least time required to blow one mile.

Joseph T. Jodka, Observer

April 1943	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level) Maximum 30.42, 18th Minimum 28.98, 14th Mean semi-daily 29.955 Range 1.44	30.48 29.38	30.71, 1911 28.99, 1907, 1929
Mean semi-daily 29.955 Range 1.44	29.99 1.11	1.52, 1930; .72, 1919
Air Temperature, in degrees F.		
Highest 72, 25th Lowest 19, 4th, 6th, 7th Mean 41.0 Range 53 Highest mean daily 55.5, 25th, 26th	79.4 22.0 45.7 57.4	90, 1941 8.5, 1923 52, 1921; 41.2, 1926
Highest 72, 25th Lowest 19, 4th, 6th, 7th Mean 41.0 Range 53 Highest mean daily 55.5, 25th, 26th Lowest mean daily 24.5, 6th Mean maximum 51.0 Mean minimum 30.9 Greatest daily range 40, 23rd Least daily range 7, 21st	56.8 34.8	
Precipitation, in inches		
Precipitation 3.66 Snow 1.75 Maximum precip. in 24 hrs. 0.94, 19th 20th	3.35 2.17	6.89, 1929; .55, 1941 11, 1891; 0, 1910, 1934
Number of days with .01 or more 14	11	18, 1909; 3, 1892
Wind, in miles		
Total movement 6200 Greatest daily movement 481, 6th Least daily movement 62, 1st Mean velocity 8.6 Maximum velocity 39, 28th	5404	8208, 1908; 3853, 1917
Mean velocity . 8.6 Maximum velocity . 39, 28th	7.5 31.4	40, 1935, 1938, 1940
Wind direction		
Prevailing direction W North, days 2 Northeast, days 1 East, days 0 Southeast, days 5 South, days 1 Southwest, days 2 West, days 10 Northwest, days 9	WNW	
Weather		
Mean relative humidity, percent 59.4 Mean cloudiness, percent 55.7 Number of clear days 10 Number of fair days 12 Number of cloudy days 6 Number hours bright sunshine 247 Percent of possible hours of bright sunshine 61.4	61.6 51.8 11 9 10 220 54.7	75, 1901; 34, 1925, 1927 23, 1941; 3, 1898, 1902 18, 1915; 2, 1894, 1901 22, 1901; 1, 1941 329, 1941; 103, 1901
Thunder and lightning 27th, 28th Last snow April 14		Mar. 14, 1910; May 11, 1907

Note: The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1942.

The weather during April was very cold with a mean temperature of 41 degrees. This is the lowest mean temperature for April recorded at this station since it was established in 1889. The normal mean temperature for April is 45.7 degrees. The lowest temperature was 19 degrees which occurred on the 4th, 6th and 7th. This is not unusually cold for this month. The domestic heating load for the month was 721 degree-days compared to a normal of 579 degree-days. The total heating load for the season is now 6735 degree-days while the normal heating load to the end of April is 6511 degree-days.

The total precipitation during April was 3.66 inches; the normal precipitation is 3.35 inches. The snowfall was 1.75 inches, the normal being 2.17 inches. The total snowfall for the past winter was 33.50 inches. The normal snowfall to the end of April is 36.95 inches. There were 247 hours of bright sunshine compared to a normal of 220 hours. The total wind movement was 6200 miles, while the normal for April is 5404 miles. The highest wind velocity was 39 miles per hour on the 28th. A very low barometer of 28.98 inches, corrected to sea level, was recorded on the 14th. This is the lowest barometer recorded at this station in April.

Dr. Shaw comments as follows: "Rarely has a month of April passed with so little advance in vegetation as in April, 1943. Buds have enlarged and grass has become green at the close of the month, but the stage of development is at least two weeks behind that of the corresponding date of last year. There is as yet no indication of severe winter injury to fruit plants. There will be no peach crop in Massachusetts. There is some tree injury, but at present there is no indication that it will prove to be severe. The wood shows some brown discoloration and the branch tips are killed in some cases. Unless there is unseasonably warm weather in early May, blooming dates of all fruits will be later than the average. This is likely to make harvesting dates later than usual, but the danger of injury to blossoms and young fruits from spring frosts will be decreased. Apple fruit buds are swelling and indicate a light bloom which would be expected considering the heavy crop of last season."

Meteorological Series

Bulletin No. 653

May 1943

Meteorological Observations

FOR

MAY

1943

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23' 31'' N. Longitude, 72° 31' 48'' W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Те	mpera	iture		ty	l «		Wind			Precipitation			
	Maximum		Minimum			Humidi	r of Hours Sunshine	bn	1	es					
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number o Bright Su	Prevailing Direction	Total Movement	Max. Miles	Began	Ended	Water	Snow	
1 2 3 4 5 6 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 29 20 20 20 20 20 20 20 20 20 20 20 20 20	44 62 55 53 68 66 81 72 55 52 71 68 72 77 79 70 62 77 77 72 68 79 75 68 72 77 78 80	2 P P 3 P P 12 M 4 P P 1 P P 3 P P 1 P P 3 P P 1 P P 3 P P 1 P P 3 P P 1 P P 3 P P 3 P P 1 P P 3 P P 3 P P 1 P P 3	32 31 42 34 37 46 24 47 48 47 46 44 44 44 47 47 49 49 49 49 48 52 51 51 51 51 51 51 51 51 51 51 51 51 51	12 P 4 A A P 5 A A A A A A A A A A A A A A A A A A A	38.0 46.5 48.5 51.0 51.5 63.5 66.0 59.5 51.0 50.5 57.0 67.0 67.0 67.0 69.5 71.5 69.5 71.0 61.5 63.0 63.0 63.0 63.0 63.0 63.0 63.0 63.0	50 44 91 62 63 65 75 48 90 94 94 95 67 67 67 67 67 67 67 67 67 67 67 67 67	14.0 12.8 0.0 4.6 13.7 4.8 14.4 0.0 0.0 12.5 14.5 14.6 4.8 7.2 7.8 8.8 14.7 1.0 0.8 14.8 15.7 14.6 15.7 14.6 15.7 16.8 16.8 16.8 16.8 16.8 16.8 16.8 16.8	nw se se nw nw nw se se nw nw se se nw nw se se nw nw se se nw sw se se se nw sw se se se se se nw nw se se se se se se se se se se se se se	459 184 208 214 223 81 146 108 109 80 57 57 65 221 139 119 164 244 78 100 100 113 192 120 120 120 121 120 121 120 121 121	32 18 17 20 28 13 17 32 18 11 12 13 28 22 20 14 39 13 10 11 18 14 11 14 14 14 16 16 17 18 18 19 19 19 19 19 19 19 19 19 19	0 A 7 P 6 P 4 P 5 A 0 A 0 A 0 A 0 A 0 A 2 A 3 P	9 P 8 P 9 P 12 M 12 P 12 P 12 P 12 P 8 A 12 P 8 A	T .78 .03 .04 .42 .08 .81 .57 .09 .655 .17 .72 .05 T		

^{*}Based on least time required to blow one mile.

Joseph T. Jodka, Observer

May 1943	Norma	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30,54, 16th Minimum 29,54, 1st Mean semi-daily 30,039 Range 1,00	30.39 29.51 29.96 0.88	30.62, 1936 29.10, 1938
Air Tomporature in dogrees F		
Highest 84, 17th Lowest 31, 2nd Mean 57.8 Range 53 Highest mean daily 71.5, 17th Lowest mean daily 38.0, 1st Mean maximum 68.5 Mean minimum 47.0 Greatest daily range 38, 15th Least daily range 4, 11th	86.1 31.2 57.1 54.9	94.5, 1896, 1911 24.0, 1900
Precipitation, in inches	3.00	5 44 4004 40 400
Precipitation 5.62 Maximum precipitation in 24 hrs. 0.92, 21st, 22nd	3.60	7.44, 1931; .48, 1903
Number of days with .01 or more	12	20, 1901; 5, 1903
Wind, in miles Total movement 4420 Greatest daily movement 459, 1st Least daily movement 51, 30th	4504	5946, 1907; 2180, 1894
Least daily movement51, 30thMean hourly velocity5.9Maximum velocity39, 17th	6.1 28.8	45, 1935
Wind, direction		
Prevailing direction W North, days 1 Northeast, days 0 East, days 2 Southeast, days 9 South, days 2 Southwest, days 3 West, days 0 Northwest, days 14	W	-
Weather	co =	
Mean relative lumidity, percent 68.5 Mean cloudiness, percent 60 Number of clear days 14 Number of fair days 11 Number of cloudy days 6 Number hours bright sunshine 274 Percent of possible hours of bright sunshine 60.5	60.7 52.1 11 11 9 252 55.6	70, 1901, 1902; 30, 1923 20, 1923, 1941; 0, 1927 17, 1907, 1926; 5, 1923 20, 1927; 2, 1941 338, 1941; 137, 1927
Thunder and lightning 3rd, 8th, 12th, 17th, 18th, 19th, 27th .ast frost May 5	May 14	June 8, 1932; Apr. 23, 1904

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1942.

The outstanding feature of May weather was the large number of rainy days, with the rainfall considerably above normal. Rain fell on 15 days whereas the normal number of days of rain during May is 12. The total precipitation during the month was 5.62 inches, the normal for May being 3.60 inches. The total precipitation since January 1st is 16.90 inches, while the normal for this period is 17.45 inches.

The mean temperature during the month was 57.8 degrees compared to a normal of 57.1 degrees. The highest temperature was 84 on the 17th and the lowest temperature 31 on the 2nd. The domestic heating load during the month was 243 degree-days, while the normal for May is 267. The total heating load for the winter season is 6978 degree-days and the normal for this period is 6778 degree-days.

There were 274 hours of bright sunshine, the normal for May being 252 hours. The total wind movement for the month was 4420 miles compared to the normal for May of 4504 miles. The highest wind velocity was 39 miles on the 17th. The last frost occurred on May 5. The normal date of last frost is May 14.

Dr. J. K. Shaw, research pomologist, makes the following statement on the progress of vegetation: "At apple blossom time, the season, as measured by the date of full bloom of the McIntosh apple, was two weeks later than those of 1941 and 1942. The progress of all vegetation has been delayed by cool weather and frequent rains. Planting has been delayed by bad weather and wet soils. Conditions have favored the development of apple scab and made control difficult. Peach bloom was entirely wanting in our orchards and nearly so in all northern peach areas. Peaches will be scarce this season. Apple bloom was light on trees that bore heavily last year. The apple crop will be less than the heavy crop of last year. Peach trees were more or less injured by the severe winter cold and other tender or half-hardy trees show some injury, but at present, no material loss of trees is indicated. There is an abundant water supply in the soil and if warm sunny weather comes soon, fruit trees will progress rapidly."

Meteorological Series

Bulletin No. 654

June 1943

Meteorological Observations

FOR

JUNE

1943

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23' 31'' N. Longitude, 72° 31' 48'' W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Te	mpera	ture		ty	s		Wind			Precipita	ation	
Day	Max	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 22 3 4 4 5 6 6 7 8 9 10 11 12 13 14 15 16 6 17 18 19 20 21 22 23 24 25 26 27 28 29 30	62 85 90 94 81 76 61 73 75 73 78 82 84 75 81 81 91 88 92 94 94 95 78 68	1 P 3 P 2 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 4	55 58 63 69 68 49 48 50 52 49 55 54 61 62 63 63 63 63 63 64 65 66 66 66 66 66 67 47 16 47 16 47 16 47 16 47 16 47 16 47 16 47 16 47 16 16 16 16 16 16 16 16 16 16 16 16 16	5 5 A A A A A 5 5 A A A A A 5 5 5 A A A A A A 5 5 5 A A A A A A 5 5 5 5 6 5 A A A A	58.5 71.5 76.5 81.5 74.5 62.5 54.5 61.0 66.5 71.0 71.5 76.0 64.5 77.5 75.5 73.0 80.0 81.5 83.0 71.0	96 87 63 70 63 88 62 64 74 66 68 75 78 71 70 72 72 75 71 76 68 74 76 78 85 77	0.0 9.3 15.1 12.8 15.2 12.9 0.0 14.6 18.7 15.2 15.3 15.1 13.7 8.5 15.1 12.2 11.5 12.3 11.9 15.3 15.3 15.3 15.3 15.3 15.3 15.3 15.3	se s nw se nw se nw se se nw sw sw sw sw sw sw sw sw sw sw sw sw sw	93 119 104 93 237 99 97 68 61 141 146 83 166 105 191 201 104 105 114 105 75 163 100 94 155 116 223	10 17 13 13 13 25 12 8 8 6 6 15 13 11 13 13 12 18 11 17 32 10 15 10 11 11 13 13 13 13 13 13 13 13 13 13 13	9:30 A 6:30 A 7 A 7 P 6:30 P 6:30 P 5:30 P 2 P	8P 8A 0:30A 5:30P 11:30P 7P 11:30P 6P 3P 8P 9:30P	.54 .05 .48 .16 .46 .28 .14 .16	

^{*}Based on least time required to blow one mile.

JUNE, 1943	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum	30.30	30.54, 1889
Minimum 29.46, 2nd	29.55	29.24, 1902
Mean semi-daily 29.942	29.95	
Range	0.75	
Air Temperature, in degrees F.	01.3	101 1016
Highest	91.2 40.0	101, 1919 34, 1891
Mean71.1	65.7	54, 1691
Range 47	51.3	
Highest mean daily 83.5, 27th	01.0	
Lowest mean daily54.5, 7th		
Mean maximum82.4		
Mean minimum60.0	i i	
Greatest daily range34, 24th		
Least daily range7, 1st		
Precipitation, in inches	275	0.69 1022 0.76 1009
Precipitation 2.38 Snow	3.75	9.68, 1922; 0.76, 1908
Maximum precipitation in 24 hrs.		
59, 1st, 2nd Number of days with .01 or more11	11	17, 1922; 4, 1908
Wind, in miles		
Total movement	3585	4571, 1908; 1409, 1906
Greatest daily movement237, 5th	0000	, , ,
Least daily movement61, 9th		
Mean hourly velocity5.3	5.0	49, 103
Maximum velocity32, 20th	24.6	48, 1939
Wind, direction		
Prevailing directionWSW	WSW	
North, days		
Northeast, days		
Southeast, days		
South, days4		
Southwest, days3		
West, days5	1 1	
Northwest, days9		
Weather		
Mean relative humidity, percent73.6	66.9	71 1002 30 100
Mean cloudiness, percent	51.1	71, 1903; 28, 1908
Number of clear days6	10 12	22, 1908, 1941; 1, 192 23, 1912; 3, 194
Number of fluir days	8	22, 1903; 1, 192.
Number hours bright sunshine 356	257	362, 1908; 102, 190
Percent of possible hours of bright	W.J.	, 2000, 100, 100
	510	
sunshine	54.0	
	54.0	

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1942.

The weather during June was unusually hot with less rain than normal. The mean temperature for the month was 71.1 degrees which is the highest mean temperature for June recorded at this station. The normal mean temperature is 65.7 degrees. The nine days of June 20th to June 28th had maximum temperatures in the nineties except for two days when the maximum was 86 and 88. The highest temperature was 95 degrees which occurred on the 26th and 28th. In spite of the high temperatures throughout the month, there were 8 days when the mean temperature was less than 65 degrees. The heating load for the month was accordingly 33 degree-days. This brings the total heating load for the season to 7011 degree-days. The normal heating load for the year is 6846 degree-days, so the past season was 2.4 percent colder than normal.

The rainfall during June was 2.38 inches compared to a normal of 3.75 inches. The total rainfall since January 1 is now 19.28 inches, the normal for this period being 21.20 inches. There was a total of 356 hours of bright sunshine compared to a normal of 257 hours.

Dr. J. K. Shaw, research pomologist, gives the following statement regarding vegetation: "The cool, wet weather of this spring came to an end for the last ten days of June and was succeeded by an unusually hot period with little or no rain. The sudden change to hot, sunny days caused shoots of succulent foliage to wilt and even show dry, dead areas. However, the injury is not enough to affect crops. Most fruit crops continued to make normal progress. The strawberry crop was a little late and below normal. Raspberries will also be light due to winter injury to the canes. The wet, cool spring favored a heavy hay crop and so far, harvesting weather has been favorable. Fruit trees show considerable wood injury from the severe cold of last winter, yet few have been entirely killed. The wood of peach trees in the nursery is almost completely dead, but most of the trees are growing vigorously. Evidently the cambium layer was not injured and plenty of rain enabled the trees to start a good recovery. The same is true of many varieties of sweet cherries, but recovery is not so good. While cold injury is rather more severe than usual, no great damage is yet apparent except for the loss of the peach crop."

Meteorological Series

Bulletin No. 655

July, 1943

Meteorological Observations

FOR

JULY

1943

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23' 31'' N. Longitude, 72° 31' 48'' W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Te	mpera	ture		ty	s		Wind]	Precipit	ation	
	Max	mum	Mini	mum		Immidi	of Hours Sunshine			SC 61				
Бау	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Bright Sum	Prevailing Direction	Fixal Movement	Max. Miles per Hour*	Began	Ended	Wat.r	wou
1 2 3 3 4 5 6 6 7 7 8 8 9 10 11 12 13 14 15 16 17 18 18 19 20 21 22 23 24 25 26 27 28 29 30 30 30 30 30 30 30 30 30 30 30 30 30	73 80 81 74 81 73 82 83 86 83 87 81 86 83 87 81 84 82 84 84 84 84 84 84	2 P P 3 P 4 P P 3 P P P P P P P P P P P P	46 49 01 66 61 58 61 65 67 72 70 68 64 53 64 55 64 64 62 66 62 61 56	5 A A A A A A A A A A A A A A A A A A A	56.50.50.5.5.5.5.6.5.0.5.0.5.0.5.0.5.0.5	8775 8775 8775 8775 8775 8775 8775 8775	15.2 15.7 2.17 16.1 16.1 16.1 17.1 17.1 17.1 17.1 18.1 18.1 19.1	nw nw se se se nw h h h k s s s s e v w se s s s e v w s e s e v s e v s e u s e s e u s e s e u s e s e u s e s e	150 110 150 150 150 150 150 150 150 150	23	4:30 f 12:30 p 10:30 x 1:30 p	9 F 12 F 6 F	.03 .03 .03 .01 .05 .06 .26 .27 1.05 .45 .06 .51 .01 .01 .01	

^{*}Based on least time required to blow one mile

July 1943	Normal	Extremes
Barometer, in inches (Readings reduced to sea level) Maximum	30.27 29.59 29.96 0.68	30.50, 1892 29.27, 1932 0.97, 1892; 0.47, 1938
Air Temperature, in degrees F. Highest 92, 10th Lowest 43, 2n.1 Mean 71.8 Range 49 Highest mean daily 81, 12th Lowest mean daily 59.5, 1st Mean maximum 83.0 Mean minimum 60.7 Createst daily range 37, 2nd, 20th Least daily range 5, 29th Precipitation, in inches	93.9	104, 1911 40, 1890, 1898 74.7, 1921; 66.3, 1891
Precipitation, in inches Precipitation 6.18 Snow 6.18 Maximum precipitation in 24 hrs. 140, 25th		14.51, 1897; 0.70, 1929
Number of days with .01 or more 16	11	20, 1915; 4, 1924
Wind, in miles Total movement 296a Gree est daily movement 213, 5th Least daily movement 39, 17th Mean hourly velocity 4.0 Maximum velocity 25, 31, t	3422 4.6 28.7	5097, 1909; 1109, 1894 44, 1936
Wind, direction Prevailing direction WSW North, days 1 Northeast, days 0 East, days 0 Southeast, days 7 South, days 7 South, days 7 South, days 7 South, days 7 Northwest, days 3 Northwest, days 11	SW	11, 1500
Wenther Mean relative humi lity, percent 83.2 Mean cloudiness, percent 70 Number of clear days 20 Number of fair days 9 Number of cloudy days 342 Number hours bright senshine 342 Percent of possible hours of bright sunshine 77 Thunder and lightning 11th 13th 20th, 21st, 22nd, 27th, 31st	10 14 7 21 268 1 58	69, 1914; 31, 1924 22, 1923; 0, 1915 24, '09, '21; 5, '89, '23, '28, '36 18, 1880; 0, 1910 371, 1910; 180, 1931

Note—The first column in the above summary gives observations made during the month. The second column gives the averages lased on observations made from 1889 to 1978, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1942.

July weather was slightly warmer than normal, with heavy showers during the latter part of the month. The mean temperature for the month was 71.8 degrees compared to a normal of 70.8 for July. It will be recalled that the mean temperature for June was 71.1 degrees. The highest temperature during the month was 92 degrees on the 10th and the lowest, 43 on the 2nd. The rainfall during the month was 6.18 inches while the normal for July is 4.10 inches. Only one other year since 1928 had greater rainfall in July. In 1938, 7.45 inches was recorded and in 1928, 6.23 inches. The total rainfall since January 1 is now 25.46 inches compared to a normal of 25.30 inches for this period. There were 342 hours of bright sunshine during the month, while the normal for July is 268 hours. Twenty days classified as "clear", nine as "fair" and two as "cloudy". Thunderstorms occurred on seven days.

Dr. J. K. Shaw of the Pomology Department makes the following report: "Vegetation made good progress during July. Dry weather during the early part of the month checked the progress of shallow-rooted plants, and excessively drained lawn soils showed dry, brown areas of grass. Frequent showers during the latter part of the month corrected this deficiency and the month closes with the soil well supplied with moisture. A hail storm did considerable damage to tobacco and fruit crops in a limited area in Amherst.

"The strawberry crop was below normal and winter injury to raspberry canes has reduced the crop. The blueberry crop is about normal. Apple scab on unsprayed McIntosh is very severe owing to frequent rains while the foliage was expanding. It was well controlled on properly sprayed trees but there is likely to be more scabby McIntosh than usual."

Meteorological Series

Bulletin No. 656

August, 1943

Meteorological Observations

FOR

AUGUST

1943

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Te	прета	ture		ty	v		Wind		I	Precipita	ation	
Day	Maxies O	imum Jim	Mini	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Fotal Mov.ment	Max. Miles per Hour*	Began .	Endad	Water	Snow
1	87 92 89 81 75 81 75 82 82 82 82 86 80 89 78 75 75 82 83 83 83 85 76	5 P 2 P 4 P 0 A A P 12 M 4 P 12 M 4 P 12 M 4 P 12 M 5 P 12 M 5 P 12 M 5 P 15 P 15 P 15 P 15 P 15 P 15 P 15	57 62 70 69 61 61 60 64 65 75 55 563 60 45 748 45 55	4 A A A A A A A A A A A A A A A A A A A	72.0 77.0 79.5 75.0 68.0 63.0 71.5 71.5 64.0 69.5 70.0 72.5 68.0 69.5 64.0 64.0 64.0 70.5 72.0 70.0 70.0 70.0 70.0 70.0 70.0 70.0	81 75 88 87 88 87 88 87 88 87 88 88	11.2 14.5 9.4 5.4 9.0 10.3 12.4 0.0 7.8 3.7 2.8 10.9 14.0 5.4 13.9 13.8 13.8 13.8 13.5 6.2 13.5 13.5 5.6 8.3 10.2 13.3	sw se se se w mw sw sw sw nw sw se se se se se se se se se se se se se	46 101 101 124 135 40 53 52 68 87 141 162 102 123 133 84 64 69 98 62 168 39 87 54 70 74	6 11 11 11 13 22 14 4 10 7 12 12 12 14 13 18 13 13 12 22 14 16 6 6 6 6 7 7	4 P 0 A 9:30 P 6 A 4 P 9 P 0 A 4:30 A	8 P 1 A 11 P 10 A 5 P 12 P 2 A 5 :30 A		

^{*}Based on least time required to blow one mile.

Stanley Kisiel, Observer

August, 1943	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.15, 29th Minimum 29.51, 17th Mean semi-daily 29.910 Range 0.64	30.32 29.61 30.00 0.71	30.50, 1934 28.87, 1930
Air Temperature, in degrees F.		
Highest 92, 2nd Lowest 45, 27th Mean 68.9 Range 47 Highest mean daily 79.5, 3rd Lowest mean daily 56.0, 27th Mean maximum 80.2 Mean minimum 57.7 Greatest daily range 36, 20th Least daily range 8, 10th	91.6 43.4 68.6 48.1	100, 1918 37, 1894, 1908 34, 1940
Precipitation, in inches		0 10 1030 - 21 1001
Precipitation 2.49 Snow Maximum precipitation in 24 hrs	4.08	8.40, 1928; .31, 1894
Number of days with .01 or more 11	11	16, 1892, 1933; 4, 1899
Wind, in miles		1371 1010 1020 1001
Total movement 2819 Greatest daily movement 169, 14th Least daily movement 38, 20th	3127	4,271, 1910; 1,920, 1894
Mean hourly velocity	4.2	40, 1941
Wind, direction		
Prevailing direction W North, days 0 North ast, days 0 Fast, days 2 Southeast, days 9 South, days 2 Southwest, days 5 West, days 4 Northwest, days 9	SW	
Weather	·	
Mean relative humidity, percent 77.4 Mean cloudiness, percent 56	70.2 49.7	67, 1901; 27, 1923
Number of clear days18 Number of fair days10	9 13	23, 1941; 0, 1915, 1929 25, 1912; 3, 1939
Number of cloudy days 3 Number hours bright sunshine 307	9 237	18, 1901, '28; 2, 1910, '23 332, 1941; 152, 1915, 1929
Percent of possible hours of bright	55.2	1711, 102, 1710, 1727
sunshine 71.5 Thunder and lightning 3rd, 12th		22.104.107.0
First frost	Sept. 21	Aug. 22, 94, 95; Oct. 13, 709

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1942.

The weather during August was very near normal in most respects. The rainfall was less than usual with a total precipitation of only 2.49 inches, while the normal for August is 4.08 inches. The total precipitation since January 1 is now 27.95 inches, the normal for that period being 29.38 inches. The mean temperature for the month was 68.9 degrees. The normal mean temperature for August is 68.6 degrees. There were 307 hours of bright sunshine which is well above the average, the normal for August being 237 hours. Wind travel was slightly less than usual, with only 2819 miles. No high winds were recorded.

Dr. J. K. Shaw, research pomologist, makes the following report on vegetation: "The weather during August was characterized by much cloudy, humid weather for the early part of the month. This resulted in some rotting and other disease injury to fruit and vegetable plants. The weather later in the month was less humid with many fair or clear days. The moisture supply continues ample. The appearance of dried out areas in lawns is one of the first signs of lack of rainfall. Except for a period in early July, lawns have presented a good appearance all summer. With the approach of shorter days and cooler weather and consequently a lessened demand for water by vegetation, there can be little damage from drouth this year. The apple crop is sizing and coloring fairly well, and the prospects are for a crop considerably below the unusually large crop of last year but not above the average of recent years."

Meteorological Series Bulletin No. 657

September 1943

Meteorological Observations

FOR

SEPTEMBER

1943

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

> Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Те	mpera	ture		ty	s		Wind		I	recipita	ition	
	Max	imum	Min	mum		Jumidi	r of Hours Sunshine	b 0		es				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number o Bright Su	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 3 4 4 5 5 6 6 7 8 8 9 9 10 11 12 13 14 15 16 17 7 18 19 20 21 22 3 23 24 25 26 26 28	83 69 65 70 79 88 91 73 81 73 74 68 71 76 64 72 75 80 67 67 64 72 75 69 69 69 69 69 69 69 69 69 69 69 69 69	5 P 2 A A P 12 M 4 P P 3 P P 1 P P 1 P P 2 P P 1 P P 2 P P 2 P P 2 P P 2 P P 2 P P 2 P P 2 P P 1 P P 2 P P 2 P P 1 P P 2 P P 2 P P 1 P P 2 P P 1 P P 2 P P 1 P P 2 P P 1 P P 2 P P 1 P P 2 P P P P	59 58 56 59 64 70 69 58 54 40 38 41 46 38 34 46 38 34 46 46 47 47 48 48 48 48 48 48 48 48 48 48 48 48 48	5 A A 12 P 5 A A 2 A A 12 P 7 A A 4 A A 12 P 7 A A 12 P 7 A A 12 P 7 A A 12 P 7 A A 12 P 7 A A 12 P 7 A A 12 P 7 A A 12 P 7 A A 13 P 7 A A 14 A A 15 P 7 A 15 P 7 A 15 P	71.0 63.5 60.5 64.5 71.5 79.0 65.5 63.0 54.0 54.5 58.5 67.0 56.5 51.0 53.0 60.5 61.5 63.0 55.0 60.5	87 80 82 84 91 91 83 85 77 70 67 79 80 86 87 74 66 70 74 81 82 71 71 62	6.4 0.3 0.6 3.5 3.4 7.2 6.6 2.4 3.6 10.5 12.7 12.6 8.6 12.4 10.3 7.3 9.6 6.4	se ne nw nw se sw nw sw nw se nw nw nw nw nw nw nw nw nw nw nw nw nw	40 85 104 79 52 117 106 75 56 70 123 108 118 128 50 83 69 111 116 93 154	6 13 7 16 8 8 13 10 22 11 11 12 7 12 13 22 17 17 17 7 8 9 10 17 13 13 13	3:30 F 6 F 6 A	8 A 8 A 8 A 8 A 8 A	.11 T T T .78	
26 27 28 29 30	62 74 82 69 57	3 P 4 P 3 P 2 P 3 P	34 39 39 52 45	5 A 6 A 7 A 7 A 1 A	48.0 56.5 60.5 60.5 51.0	59 56 71 59 79	12.0 12.0 11.9 11.7 0.3	W S S nW DW	111 120 47 141 177	7 13 7 13 17	12 M	9:30 г	.04	

^{*}Based on least time required to blow one mile.

September, 1943	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.55, 29th Minimum 29.78, 7th Mean semi-daily 30.123 Range 0.77	30.45 29.57 30.06	30.65, 1924 28.41, 1938
_	.88	1.99, 1938; .57, 1910
Air Temperature, in degrees Highest 91, 7th Lowest 34, 19th, 26th Mean 61.1 Range 57 Highest mean daily 80, 7th Lowest mean daily 48, 26th Mean maximum 73.0 Mean minimum 49.3 Greatest daily range 43, 28th Least daily range 9, 3rd	87.7 33.2 61.7 54.5	97, 1895, 1929 24.5, 1914
Precipitation, in inches Precipitation 2.40	4.26	14.55, 1938; .52, 1914
Snow Maximum precipitation in 24 hrs. 1.26, 15th Number of days with .01 or more	10	16, 1933; 3, 1903
Wind, in miles		
Total movement 2867 Greatest daily movement 177, 30th Least daily movement 40, 1st Mean hourly velocity 4.0 Maximum velocity 22, 9th, 16th	1 1	4,686, 1896; 1,414, 1894
	25.9	80, 1938
Wind, direction Prevailing direction WNW North, days 0 Northeast, days 1 East, days 1 Southeast, days 6 South, days 3 Southwest, days 2 West, days 2 Northwest, days 15		
Weather	526	
Mean relative humidity, percent 75.2 Mean cloudiness, percent 61 Number of clear days 12 Number of fair days 12 Number of cloudy days 7 Number hours bright sunshine 228 Percent of possible hours of bright sunshine 61.1 Thunder and lightning 7th, 9th	73.6 49.9 10 10 7 10 201 1 54.9	70, 1934; 27, 1908, 1914 19, 1932; 2, 1907, 1928 19, 1908; 3, 1889 20, 1934; 3, 1941 255, 1916; 106, 1934
First frost	Sept. 21	Aug. 22, '94, '95; Oct. 13, '09

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1942.

The weather during September was slightly cooler than usual, with less precipitation than normal. The total precipitation for the month was only 2.40 inches, and half of this amount fell in one storm on the 15th. The normal for September is 4.26 inches. August also had less rainfall than normal, with 2.49 inches recorded during that month. The total rainfall since the first of the year is now 30.35 inches. The normal for this period is 33.62 inches.

The mean temperature for the month was 61.1 degrees compared to a normal of 61.7 degrees for September. The highest temperature was 91 degrees on the 7th and the lowest, 34 degrees on the 19th and 26th.

The domestic heating load for the month was 163 degree-days. The normal for September is 149 degree-days.

There were 228 hours of bright sunshine compared to a normal of 201 hours for September. The total wind movement was 2867 miles, while the normal is 3271 miles. No high winds were recorded during the month.

Dr. Shaw's report on vegetation states: "The month of September is the final month of the season of vegetation. Perennial plants later go into the winter rest period. No general severe killing frost has occurred, but there was enough just after the middle of the month to partially or completely kill tender vegetation, the severity of injury depending on the location of the plants.

"The month has been dry. Coming so late in the growing season, it has not done much damage; but unless we have more abundant rains during the next two months, there will be a marked deficiency of soil moisture this winter.

"The spring was late which, combined with heavy precipitation, delayed planting. No severe drouth periods have occurred. The season has, on the whole, been quite favorable to most crops."

Meteorological Series Bulletin No. 658

October 1943

Meteorological Observations

FOR

OCTOBER

1943

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

> Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Те	mpera	iture		ty	s		Wind			Precipitat	ion	
	Max	imum	Min	imum		Tumidi	f Hour nshine	ho		es				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max, Miles per Hour*	Began	Ended	Water	Snow
1 2 3 3 4 5 5 6 6 7 7 8 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 29 30 30 30 30 30 30 30 30 30 30 30 30 30	65 71 58 57 56 70 75 80 54 61 70 75 74 69 68 54 51 52 57 54 51 54 51 54 51 55 57 57 57 57 57 57 57 57 57 57 57 57	3 P 1 P 11 A 1 P 3 P 3 P 3 P 3 P 2 P 1 P 1 A 1 D 1 A 1 P 2 P 2 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P 1	52 53 48 41 32 28 37 42 44 29 23 47 61 61 42 39 38 57 46 44 45 44 45 47 47 48 47 47 48 48 48 48 48 48 48 48 48 48 48 48 48	0 A 12 P 12 P 12 P 6 A 7 A A 12 P 6 A A 7 A A 6 A 12 P 3 A A 6 A A 7 A A 12 P 9 P 5 A 8 A A 6 A 3 A A 7 A A 13 A 14 A A 15 A A A A A A A A A A A A A A A A	58.5 62.0 53.0 49.0 44.0 49.0 57.5 58.5 62.0 41.5 42.0 48.0 53.5 65.0 55.0 44.5 52.5 51.0 49.0 44.5 40.0 52.5 51.0 52.5 51.0 52.5 51.0 52.5 51.0 52.5	92 87 67 68 73 59 69 63 54 64 62 55 68 77 95 64 68 67 77 72 87 87 87 89 90 87	0.2 3.5 2.3 2.4 6.6 11.5 11.4 11.3 11.3 11.3 11.3 2.0 3.4 0.8 5.5 4.0 0.7 6.4 1.9 0.0 0.0 0.0 1.9 10.5	nw nw nw nw n nw s nw se w nw sw sw se e w se w nw nw nw nw nw nw nw nw nw nw nw nw n	132 55 208 168 105 64 109 252 33 44 145 102 147 211 112 110 165 236 104 63 147 130 206 208 294 208 231	10 15 14 13 17 13 18 7 28 18 8 4 9 11 15 25 25 25 22 22 22	4 :30 P 4 :30 A 1 :30 A 4 A 5 :30 A 12 M	11 P 9:30 A 12 P 12 P 8 A 5 A	.64 .69 .38 .03 T	

^{*}Based on least time required to blow one mile.

October, 1943	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.53, 11th Minimum 29.15, 4th	30.53 29.42	30.68, 1929 29.00, 1926
Mean semi-daily 29.971 Range 1.38	30.06 1.11	1.47, 1926; .76, 1899
Air Temperature, in degrees F.	-o. 1	00 5 1000
Highest	79.4 23.4	90.5, 1908 17, 1936
Mean 51.4 Range 57	50.5 56.0	56.4, 1920; 43.2, 1890
Highest mean daily 67.5, 15th	30.0	
Highest mean daily 67.5, 15th Lowest mean daily 40.0, 25th Mean maximum 61.3 Mean minimum 41.5		
Mean minimum 41.5		
Greatest daily range		
Precipitation, in inches		
Precipitation 3.88	3.29	8.81, 1911; .01, 1924
Snow		
Number of days with .01 or more9	9	15, 1913; 1, 1897, 1924
Wind, in miles		
Total movement	4074	5,467, 1910; 2,540, 1894
Least daily movement _ 33, 11th		
Mean hourly velocity 6.0 Maximum velocity 28, 9th	29.5	42, 1937
Wind, direction		
Prevailing direction WNW	W	
North, days 3 Northeast, days 0	1	
East, days 1 Southeast, days 4 South, days 1		
South days 1		
Southwest, days 4		
Southwest, days 4 West, days 6 Northwest, days 12		
Weather		
Mean relative humidity, percent69.0	69.0	(7 1022 10 1021
Mean cloudiness, percent 64 Number of clear days 64	10	67, 1932; 18, 1924 21, 1938; 1, 1911
Number of fair days	10	17, 1924; 3, 1938
Number of cloudy days 1- Number hours bright sunshine 159	175.0	19, 1896; 3, 1924 232, '23, '38; 91, 1913
Percent of possible hours of bright sunshine 46.6	51.3	
Thunder and lightning 27tl First Frost Oct. 6	31.3	22 10 / 107 0 47 122
First Frost Oct. 6	Sept. 21	Aug. 22, '94, '95; Oct. 15, '39

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1942.

October was slightly warmer than usual, with more precipitation than normal. There were many pleasant days the first part of the month. The dry spell which prevailed in September continued until the 15th of October, and only half an inch of rain was recorded during the period from September 15 to October 15. However, heavy rains during the latter part of the month brought the total rainfall for October to 3.88 inches. The normal for October is 3.29 inches. The total rainfall since the first of the year is now 34.23 inches, which is 2.68 inches less than normal for this period.

The mean temperature for the month was 51.4 degrees, which is .9 degree warmer than normal. The highest temperature was 80 degrees on the 9th, and the lowest, 23 on the 11th.

The domestic heating load for October was 422 degree-days, and the total load for September and October is 585 degree-days. The normal heating load for September and October is 605 degree-days.

The first killing frost was late this year, October 6 being the date at this station. The normal date for first frost is September 21 and last year it occurred on September 29.

There was less sunshine and more wind than normal.

Meteorological Series Bulletin No. 659

November 1943

Meteorological Observations

FOR

NOVEMBER

1943

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

> Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

	Temperature		Temperature tip sau		Wind Precip				Precipitat	itation		
Maximum Minimu	n	Mean Relative Humidity	Number of Hours Bright Sunshine	ng n	nt	iles r*						
Day Degrees Time Degrees	Mean	Mean Relative	Number Bright S	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Steaw		
1 60 2 P 27 7 7 2 55 1 P 42 0 3 54 12 M 36 12 4 53 1 P 34 3 5 55 2 P 34 6 6 63 1 P 37 12 7 61 2 P 32 6 8 64 12 P 38 1 9 66 4 A 38 12 10 48 1 P 31 12 45 2 P 30 17 11 42 1 P 31 12 45 2 P 30 17 13 48 11 A 34 8 14 39 1 P 24 7 15 35 8 P 24 7 16 38 2 P 23 7 16 38 2 P 23 7 16 38 2 P 23 7 18 45 3 P 23 0 19 58 2 P 23 7 18 45 3 P 23 0 19 58 2 P 25 7 22 40 0 A 30 12 23 32 3 P 28 6 6 24 43 3 P 28 7 15 50 2 P 25 7 22 40 0 A 30 12 23 32 3 P 28 6 6 24 43 3 P 28 7 27 26 57 3 P 27 8 27 47 3 P 31 8 28 36 1 P 22 7 8 27 47 3 P 31 8 28 36 1 P 22 7 8 29 34 2 P 27 8 29 34 2 P 27 8 29 34 2 P 27 8 29 34 2 P 28 7 16 8 30 39 3 P 21 8 8 30 39 3 P 21 8 8 30 39 3 P 21 8 8 30 39 3 P 21 8 8 30 39 3 P 21 8 8 30 39 3 P 21 8 8 30 39 3 P 21 8 8 30 39 3 P 21 8 8 30 39 3 P 21 8 8 30 30 39 3 P 21 8 8 30 30 39 3 P 21 8 8 30 30 39 3 P 21 8 8 30 30 30 30 30 30 30 30 30 30 30 30 30	A 43.5 A 48.5 P 45.0 A 43.5 A 44.5 P A 50.0 A 46.5 A 37.5 A 37.5 A 39.0	67 90 60 72 73 69 69 65 65 68 45 57 80 94 563 79 88 62 100 67 58 61 67 58	10.3 0.0 5.5 3.7 1.8 0.0 7.5 0.0 6.7 0.0 9.1 5.1 9.5 2.6 0.0 9.7 6.0 9.0 9.0 9.0 9.0 9.0 9.0 9.0 9	se se se w w se se s sw nw w w w w se nw nw nw se nw nw nw nw nw nw nw nw nw nw nw se w nw nw nw se w nw nw nw nw nw nw nw nw nw nw nw nw n	63 144 172 72 72 131 53 62 244 190 236 111 237 114 84 150 73 39 47 273 407 204 119 60 89 271 63	8 11 20 13 13 14 8 22 28 12 22 18 10 10 13 4 13 22 25 14 15 11	1 P 11:30 P 0 A 8 A 0 A 1:30 P 0 A 11:30 P	6:30 P 12 P 2 A 12 P 10 A 6 P 8:30 A	.34 T .04 .16 .97 1.63	T 1.00 3.50 0.50		

^{*}Based on least time required to blow one mile.

November, 1943	Normal	Extremes
Barometer, in inches (Readings reduced to sea level) Maximum 30.48, 7th Minimum 29.59, 9th Mean semi-daily 30.070 Range 0.89	30.59 29.30 30.05 1.29	30.87, 1932 28.73, 1904 1.84, 1904; .94, 1905
Air Temperature, in degrees F. Highest	66.0	75 , 1924 , 19 38
Tiglest 16, 29th	12.9 38.9 53.0	-4, 1938 -4, 1938 44.1, 1931; 33.6, 1901
Precipitation, in inches	2.41	9.64 1027 62 1015
Precipitation 4.64 Snow 5.00 Maximum precipitation in 24 hrs.	3.41 2.34	8.64, 1927; .63, 1917 13.50, 1938
Number of days with .01 or more 10	9	22, 1921; 2, 190-
Wind, in miles Total movement	4531 6.3 30.3	5,978, 1906; 2,589, 1889 44,1938
Wind, direction		
Prevailing direction WNW North, days 1 Northeast, days 0 East, days 0 Southeast, days 7 South, days 1 Southwest, days 1 West, days 8 Northwest, days 12		
Weather Mean relative humidity, percent	70.6	
Mean cloudiness, percent	55.1 6 9 15 121	72, 1900, 1927; 34, 1917 15, 1903; 1, 1900, 1911, 1927 16, '12; 4, '89, '30, '32, '33 24, 1927; 9, 1905, 1917 182, 1903; 66, 1927
sunshine 50.5 Thunder and lightning		
First snow 15th	Nov. 6	Oct. 10, '25; Nov. 27, '3

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1942.

November had more rain and snow than usual, although the month as a whole was normal. As in October, there were many pleasant days during the month and the fall season has been an enjoyable one in most respects. The first snow of the season occurred on November 15, although a light trace was recorded on the 11th. The normal date for tirst snowfall is November 6, and last year only a trace was recorded in November. The snowstorm on the 22nd and 23rd, in which 4 inches was recorded at Amherst, was accompanied by high winds and although the storm was not considered too severe in this section, in many parts of New England it developed into one of the worst storms of the season in many years. The total snowfall for the month was 5.00 inches and the precipitation, 4.64 inches. The normal snowfall for November is 2.34 inches and the normal precipitation, 3.41 inches.

The mean temperature for the month was 38.7 degrees, while the normal for November is 38.9. Last year the mean was 38.8 degrees. The domestic heating load was 788 degree-days for November, and the normal for this month is 784 degree-days. The total load for the season through November 30 is 1373 degree-days compared to a normal of 1389 for this period.

There were 148 hours of bright sunshine for November, while the normal for this month is 121 hours.

Sleet fell on the 15th, and the roads were icy on the 21st, 22nd and 23rd.

Meteorogical Series

Bulletin No. 660

December 1943

Meteorological Observations

FOR

DECEMBER

1943

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23' 31'' N. Longitude, 72° 31' 48'' W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Те	mpera	iture	-	ty	s		Wind			Precipitat	ion	
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Stsow
1 2 3 3 4 5 6 6 7 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 30 30 30 30 30 30 30 30 30 30 30 30	42 45 40 41 44 38 39 50 42 13 34 45 36 43 37 36 43 45 36 43 43 45 36 43 45 36 43 45 46 46 47 47 47 47 47 47 47 47 47 47 47 47 47	1 P 3 P 12 M 6 P 12 M 12 P 12 M 12 P 11 A 1 P 2 P 13 P 10 A 1 P 10 A 6 P 10 A 6 P 10 A 6 P 10 A 6 P 10 A 7 P 10	22 25 31 27 23 22 24 49 33 31 6 6 8 5 4 -1 3 20 20 21 6 10 10 10 10 10 10 10 10 10 10 10 10 10	4 A A 5 A A 12 P 12 P 12 P 12 P 12 P 12 P 12 P	32.0 35.0 35.5 34.0 33.5 28.0 31.0 41.5 36.5 20.0 20.0 14.0 12.0 34.0 28.0 31.5 25.0 7.5 8.5 21.0 28.0 28.0 28.0 28.0 28.0 28.0 28.0 28	77 77 77 89 72 86 48 66 57 66 57 66 57 66 57 66 57 66 57 66 57 66 57 66 57 66 57 66 57 66 57 66 57 66 57 66 57 57 57 57 57 57 57 57 57 57 57 57 57	3.3 1.3 0.0 6.7 9.2 0.7 4.1 5.5 3.2 2.3 7.0 9.1 9.0 9.1 9.0 4.0 9.0 9.0 5.7 1.8 8.0 9.0 6.2 9.0	nw s s se w nw se w w nw nw nw se w se w	55 46 41 81 283 82 124 205 187 513 178 318 44 129 60 82 116 97 276 260 195 185 28 93 184 24 331 142 9	11 9 5 9 28 8 8 32 25 39 15 28 12 9 13 13 12 48 25 18 11 13 7 18 28 22 25 25 26 26 27 28 28 29 20 20 20 20 20 20 20 20 20 20 20 20 20	6 Р О А	11 P 8 A	.23 .03	ТТ

^{*}Based on least time required to blow one mile.

Barometer, in inches (Readings reduced to sea level) Maximum 30.55, 25th Minimum 29.36, 10th Mean semi-daily 30.051 Range 1.19 Air Temperature, in degrees F. Highest Lowest -6, 24th Mean 24.9	29.27 30.06 1.38	30.96, 1889 28.85, 1915 1.78, 1895; 1.01, 1892
(Readings reduced to sea level) Maximum 30.55, 25th Minimum 29.36, 10th Mean semi-daily 30.051 Range 1.19 *Air Temperature, in degrees F. Highest 50, 9th, 27th Lowest -6, 24th	29.27 30.06 1.38	28.85, 1915
Maximum 30.55, 25th Minimum 29.36, 10th Mean semi-daily 30.051 Range 1.19 *Air Temperature, in degrees F. Highest 50, 9th, 27th Lowest -6, 24th	29.27 30.06 1.38	28.85, 1915
Minimum 29.36, 10th Mean semi-daily 30.051 Range 1.19 * Air Temperature, in degrees F. Highest 50, 9th, 27th Lowest -6, 24th	29.27 30.06 1.38	28.85, 1915
Mean semi-daily 30.051 Range 1.19 *Air Temperature, in degrees F. Highest 50, 9th, 27th Lowest -6, 24th	30.06 1.38 54.3	
Range 1.19 *Air Temperature, in degrees F. Highest 50, 9th, 27th Lowest -6, 24th	1.38	1.78, 1895; 1.01, 1892
Highest		
Lowest		
		65.5, 1908
Mean 24.9	-1.9	-22.5, 1917
	27.5	36.9, 1891; 17.1, 1917
Range 56	56.2	
Highest mean daily 41.5, 9th Lowest mean daily 7.5, 23rd		
Lowest mean daily		
Mean maximum 35.0 Mean minimum 14.8		
Greatest daily range 32 25th		
Greatest daily range 32, 25th Least daily range 7, 11th		
Precipitation, in inches Precipitation58	3.39	7.77, 1901; .87, 1896
SnowT	8.50	26.50, '02; t, 1891
Maximum precipitation in 24 hrs.	0.00	20.30, 02, t, 1031
32, 26th, 27th	10	17, 1002, 4, 1003
Number of days with .01 or more4	10	17, 1902; 4, 1892
Wind, in miles Total movement 4411	4710	6 (01 1035 2 220 1010
Greatest daily movement 513, 11th	4/10	6,694, 1925; 3,239, 1918
Least daily movement9, 31st		
Mean hourly velocity 5.9	6.3	
Mean hourly velocity 5.9 Maximum velocity 48, 20th	31.4	48, 1938
Wind, direction		
Prevailing directionW	WNW	
North, days0		
Northeast, days0	1	
East, days0		
Southeast, days5		
South, days4		
Southwest, days0 West, days9	ĺ	
Northwest, days		
Weather		
Mean relative humidity, percent	69.4	
Mean cloudiness, percent 46	54.9	71, 1929; 39, 1919
Number of clear days	8	15, 1890; 2, 1922, 1933 16, '09; 4, '89' 30, '31, '36, '38
Number of fair days10	9	[16, 709; 4, 789, 30, 731, 736, 738
Number of cloudy days10	14	23, 33; 7, 09, 23
Number hours bright sunshine145	128	172, 1896; 63, 1933
Percent of possible hours of bright	45.2	
sunshine 51.2 Thunder and lightning —	45.2	
- I nunder and rightning		

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1942.

ANNUAL SUMMARY

Annual, 1943	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum31.05, Mar. 9	30.81	31.05, 1920
Minimum 29.15, Oct. 4	28.95	28.41, 1938
Mean semi-daily 30.017	30.01	
Range	1.85	2.47, 1938; 1.38, 1933
Air Temperature, in degrees F.		•
Highest 95, June 26 and 28 Lowest23, Feb. 16 Mean 47.3	95.7	104, 1911
Lowest ——23. Feb. 16	-12.2	-26, 1904
Mean 47.3	47.4	49.8, 1921; 44, 1904
Range 118 Highest mean daily 83.5, June 27 Lowest mean daily -13.5, Feb. 16 Mean maximum 57.3	107.8	
Highest mean daily 83.5, June 27		
Lowest mean daily		
Mean maximum 57.3	57.8	
Mean minimum 37.1	36.5	
Greatest daily range 44, Nov. 12 Least daily range 3, Jan. 18		
Least daily range3, Jan. 18		
Precipitation, in inches	ļ	
Precipitation 39.45	43.70	59.00, 1938; 30.68, 1908
Snow	47.78	89.00, 1893; 24.50, 1919
Maximum precipitation in 24 hrs.		05.00, 1055, 24.50, 1515
2.60, Nov. 8 and 9		
Number of days with .01 or more124	124	144, 1902; 96, 1924
Wind, in miles		
Total movement 50,963 Greatest daily movement 513, Dec. 11	52,223	63,571, 1908; 36,257, 1894
Greatest daily movement 513, Dec. 11		
Least daily movement 9, Dec. 31		
Least daily movement 9, Dec. 31 Mean hourly velocity 5.8 Maximum velocity 48, Dec. 20	5.8	
Maximum velocity 48, Dec. 20	39.5	80, 1938
Wind, direction		
Prevailing direction W	W	
North, days9		
Northeast, days 2		
East, days6		
Prevailing direction W North, days 9 Northeast, days 2 East, days 6 Southeast, days 74 South, days 33		
South, days 33 Southwest, days 24 West, days 64 Northwest, days 153		
Southwest, days 24		
West, days		
Northwest, days		
Weather		
Mean relative humidity, percent 71.0 Mean cloudiness, percent 58.0	67.6	
Mean cloudiness, percent	51.7	60, '98, '01, '02; 41, '08, '24
Number of clear days	116	217, 1941; 59, 1927
Number of fair days 102	123	217, 1941; 59, 1927 182, 1912; 64, 1936
Number of cloudy days 85	126	179, '01, '02; 71, 1910, 19 4 1
Number of clear days	2,353	3038, 1941; 1864, 1902
sunshine -62.8 Last snow Apr. 14 First snow Nov. 15 Last frost Apr. 14	52.8	
Last snow Apr. 14	April 15	Mar. 14, '10; May 11, '07 Oct. 10, '25; Dec. 13, '41
First snow Nov. 15	Nov. 6	Oct. 10, 25; Dec. 13, '41
First snow Nov. 15 Last frost Apr. 14 First frost Oct. 6	May 14	Apr. 23, '04; June 8, '32
First frost Oct. 6	Sept. 21	Aug. 22, '94, '95; Oct. 13, '09

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 661

January 1944

Meteorological Observations

FOR

JANUARY

1944

C. I. GUNNESS

OBSERVATORY

Latitude, 42 $^{\circ}$ 23' 31'' N. Longitude, 72 $^{\circ}$ 31' 48'' W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Те	mpera	ture		y:			Wind			Precipita	ıtion	
	Max	imum	Min	imum		Humidity	of Hours inshine	bo.	t	* es				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative I	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Srsow
1 2 3 4 5 6 7 8 8 9 10 11 12 13 14 15 16 17 18 19 20 21	37 32 36 37 38 38 30 19 32 27 38 37 30 33 33 35 33 42 41 46	12 M 2 P 12 P 4 P 2 P 10 A 12 M 1 P 3 P 4 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2	15 7 15 33 33 31 30 20 8 3 16 5 17 18 22 22 7 7 -2 9 13 28 29 20 20 20 20 20 20 20 20 20 20 20 20 20	12 P 8 A 8 A 12 P 12 P 12 P 12 P 4 A 12 P 7 A 0 A 4 A 7 A 12 P 8 A 2 A 8 A 8 A	26.0 19.5 25.5 35.0 34.5 34.0 25.0 13.5 11.0 24.0 19.0 22.5 30.0 29.5 18.5 15.5 22.0 23.0 35.0 31.0	54 57 73 62 77 92 55 70 64 62 80 61 57 76 47 47 86 92 80 80 80 80 80 80 80 80 80 80	7.1 6.7 0.0 0.0 0.0 9.3 9.3 9.3 8.3 6.7 7.2 8.7 9.5 9.5 2.8 9.6 7.8	nw nw nw nw nw nw nw nw nw se nw se nw se nw sw nw	135 47 32 274 160 194 353 232 152 152 14 194 156 55 224 18 27 1154	22 6 6 10 20 13 28 25 25 22 22 3 3 28 28 14 13 22 8 1 10 6 6 17 25 11	4 1	8 p	.80	2.25
21 22 23 24 25 26 27 28 29 30	36 41 35 37 49	10 P 3 P 11 P 2 P	29 14 8 32 34	7 A 12 P 7 A 6 A	32.5 27.5 21.5 34.5 41.5	85 59 72 87 84	0.0 6.9 5.3 0.0 0.0	nw w nw se	53 145 5 16 72	11 25 1 5 19	9:30 A 8 P 5 P	2:30 p 9:30 p 8 p	.29 .02 .03	2.50
28 29 30	42 42 43 33	1 P 0 A 11 A 0 A	30 31 11	8 A 7 A 12 P	36.0 36.5 22.0	68 68 59	9.9 4.2 9.9	se in w w	61 156 389	6 20 32	0 а	4 A	.10	1.00
31	31	8 P	9	8 1	20.0	65	3.8	se	66	11				Т

^{*}Based on least time required to blow one mile.

January, 1944	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum 30.59, 17th	30.70	31.00, 1927
Minimum 29.52, 31st	29.20	28.55, 1913
Mean semi-daily 30.098 Range 1.07	30.07	3.19 1012 0.07 1907
Range 1.07	1.50	2.18, 1913; 0.97, 1896
Air Temperature, in degrees F.		
Highest 49, 27th Lowest -2, 17th	51.1	66.0, 1932
	-7.1 24.2	-26.0, 1904 34.2, 1913; 13.9, 1918
Mean 26.7 Range 51	58.2	01.2, 1910, 10.9, 1910
Highest mean daily		
Lowest mean daily 11.0, 9th		
Mean maximum		
Greatest daily range 35, 17th		
Mean 26.7 Range 51 Highest mean daily 41.5, 27th Lowest mean daily 11.0, 9th Mean maximum 35.1 Mean minimum 18.4 Greatest daily range 35, 17th Least daily range 4, 4th		
Precipitation, in inches		
Precipitation 1.24	3.61	7.15, 1898; 1.07, 1896
Snow 5.75	13.32	33, 1898, 1923; 1.50, '08
Maximum precipitation in 24 hrs 0.80, 6th Number of days with .01 or more 5	11	15, '93, '20; 4, 1901
Number of days with lot of more		, , , , , , , , , , , , , , , , , , , ,
Wind, in miles	5055	7770, 1908; 2896, 1895
Total movement	3033	7770, 1908, 2890, 1893
Least daily movement 3, 18th		
Least daily movement3, 18thMean hourly velocity4.1Maximum velocity32, 30th	6.8	
Maximum velocity	32.7	47, 1938
Wind, direction		
Prevailing direction WNW North, days 1 Northeast, days 1 East, days 0 Southeast, days 5 South, days 2	WNW	
North, days		
Northeast, days		
Southeast, days		
Southeast, days		
Southwest, days 2 West, days 6		
South, days 2 South, days 2 Southwest, days 2 West, days 6 Northwest, days 14		•
Weather		
	70.	
Mean cloudiness, percent 54	70.1	78, 1932; 37, 1908
Number of clear days 20	9	18, 1941; 2, 191-
Number of clear days	9	18, 1890, 1926; 3, 1916
Number of cloudy days 8 Number hours bright sunshine . 187	13 137	22, 1923, 1931; 4, 1920, 1939 214, 1920; 74, 1932
Percent of possible hours of bright	137	314, 1920; 74, 1932
sunshine 63.6	46.7	
Thunder and lightning —		

Note The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1943.

The weather during January was somewhat warmer than usual with very little snow. The mean temperature for the month was 26.7 degrees, while the normal for January is 24.2 degrees. The temperature dropped below zero on one day, the 17th, when the thermometer reached 2 below zero. The heating load during January was 1186 degree-days whereas the normal for this month is 1265 degree-days. The total heating load for the season to the end of January is 3803 degree-days, the normal for this period being 3817 degree-days.

The snowfall during the month was 5.75 inches, while usually we get 13.32 inches during January. The total precipitation for the month was only 1.24 inches, the normal being 3.61 inches. The total snowfall for the winter to the end of January is 10.75 inches, the normal for the period being 24.16 inches. There were 187 hours of bright sunshine compared to a normal of 137 hours. Total wind movement was only 3914 miles, while the normal wind movement for January is 5055 miles. The highest wind velocity was 30 miles per hour on the 30th. The mean relative humidity was 70 percent.

Meteorological Series

Bulletin No. 662

February 1944

Meteorological Observations

FOR

FEBRUARY

1944

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23' 31'' N. Longitude, 72° 31' 48'' W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Те	mpera	ture		ty	s		Wind			Precipita	tion	
Day	Degrees Degrees	Time	Min	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Strow
1 2 3 4 4 5 5 6 7 7 8 9 10 11 12 13 14 4 15 5 16 16 17 18 19 20 21 22 23 24 25 26 27 28 29	32 38 34 40 43 32 27 26 28 19 23 45 27 43 45 29 43 38 44 51 40 36 37 45 39 45 39 45 39 45 39 45 39 45 39 45 45 45 45 45 46 46 46 47 47 47 47 47 47 47 47 47 47 47 47 47	4 A A A A A A A A A A A A A A A A A A A	10 7 24 15 10 18 14 7 5 16 10 11 3 33 33 32 29 24 20 32	12 P 8 A 7 A 8 A 8 A 12 P 7 A 12 P 7 A 12 P 7 A 12 P 7 A 12 P 7 A 12 P 7 A 6 A 6 A 7 A 7 A 8 A 7 A	21.0 19.5 31.0 24.5 25.0 30.5 23.0 17.0 15.5 22.0 14.5 17.0 19.0 25.5 22.5 32.0 34.0 35.5 22.5 32.0 34.0 35.5 32.0 35.5 32.0 35.5 32.0 35.5 32.0 35.5 36.5 36.5 36.5 36.5 36.5 36.5 36.5	62 59 600 444 7550 752 642 89 647 745 65 847 653 757 75	7.2 10.0 7.4 10.1 5.2 7.8 2.8 10.2 5.5 9.0 0.8 3.1 10.5 4.1 5.9 4.1 5.9 4.1 5.9 4.1 5.9 0.0 0.0 0.0 0.3	w w w w se nw nw se sw w se nw nw w se nw nw w se nw w se nw w se nw w se nw w se nw w se nw w se nw nw nw nw nw nw nw nw nw nw nw nw nw	331 146 193 137 142 325 73 197 81 218 192 86 401 179 68 259 104 135 126 200 58 25 29 8	32 20 20 20 15 12 22 32 15 18 7 22 18 20 18 24 25 10 25 17 17 13 12 20 25 18 17 17 17 17 18 18 19 20 20 20 20 20 20 20 20 20 20 20 20 20	6 A 0 A 4:30 P 0 A 7:30 P 0 A 9:30 P 0 A	12 P 2 P 12 P 5 A 12 P 3 A	.25 .32 .32 .33 .17 .15	T 3.00 3.50 0.25 T T 0.25

^{*}Based on least time required to blow one mile.

February, 1944	Normal	Extremes
Barometer, in inches (Readings reduced to sea level) Maximum 30.52, 19th Minimum 29.24, 15th Mean semi-daily 30.052 Range 1.28	30.63 29.24 30.04 1.38	31.05, 1920 28.56, 1895 1.89, '00, '08; .88, '13 '31
Air Temperature, in degrees F. Highest 51, 24th Lowest -5, 14th Mean 25,9 Range 56 Highest mean daily 40, 24th	50.5 -7.4 23.7 58	65.0, 1930 -23, 1943
Air Temperature, in degrees F. Highest 51, 24th Lowest -5, 14th Mean 25,9 Range 56 Highest mean daily 40, 24th Lowest mean daily 14.5, 11th, 19th Mean maximum 35.5 Mean minimum 10.3 Greatest daily range 48, 14th Least daily range 2, 27th	32.8 14.5	
Precipitation, in inches Precipitation 2.34 Snow 7.50 Maximum precipitation in 24 hrs		8.12, 1900; 0.62, 1901 48.75, 1893; 0.50, 1937
Number of days with .01 or more 10	10	15, '93, '20; 4, 1901
Wind, in miles Total movement 4777 Greatest daily movement 25, 27th Mean velocity 6.9 Maximum velocity 48, 15th	4776	6445, 1896; 3438, 1892
Mean velocity 6.9 Maximum velocity 48, 15th	7.1 31.7	48, 1934, 1937
Wind, direction Prevailing direction WNW North, days 0 Northeast, days 0 East, days 0 Southeast, days 7 South, days 0 Southwest, days 1 West, days 8 Northwest, days 13	WAIN	
Weather Mean relative humidity, percent Mean cloudiness, percent Soft Number of clear days Number of fair days Number of cloudy days Number hours bright sunshine Percent of possible hours of bright sunshine Thunder and lightning Soft Mean relative humidity, percent Soft Soft Soft Soft Soft Soft Soft Sof	66.6 50.9 10 8 10 158	66, '90, '27; 31, 1905 19, 1941; 2, 1927 16, 1920; 2, 1936 16, 1894; 2, 1920 221, 1924; 110, '27, '38

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that lumidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1943.

The weather during February was unusually pleasant. The temperature was not as low as usual, there was less rain and snow than normal, and a greater amount of sunshine than usual was recorded. Roads were unusually free from ice.

The mean temperature for the month was 25.9 degrees, the normal being 23.7 degrees. The lowest temperature was -5 degrees, the normal minimum for February being -7.4 degrees. It is interesting to recall that the temperature fell to -23 degrees on February 16 in 1943. The heating load for the month was 1134 degree-days, the normal being 1165 degree-days. The total heating load for the season to the end of February is 4937 degree-days, while the normal for that period is 4982 degree-days.

Only 7.50 inches of snow fell during the month while the normal snowfall for February is 13.99 inches. The total precipitation was 2.34 inches compared to a normal of 3.19 inches. The total snowfall for the winter is now 18.25 inches, while usually we get 38.15 inches to the end of February. There were 170 hours of bright sunshine during February compared to a normal of 158 hours. The mean relative humidity was 66.2 percent, the normal is 66.6 percent. The maximum wind velocity was 48 miles per hour on the 15th.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 663

March 1944

Meteorological Observations

FOR

MARCH

1944

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23' 31'' N. Longitude, 72° 31' 48'' W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Te	mpera	ture		ty	s		Wind			Precipita	tion	
	Max	imum	Min	imum		Humidi	f Hour nshine	to.		es				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8 9 10 11 12 13	34 27 35 27 35 47 40 43 31 30 38 57 51	2 P 1 P 2 P 4 P 4 P 3 P 3 P 2 P 12 M 4 P 1 P 4 P	16 10 15 10 5 8 30 27 18 14 11 27 33	12 P 6 A 3 A 12 P 7 A 2 A 12 P 12 P 7 A 6 A 12 P	25.0 18.5 25.0 18.5 20.0 27.5 35.0 24.5 22.0 24.5 42.0 42.0	52 37 69 40 36 36 93 55 48 55 62 51 86	11.2 11.2 6.0 8.0 11.4 9.5 0.0 10.3 11.6 10.0 11.7 9.5 0.0	nw nw se nw nw s se sw w w se se sw	484 338 53 173 241 173 106 206 292 181 115 154 108	32 25 7 17 28 13 13 20 25 17 14 17 32	0 A	1 Р 12 Р 10:30 A	.83	T 1.50
14 15 16 17 18	46 38 37 42 44	3 P 12 M 1 P 2 P 1 P	26 31 33 35 19	7 A 7 A 0 A 0 A 12 P	36.0 34.5 35.0 38.5 31.5	43 66 99 89 50	11.8 0.3 0.0 0.0 5.6	nw se se se w	346 93 43 79 279	32 8 5 15 28	12 M 0 A 0 A	12 P 12 P 2:30 P	.25 .46 .35	1.00
19 20	25 28 46 44	9 P 12 M 1 P 2 P	9 18 20 16	6 A 5 A 5 A 6 A	17.0 23.0 33.0 30.0	43 88 65 47	9.0 5.0 9.8 12.2	nw nw nw nw	238 195 121 89	28 18 12 8	9:30 P 0 A	12 P 8 P	.04	.50 5.00
21 22 23 24 25 26 27 28 29 30 31	40 50 63 51	9 P 4 P 3 P 3 P	29 34 28 35	6 A 2 A 6 A 12 P	34.5 42.0 45.5 43.0	90 50 61 47	0.0 9.3 12.4 12.4	se w se w	64 255 126 270	5 32 17 25	9 A	10 р	.78	Т
27 28	49 46	12 M 2 P	34 32 27	0 A 12 P	41.5 39.0	64 45	1.4	se nw	245 110	15 40 20	12:30p 8 p	5:30 p	.36	.25
29 30 31	48 37 45	12 м 4 р 1 р	32 29	5 A 1 A 12 P	37.5 34.5 37.0	51 100 69	4.7 0.0 10.0	e nw w	119 146 146	10 20	0 A 7 P	10 P 8 P	.39	.23 T

^{*}Based on least time required to blow one mile.

STANLEY KISIEL, Observer

March, 1944	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum 30.73, 22nd	30.57	31.05, 1943
Minimum 29.43, 23rd	29.24	28.47, 1914
Mean semi-daily 30.099 Range 1.30	30.00 1.34	2.10, 1914; 0.85, 1915
Air Temperature, in degrees F.	(20	70.5 1007 1031
Highest	63.8 6.1	79.5, 1907, 1921 -7.5, 1906
Mean 32.0	34.4	7.5, 1900
Range 58	57.6	•
Highest mean daily 45.5, 25th		
Lowest mean daily	43.2	
Mean minimum 22.9	25.4	
Greatest daily range		
Least daily range4, 16th		
Precipitation, in inches		
Precipitation 4.36	3.70	7.89, 1942; 0.12, 1915
Snow 8,25	7.47	27, 1899; 0, 1921
Maximum precipitation in 24 hrs83, 7th	11	17, 1890, 1913, 1936; 3, 1915
Number of days with .01 or more13	11	17, 1090, 1913, 1930, 3, 1913
Wind, in miles		
Total movement 5538	5944	8182, 1896; 3006, 1905
Greatest daily movement 484, 1st Least daily movement 43, 16th		
Mean velocity 7.4	7.7	
Maximum velocity40, 28th	31.2	48, 1932, 1939, 1941, 1942
Wind, direction	-	
Prevailing direction WSW	WNW	
North days 0		
Northeast, days0		
East, days1 Southeast, days10		
South, days		1
Southwest, days1	1	
West, days6		
Northwest, days12		
Weather		
Mean relative humidity, percent	64.2	
Mean cloudiness percent 57	51.8	68, 1901; 27, 1915 22, 1924; 3, 1901
Number of clear days 19	11	22, 1924; 3, 1901
Number of fair days 4 Number of cloudy days	10	17, 1900; 1, 1943 21, 1901; 1, 1915
Number of clear days	199	292, 1924; 93, 1901
Percent of possible hours of bright		
sunshine 60.9	53.6	
Thunder and lightning		

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1943.

The weather during March was nearly normal in most respects. It was slightly colder than usual and there was more rain and more sunshine than normal. The mean temperature for the month was 32 degrees, the normal being 34.4 degrees. The lowest temperature was 5 degrees on the 5th. The normal minimum for March is 6.1 degrees. The domestic heating load for the month was 1022 degree-days compared to a normal of 950 degree-days. The heating load for the season to the end of March is 5959 degree-days, the normal for this period being 5932 degree-days.

The total precipitation for the month was 4.36 inches, with 8.25 inches of snow. The normal precipitation for March is 3.70 inches and the normal snowfall 7.47 inches. The total precipitation since January 1 is 7.94 inches, while the normal for the three months is 10.50 inches. We have had a total of 26.50 inches of snow during the winter, the normal snowfall to the end of March being 45.62 inches.

There were 226 hours of bright sunshine, the normal being 199 hours. The highest wind velocity was 40 miles per hour on the 28th.

Dr. J. K. Shaw, research pomologist, makes the following statement: "There has been no severe cold during the past winter but snowfall was deficient and the ground bare most of the time. We should look for no material winter injury to tree fruits and few peach fruit buds have been killed. Therefore, a good peach crop is to be expected as spring frosts rarely kill the flowers or young fruits in Massachusetts. The absence of a snow cover is not good for strawberry plants but well-mulched beds should not suffer. The present state of vegetation is somewhat retarded, and we do not expect an early spring. Resumption of growth depends on future rather than past temperatures and a few days of warm weather will bring a normal stage of progress."

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 664

April 1944

Meteorological Observations

FOR

APRIL

1944

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23' 31'' N. Longitude, 72° 31' 48'' W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Te	mpera	ture		ty	s		Wind			Precipita	tion	
		imum 		imum 		Mean Relative Humidity	Number of Hours Bright Sunshine	ing	ent	files ar*				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relativ	Number Bright	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 4 5 5 6 7 7 8 8 9 10 11 12 13 14 4 15 5 16 6 17 7 18 19 20 22 23 24 22 5 26 27 28 29 30 31	51 52 43 40 41 48 59 42 64 53 40 40 41 53 40 40 41 53 40 40 41 48 59 40 41 48 59 60 40 59 60 60 60 60 60 60 60 60 60 60 60 60 60	3 P 12 M 5 P 6 P 6 P 12 M 14 P 12 P 14 P 14 P 14 P 14 P 14 P 15 P 16 P 17 P 18 P 18 P 18 P 18 P 18 P 18 P 18 P 18	26 22 28 23 26 25 34 33 32 36 36 37 37 32 35 37 37 32 35 36 37 37 37 37 38 38 39 39 39 39 39 39 39 39 39 39 39 39 39	12 P 6 A 6 A 6 A 12 P 4 A 5 A 6 A 12 P 5 A 8 A 12 P 3 A 5 A 12 P 3 A 5 A 12 P 3 A 5 A 5 A 7 A 8 A 8 A 8 A 8 A 8 A 8 A 8 A 8 A 12 A 8 A 8 A 8 A 8 A 8 A 8 A 8 A 8 A 8 A 8	38.5 42.0 35.5 31.5 33.5 46.5 37.5 48.0 41.5 37.5 42.0 38.5 47.5 48.5 48.5 48.5 48.5 48.5 48.5 48.5 48.5 48.5 50.5 48.5 50.5	52 50 42 56 63 49 74 87 65 100 42 37 59 96 41 50 99 53 81 88 83 67 45 38 34	10.4 6.1 12.3 8.3 2.9 12.5 13.0 0.0 9.3 1.8 7.6 0.0 8.0 13.3 0.8 0.0 4.7 13.5 13.6 0.0 13.6 9.0 0.0 13.3 13.5 13.5 13.6 13.6 13.6 13.6 13.6 13.6 13.6 13.6	sw w nw w w nw se nw w w w nw nw nw nw nw nw nw nw nw nw	181 194 231 67 80 159 256 32 56 130 97 94 364 357 61 89 125 61 39 115 95 124 1100 125 324 173 108	18 20 17 8 9 20 28 6 10 28 13 8 39 32 13 8 20 20 20 11 13 8 13 11 17 17 20 18 18 18 19 19 19 19 19 19 19 19 19 19 19 19 19	8 A 0 A 4 P 4 A 2 P 0 A 5:30 A 8 P 0 A 9 A 0 A	1 P 3:30 A 9 P 2 P 12 P 9 P 12 P 10:30 P 10 A 2 A	.29 .06 .23 .34 .74 .38 .22 .11 1.23 .01	

^{*}Based on least time required to blow one mile.

STANLEY KISIEL, Observer

WIGHTIEF 90		
April, 1944	Normal	Extremes
Barometer, in inches		
· ·		
(Readings reduced to sea level)	30.48	20.71 1011
Maximum 30.41, 22nd, 23rd	29.38	30.71, 1911
Maximum 30.41, 22nd, 23rd Minimum 29.37, 12th Mean semi-daily 29.996	29.99	28.98, 1943
Range104	1.11	1.52, 1930; .72, 1919
Kange104		1.02, 1700 ; .72, 1717
Air Temperature, in degrees F.	70.4	00 1011
Highest 74, 30th	79.4 22.0	90, 1941 8.5, 1923
Lowest23, 4th	45.7	52, 1921; 41, 1943
Mean 42.9	57.4	32, 1721, 41, 1943
Range 51 Highest mean daily 56, 30th Lowest mean daily 31.5, 4th Mean maximum 53.1	37.4	
Highest mean daily 56, 30th		
Lowest mean daily 31.5, 4th	56.8	
Mean maximum 53.1	34.8	
Mean minimum 32.6	34.0	
Greatest daily range 41, 20th Least daily range 5, 12th		
Least daily range		
Precipitation, in inches		
Precipitation 3.66	3.35	6.89, 1929; .55, 1941
Snow	2.17	11, 1891; 0, 1910, 1934
Maximum precipitation in 24 hrs. 1.23, 24th		, , , , , , , , , , , , , , , , , , , ,
Number of days with .01 or more	11	18, 1909; 3, 1892
Wind, in miles		
	5404	8208, 1908; 3853, 1917
Total movement 4306 Greatest daily movement 364, 13th		, , , , , , , , , , , , , , , , , , , ,
Least daily movement		
	7.5	
Mean hourly velocity 6.0 Maximum velocity 39, 13th	31.4	40, 1935, 1938, 1940
Wind, direction		
Prevailing directionWNW	WNW	
North, days1	******	
Northeast, days0		
East, days1		
Southeast, days 1		•
South, days	1	1
South, days 1 Southwest, days 1		
West, days 12		
Northwest, days 13		
Weather		
Mean relative hymidity percent 61.7	61.6	75, 1901; 34, 1925, 1927
Mean cloudiness percent 58	51.8	23, 1941; 3, 1898, 1902
Number of clear days 13	11	18, 1915; 2, 1894, 1901
Number of fair days 6	9	22, 1901; 1, 1941
Mean cloudiness, percent 58 Number of clear days 13 Number of fair days 6 Number of cloudy days 11 Number hours bright sunshine 220 Percent of possible hours of bright	10	329, 1941; 103, 1901
Number hours bright synshine 220	220	022, 1211, 103, 1901
Percent of possible hours of bright	550	1
	54.7	
sunshine 54.7 Thunder and lightning 16th, 24th Last snow March 29	""	
Last snow March 29	Apr. 15	Mar. 14, 1910; May 11, 1907
		1

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1943.

The weather during April was cold with a mean temperature of 42.9 degrees. The normal mean temperature for April is 45.7 degrees. The lowest temperature was 23 degrees on the 4th and the highest, 74 degrees on the 30th. The domestic heating load for April was 664 degree-days compared to the normal of 579 degree-days. The total heating load for the season is now 6623 degree-days while the normal heating load to the end of April is 6511 degree-days.

The total precipitation during April was 3.66 inches. There was no snowfall. Normal precipitation for April is 3.35 inches; normal snowfall, 2.17 inches. The total snowfall for the winter was 23.50 inches compared to the normal of 47.79 inches. Last winter's snowfall was 40.50 inches. There were 220 hours of sunshine. The normal is also 220 hours. The total wind movement was 4306 miles, while the normal is 5404 miles.

Dr. J. K. Shaw makes the following comments: "Vegetation made slow progress during April because of much cold weather. The month closes with early leafing trees showing only tiny leaves just breaking from the bud. Apple buds have not shown any color. The season is a week or more later than average. The rainfall will help to restore the depleted water reserves of the soil, and the later the buds break the less danger from spring frosts. Severe winter injury has become apparent in blueberry plants, and raspberry canes have been injured. No winter injury to tree fruits is yet apparent, and we can feel quite confident that none will appear. Apple and peach fruit buds are evident and there is promise of a heavy bloom. Some evergreen trees show winter injury. The ground froze deeply and was hardly completely thawed out at the end of the month. This condition together with winds may have been factors in causing winter injury. Except for late planting, the crop season of 1944 starts off fairly well.'

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 665

May 1944

Meteorological Observations

FOR

MAY

1944

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23' 31'' N. Longitude, 72° 31' 48'' W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST. MASS.

		Te	mpera	ture		ty	s		Wind			Precipita	tion	
	Max	imum	Min	imum		Humidity	f Hours nshine			8				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative F	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 4 5 5 6 6 7 7 8 9 10 11 12 13 14 4 15 16 17 18 19 20 21 22 23 24 25 6 27 28 29 30 31	76 62 91 90 91 83 79 65 72 78 80 83 79 75 82 65 67 73 82 65 67 77 88 88 88 67 77 88 88 89 67 77 88 89 89 89 89 89 89 89 89 89 89 89 89	2 P 4 P 3 P 3 P 4 P 4 P 4 P 4 P 4 P 4 P 1 P 4 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P 1	50 443 49 527 56 44 39 49 154 57 49 46 37 33 58 58 54 53 55 56 56 56 56 57 56 57	12 PAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA	63.0 63.0 62.0 69.5 71.5 70.0 67.5 54.5 53.5 68.5 69.0 64.0 64.0 64.0 64.0 61.0 62.0 64.0 67.5 70.0 67.5 70.0 67.5 70.0 67.5 70.0 67.5 70.0 67.5 70.0 67.5 70.0 67.5 68.5 69.0	46 71 68 59 53 67 59 56 56 57 64 56 76 41 38 72 83 72 83 72 83 75 85 85 85 85 85 85 86 86 87 87 87 87 87 87 87 87 87 87 87 87 87	14.0 5.1 9.3 13.3 14.2 14.2 8.5 14.3 5.6 14.4 14.5 13.0 14.6 14.7 14.7 14.7 13.7 0.8 3.7 2.0 5.0 4.7 11.7 15.0 15.0 15.0 15.0 14.4	n s se se se se se se se sw sw sw s w s w	100 112 157 126 131 169 280 211 84 112 65 72 121 216 94 60 62 233 136 126 88 120 73 36 72 113 112 112 112 112 130	13 13 13 14 14 14 18 28 18 11 9 6 10 20 20 17 18 22 17 18 10 10 10 10 10 10 10 10 10 10 10 10 10	7 P 4:30 P 8:30 A 8:30 P 0 A	10 P 8 P 5 P 6:30 P 10 P 8 P	.14 .46 T	

^{*}Based on least time required to blow one mile.

STANLEY KISIEL, Observer

May 1944	Normal	Extremes
Barometer, in inches (Readings reduced to sea level) Maximum 30.46, 11th Minimum 29.65, 22nd Mean semi-daily 30.089 Range 0.81	30.39 29.51 29.96 0.88	30.62, 1936 29.10, 1938
Air Temperature, in degrees F. Highest 93, 31st Lowest 30, 19th Mean 63.4 Range 63 Highest mean daily 75, 31st Lowest mean daily 48.5, 19th Mean maximum 77.1 Mean minimum 49.8 Greatest daily range 44, 21st Least daily range 7, 22nd, 23rd	86.1 31.2 57.1 54.9	94.5, 1896, 1911 24.0, 1900
Precipitation, in inches Precipitation	3.60	7.44, 1931; .48, 1903
Snow	12	20, 1901; 5, 190;
Wind, in miles Total movement 3697 Greatest daily movement 280, 7th Least daily movement 36, 24th Mean hourly velocity 5.0 Maximum velocity 28, 7th	6.1	5946, 1907; 2180, 189 45, 19.
Wind, direction SSW Prevailing direction SSW North, days 2 Northeast, days 1 East, days 8 Southeast, days 8 South, days 7 Southwest, days 5 West, days 3 Northwest, days 4	W	
Weather Mean relative humidity, percent 62.2 Mean cloudiness, percent 53.1 Number of clear days 21 Number of fair days 7 Number of cloudy days 3 Number hours bright sunshine 350 Percent of possible hours of bright sunshine 77.5 Thunder and lightning 13th, 15th	52.1 11 11 3 9 252 55.6	70, 1901, 1902; 30, 192 20, 1923, 1941; 0, 192 17, 1907, 1926; 5, 192 20, 1927; 2, 194 338, 1941; 137, 192

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1943.

May of this year was the warmest, sunniest and very nearly the dryest May on record here at the station; and yet on the morning of the 19th a killing frost occurred which caused serious damage to crops in this section.

The mean temperature for the month was 63.4 degrees, compared to a normal of 57.1 degrees. This is the highest mean temperature for May recorded at this station in 56 years. The nearest approach to this record is a mean temperature of 62.5 degrees recorded in 1911. The highest temperature was 93 degrees on the 31st and the lowest, 30 degrees on the 19th. The first and last weeks of the month were exceptionally warm. The domestic heating load for the month was 112 degree-days, the normal for May being 207 degree-days. The total heating load for the season to the end of May is 6735 degree-days while the normal for this period is 6778 degree-days.

The total precipitation for the month was 1.35 inches, which is 2.25 inches less than normal for May. Rain fell on only four days. The total precipitation since the first of the year is now 12.95 inches, while the normal for this period is 17.45 inches. The outstanding feature of May weather last year was the large number of rainy days, 15 in all, with a total rainfall of 5.62 inches.

There were 350 hours of bright sunshine compared with a normal of 199 hours. There were no high winds during the month.

Dr. J. K. Shaw, research pomologist, makes the following interesting comments regarding the progress of vegetation: "At the end of April the development of vegetation was a week or more behind normal. During the first half of May there were many days when the temperature was unusually high and the date of bloom of fruit plants was two or three days earlier than normal. Warm weather continued through the blooming period, bees were active and prospects were for a heavy set of fruit. Both apples and peaches blossomed heavily. On May 17 the maximum temperature was 85 degrees. On the afternoon of the 18th the temperature rose to 58 degrees and then fell to a minimum of 30 degrees on the morning of the 19th. This caused severe damage to tree fruits and injured tender vegetation. Orchards on high sites with good air drainage apparently escaped injury but those on low sites or in "frost pockets" suffered complete loss of the setting fruits. Some soft, tender new shoot tips were killed and there was injury to all tender vegetation that had been encouraged by the warm days preceding May 18. This is the second time in 37 years that there has been extensive spring frost injury in our orchards. Rainfall during the month was deficient. While the month closes with no serious drouth injury, shallow-rooted plants will soon suffer if rains do not come early in June."

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 666

June 1944

Meteorological Observations

FOR

JUNE

1944

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23' 31'' N. Longitude, 72° 31' 48'' W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Те	mpera	ture		ity	s		Wind			Precipitat	ion	
Day	Max	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 3 4 4 5 6 6 7 7 8 9 10 11 12 13 13 14 15 16 6 17 18 19 20 21 22 23 24 25 26 29 30 31	80 89 76 70 74 83 73 74 66 78 80 77 71 68 80 86 81 62 64 64 64 81 85 70 68 80 90 86 82 82	3 P 4 P 8 P 5 3 P 2 P 4 P 3 P 3 P 0 A 4 P 3 P 3 P 11 A 4 P 8 A 4 P 8 A 4 P 8 A 12 M 6 P 3 P 3 P 8 A 4 P 8 A 8 A 8 A 8 A 8 A 8 A 8 A 8 A 8 A 8 A	62 59 52 47 45 55 48 40 54 54 54 55 62 60 60 55 58 66 67 62	4 A A A A A A A A A A A A A A A A A A A	71.0 74.0 64.0 58.5 59.5 69.0 60.5 57.0 66.0 66.0 66.0 73.0 69.5 59.0 69.5 58.5 68.0 61.0 65.0 64.0 69.0 64.0 67.0 67.0 67.0 67.0 67.0 67.0 67.0 67	81 60 59 47 56 67 63 53 53 94 69 60 93 74 63 91 97 97 94 87 70 74 72 71 59	8.7 15.1 11.0 15.1 11.6 10.9 15.2 13.1 0.0 10.6 13.4 14.6 8.0 0.0 0.0 0.0 0.0 0.4 11.1 18.9 0.0 0.3 7.7 15.3 12.5 15.1	nw nw nw se se se w w se nw se se w nw se nw nw se ne nw nw se	77 45 231 93 199 240 200 93 87 211 117 155 103 130 103 1128 99 94 116 74 173 220 90 59	17 5 22 8 15 15 10 11 13 10 15 17 18 9 7 39 13 18 9 14 11 18 9	1 A 8 A 2:30A 5 A 1:30A 8 A 4 A 3:30A	3:30A 9 P 6:30A 12 P 8 P 9:30P 6 A 10:30P	.03 .81 .22 .44 1.12 .05 1.84	

^{*}Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

MONTHE 50		
JUNE, 1944	Normal	Extremes
Barometer, in inches (Readings reduced to sea level) Maximum 30.28, 5th, 9th Minimum 29.41, 24th Mean semi-daily 29.896 Range 0.87 Air Temperature, in degrees F. Highest 95, 29th Lowest 40, 8th Mean 66.3 Range 55 Highest mean daily 81, 29th Lowest mean daily 57, 8th	30.30 29.55 29.95 0.75 91.2 40.0 65.7 51.3	30.54, 1889 29.24, 1902 101, 1919 34, 1891
Mean maximum 77.6 Mean minimum 55.1 Greatest daily range 34, 8th Least daily range 6, 19th Precipitation, in inches		
Precipitation 4.70 Snow —	3.75	9.68, 1922; 0.76, 1908
Maximum precipitation in 24 hrs. 1.84, 24th Number of days with .01 or more 8	11	17, 1922; 4, 1908
Wind, in miles Total movement 3886 Greatest daily movement 240, 6th Least daily movement 45, 2nd	3585	4571, 1908; 1409, 1906
Least daily movement45, 2ndMean hourly velocity5.4Maximum velocity39, 16th	5.0 24.6	48, 1939
Wind, direction Prevailing direction WSW North, days 1 Northeast, days 1 East, days 0 Southeast, days 11 South, days 2 Southwest, days 1 West, days 4 Northwest, days 10	WSW	
Weather Mean relative humidity, percent	66.9	
Mean cloudiness, percent 70.00 Mean cloudiness, percent 61.3 Number of clear days 16 Number of fair days 66 Number of cloudy days 8 Number hours bright sunshine 279 Percent of possible hours of bright sunshine 61.1 Thunder and lightning 19th, 20th, 21st, 22nd, 24th Last frost May 19	51.1 10 12 8 257 54.0	71, 1903; 28, 1908 22, 1908, 1941, 1943; 1, 1927 23, 1912; 3, 1941 22, 1903; 1, 1923 362, 1908; 102, 1903

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1943.

The weather during June was near normal in all respects. The mean temperature for the month was 66.3 degrees which is only 0.6 of a degree higher than normal. The highest temperature was 95 degrees on the 29th and the lowest 40 degrees on the 8th. The domestic heating load during the month was 64 degree-days which is 4 less than normal. The domestic heating load for the 1943-44 season was 6799 degree-days while the normal is 6846 degree-days.

The total precipitation for the month was 4.70 inches, the normal for June being 3.75 inches. The total precipitation since January 1 is now 17.65 inches, while the normal for this period is 21.20 inches. There were 279 hours of bright sunshine compared to a normal of 257 hours. The mean relative humidity was 70.8 percent, the normal being 66.9 percent. The highest wind velocity was 39 miles per hour on the 16th.

Dr. J. K. Shaw gives the following report: "The dry period of May continued into June. Light rains on June 7th, 10th and 18th kept matters from growing worse, and during the period from June 19 to June 24, abundant rains removed any danger of drought injury for some time to come. The dry weather delayed germination of newly-planted seeds and plant setting, and injured shallow-rooted plants. The early hay crop was very short. The heavy rains came in season to increase growth of later fields and encourage a better second crop. Tree fruits did not suffer from lack of water.

"Further reports of the frost of May 19 indicate serious losses of the apple crop. Injury was confined to low frosty locations, while orchards on higher sites escaped any injury. Other tree fruits seem to have suffered but little. The crop of cultivated blueberries was very seriously injured and strawberries were extremely scarce."

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 667

July, 1944

Meteorological Observations

FOR

JULY

1944

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23' 31'' N. Longitude, 72° 31' 48'' W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION

AMHERST, MASS.

		Te	mpera	ture		ty	s		Wind			Precipita	tion	
	Max	imum	Min	imum		Iumidi				s				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8	83 84 83 89 89 88 91 93	4 P 5 P 2 P 1 P 3 P 2 P 3 P	53 57 51 57 56 64 60 64	5 A 5 A 5 A 4 A 4 A 4 A 5 A	68.0 70.0 67.5 70.0 72.5 76.0 75.5 78.5	49 47 60 67 69 62 66 63	15.3 15.3 8.8 7.7 12.5 13.2 15.2 15.2	nw nw sw sw sw nw s	117 122 56 71 67 109 59 93	14 10 10 11 14 13 8 7	2 A	10:30р	.56	
9 10 11 12 13 14 15	93 92 92 88 86 86 87	2 P 1 P 1 P 12 M 3 P 5 P 4 P	70 66 66 71 62 59 61	5 A 5 A 4 A 3 A 12 P 5 A 5 A	81.5 79.0 79.0 79.5 74.0 72.5 74.0	64 67 74 72 63 61 67	15.1 10.7 7.9 7.8 15.0 14.0 8.4	s se se se nw nw sw	110 91 88 188 141 40 58	10 11 15 18 14 4 7	2:30 P 8 P 12:30A	3 P 11 P 4 A	T .06 .22 .08	
16 17 18 19	87 79 81 82	1 P 4 P 4 P	60	12 P 12 P 5 A 4 A	73.5 67.0 66.5 68.0	77 58 61 63	5.6 15.0 15.0 11.6	se nw nw se	139 119 53 112	32 11 6 17	3 Р	4 P	.45 T	
20 21 22 23 24 25 26	79 77 82 86 90	3 P 1 P 6 P 4 P	61 55 50 57 61	3 A 12 P 3 A 5 A 5 A	70.0 66.0 66.0 71.5 75.5	84 68 57 58 65	3.6 7.2 14.9 14.8 14.8	se w w nw	90 114 113 55 79	10 18 15 10 12	5:30 A 0 A	12 P 3:30 A	.27 .15	
25 26 27 28	83 90 76 87	5 P 3 P 5 P	65 57	12 p 5 a 12 p	74.0 73.5 71.5 76.0	69 58 95 73	10.8 11.5 1.4 4.6	nw s nw se	121 70 62 34	14 17 15 6	2 A 10 P 2 A	2:30 A 10:30P 8 P	.04 .01 1.24	
28 29 30 31	87 80 86 86	2 P 12M 3 P 6 P	65 68 62	4 A 6 A 2 A 5 A	72.5 77.0 74.0	79 81 73	3.4 4.3 14.4	se se sw	78 65 33	15 25 6	1 р 12 м	3 P 5 P	.57 .23	

^{*}Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

July 1944	Normal	Extremes
Barometer, in inches (Readings reduced to sea level) Maximum	30.27 29.59 29.96 0.68	30.50, 1892 29.27, 1932 0.97, 1892; 0.47, 1938
Air Temperature, in degrees F. Highest 93, 8th, 9th Lowest 50, 22nd Mean 72.9 Range 43 Highest mean daily 81.5, 9th Lowest mean daily 66, 21st, 22nd Mean maximum 85.5 Mean minimum 60.4 Greatest daily range 33, 3rd, 5th, 26th Least daily range 9, 27th	93.9 46.4 70.8 47.7	104, 1911 40, 1890, 1898 74.7, 1921; 66.3, 1891
Precipitation, in inches Precipitation	4.10	14.51, 1897; 0.70, 1929 20, 1915; 4, 1924
	11	20, 1913, 4, 1924
Wind, in miles Total movement 2747 Greatest daily movement 188, 12th Least daily movement 33, 31st Mean hourly velocity 3.7	3422 4.6	5097, 1909; 1109, 1894
Maximum velocity	28.7	44, 1936
Wind, direction Prevailing direction SW North, days 0 Northeast, days 0 East, days 0 Southeast, days 9 South, days 5 Southwest, days 5 West, days 2 Northwest, days 10	SW	
Weather		
Mean relative humidity, percent 66.8 Mean cloudiness, percent 53 Number of clear days 19 Number of fair days 6 Number of cloudy days 6 Number hours bright sunshine 335 Percent of possible hours of bright sunshine 72.5 Thunder and lightning 4th, 11th, 16th 20th, 21st, 25th, 27th, 29th, 30th	68.4 50.9 10 14 7 268 58	70, 1943; 31, 1924 22, 1923; 0, 1915 24, '09, '21; 5, '89, '23, '28 '36 18, 1889; 0, 1910 371, 1910; 180, 1931

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1943.

The weather during July was warmer than usual, with more sunshine and less wind than normal. Eight heavy thunderstorms during the month brought much needed rainfall.

The mean temperature for the month was 72.9 degrees, the normal for July being 70.8 degrees. The highest temperature was 93 degrees on the 8th and 9th, and the lowest 50 degrees on the 22nd. A 21-day period beginning June 26 and ending July 16, when the maximum temperatures were in the 80's and 90's every day without a break, made a rather uncomfortable warm spell. In 1911 there was a 29-day period when the daily maximum temperatures, with the exception of one day, ranged from 80 degrees to 104 degrees. On three consecutive days during that period the afternoon temperatures reached 102.5, 104, and 101.5 degrees respectively. Other warm Julys were recorded in 1921 and 1935.

The total precipitation was 3.88 inches, compared with the normal for July of 4.10 inches. The total precipitation since the first of the year is now 21.53 inches, while the normal for this period is 25.30 inches.

There were 335 hours of sunshine, the normal being 268 hours. The total wind movement was 2747 miles as compared to the normal for July of 3422 miles.

Dr. Shaw makes the following comments: "The progress of vegetation during July was about normal. There were hot periods and, at times, lack of rain threatened to cause drought conditions. However, timely showers relieved the situation, and the month closed with enough moisture in the soil to maintain normal growth of crops for a week or more. Early varieties of apples matured somewhat earlier than normal. Hail on July 30 damaged crops a little to the north of us. The hay crop was generally small and there has not yet been enough rain to grow a very heavy second crop."

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 668

August, 1944

Meteorological Observations

FOR

AUGUST

1944

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23' 31'' N. Longitude, 72° 31' 48'' W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Te	mpera	ture	-	ty	s		Wind			Precipita	tion	
	Max	imum	Min	imum		Humidity	f Hour nshine			se				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative F	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8	89 85 85 96 94 81	2 P 3 P 5 P 4 P 2 P 3 P	60 68 67 64 70 70	5 A 5 A 12 P 3 A 5 A 7 A	74.5 76.5 76.0 80.0 82.0 75.5	70 84 82 78 67 77	10.9 6.2 8.7 10.8 7.0 6.4	w se ne s sw nw	30 55 97 26 89 122	10 12 5 12 13	7:30 A	9:00 a	.16	
7 8 9 10 11 12 13	71 81 85 93 97 99	1 P 5 P 5 P 4 P 3 P 3 P 2 P	62 57 49 57 63 68 72	5 A 6 A 5 A 6 A 6 A	66.5 69.0 67.0 75.0 80.0 83.5 84.0	80 58 61 68 66 66 83	0.7 14.2 14.2 14.2 14.1 14.1 10.3	nw nw sw sw s se se	124 134 43 81 66 126 109	10 14 4 10 10 11 8	2:30 A	3:00 A	.02	
14 15 16 17 18 19 20	94 93 95 94 88 75	12 M 5 P 3 P 1 P 5 P 3 P	74 67 73 70 59 50 45	4 A A A A A A P 12 P 6 A	84.0 80.0 84.0 82.0 73.5 62.5 62.0	87 82 79 79 71 67 53	3.7 10.7 6.4 8.7 9.3 13.8 13.7	e e e e w	97 55 88 87 134 134 62	36 7 10 25 25 17 7	1:00 P 2:00 P 6:00 A	4:30 P 10:00 P 6:30 A	2.18 T 1.72 .02	
21 22 23 24 25 26 27	85 83 80 73 72 74	2 P 4 P 3 P 3 P 3 P 2 P	58 65 60 56 45 43	5 A 6 A 6 A 6 A 5 A 6 A	71.5 74.0 70.0 64.5 58.5 58.5	65 83 62 79 64 64	7.7 5.3 9.0 3.5 13.5 13.5	se se se sw nw nw	153 132 107 156 84 101	13 9 25 17 12 15	10:00 P 0 A 11:30 P 2 A	11:30 P 3 A 12 P 4 P	.04 .04 .03 .12 T	
27 28 29 30 31	79 80 77 79 86	3 P 3 P 1 P 4 P 3 P	47 45 59 52 54	5 A 5 A 5 A 5 A 5 A	63.0 62.5 68.0 65.5 70.0	58 59 68 67 67	13.5 8.8 4.5 13.3 10.3	SW SW SW SW	107 50 128 150 51	11 5 13 18 7			Т	·

^{*}Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

August, 1944	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.36, 20th Minimum 29.63, 17th Mean semi-daily 29.986 Range 0.73	30.32 29.61 30.00 0.71	30.50, 1934 28.87, 1930
Air Temperature, in degrees F.		
Highest 99, 12th Lowest 43, 26th Mean 72.4 Range 56 Highest mean daily 84, 13th, 14th, 16th Lowest mean daily 58.5, 25th, 26th Mean maximum 85.1 Mean minimum 59.6 Greatest daily range 36, 9th, 10th Least daily range 9, 7th	91.6 43.4 68.6 48.1	100, 1918 37, 1894, 1908 34, 1940
Precipitation, in inches		
Precipitation 4.33 Snow —	4.08	8.40, 1928; .31, 1894
Maximum precipitation in 24 hrs2.18, 14th Number of days with .01 or more9	11	16, 1892, 1933; 4, 1899
Wind, in miles		
Total movement 2978 Greatest daily movement 156, 24th Least daily movement 26, 4th Mean hourly velocity 4.0 Maximum velocity 36, 14th	3127 4.2 22.7	4,271, 1910; 1,920, 1894 40, 1941
Wind, direction		
Prevailing direction SW North, days 0 Northeast, days 1 East, days 4 Southeast, days 6 South, days 2 Southwest, days 8 West, days 2 Northwest, days 8	SW	
Weather		
Mean relative humidity, percent 70.8 Mean cloudiness, percent 55 Number of clear days 17 Number of fair days 11 Number of cloudy days 3 Number hours bright sunshine 301 Percent of possible hours of bright	70.2 49.7 9 13 9 237	67, 1901; 27, 1923 23, 1941; 0, 1915, 1929 25, 1912; 3, 1939 18, 1901, '28; 2, 1910, '23 332, 1941; 152, 1915, 1929
sunshine	55.2	
Thunder and lightning 14th, 17th, 22nd First Frost	Sept. 21	\ug. 22, '94, '95; Oct. 13, '09

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1943.

The weather during August was marked with several extremes. The first three weeks were very warm with 10 days on which the temperature exceeded 90 degrees, the hottest being 99 degrees on August 12. This has been exceeded only once in August in the history of this station. This was in 1918 when the maximum temperature was 100 degrees. The weather cooled off during the last week in August so that the mean temperature for the month was 72.4 degrees. This has been exceeded only twice since 1889; in 1937 when the mean for August was 73.3 degrees and 1939 when it was 73.5 degrees. The mean temperature for the three summer months of the past season was 70.5 degrees. This equals the previous record for hot summer months which was reached in 1939.

The rainfall during the month was 4.33 inches. Of this amount 2.18 inches fell on the afternoon of the 14th. During part of this storm rain fell at a greater intensity than had previously been recorded at this station. A total of 2.02 inches fell in one-half hour and 1.10 inches fell in 10 minutes. The rain on the 14th was followed by another heavy rain on the 17th when 1.72 inches fell. There were 301 hours of bright sunshine, the normal being 237 hours. The mean relative humidity for the month was 70.8 percent. The highest wind velocity was 36 miles per hour on the 14th.

Dr. J. K. Shaw makes the following comments: "The weather during August was characterized by extremely hot periods and it was on the dry side much of the time. The month closes with the soil pretty dry. Timely rains broke the drought before disastrous results took place but many crops suffered injury. Fruit plants stood the dry weather with little injury; and apples, peaches, and other fruits are sizing and coloring well. Except in orchards on low, frosty sites, where the freeze of May 19 was disastrous, all tree fruits promise good crops of excellent fruit."

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 669

September 1944

Meteorological Observations

FOR

SEPTEMBER

1944

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23' 31'' N. Longitude, 72° 31' 48'' W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Te	mpera	ture		ty	S		Wind		Precipitation				
	Max	imum	Minimum		Minimum			Iumidi	r of Hours Sunshine			တ္			
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number o Bright Su	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water		
1 2 3	83 89 81	12 M 1 P 3 P 12 M	65 65 61 53	5 A 11 P 12 P 5 A	74.0 77.0 71.0 66.0	81 69 66 72	2.7 9.8 13.1 8.4	se w nw	69 87 158 59	7 13 15 8	5 F 9:30 F 0 A	10 p	.35		
2 3 4 5 6 7 8 9	79 76 81 77 71 67 76	12 M 1 P 3 P 1 P 3 P 2 P 3 P	62 58 53 51 49	12 P 4 A 12 P 5 A 12 P 6 A	69.0 69.5 65.0 61.0 58.0 60.0	90 76 65 64 73 71	1.9 5.6 12.9 7.2 2.6 12.8	sw sw se w w	55 62 185 191 81 49	8 12 20 22 10 7	2:30 F	3 P	.22		
11 12 13 14 15 16 17	81 70 63 76 84 80 82	1 P 1 P 4 P 3 P 1 P 2 P 3 P	45 50 57 62 61 55 56	6 A 5 A 0 A 12 P 5 A 5 A	63.0 60.0 60.0 69.0 72.5 67.5 69.0	67 80 98 98 74 78 72	12.8 0.4 0.0 0.0 6.5 6.4 12.6	nw nw ne w nw e	39 54 109 205 191 40 75	8 8 55 48 10	5:30 F 4 A 1 A 0 A	10 p 12 p			
18 19 20 21 22 23 24 25 26 27	71 72 74 78 69 59 64	4 P 2 P 1 P 2 P 4 P 2 P 2 P 3 P	58 53 64 61 47 36 33	12 P 5 A 12 P 12 P 12 P	64.5 62.5 69.0 69.5 58.0 47.5 48.5	81 92 89 88 57 66 74	6.4 1.0 0.0 0.8 11.2 12.1 12.1	e sw se se w nw sw	35 38 126 226 183 50	5 4 13 28 20	10 г	11 P	.41		
26 27 28 29 30	67 74 74 68 65 66	3 P 1 P 2 P 1 P 0 A 3 P	34 44 52 60 41 36	5 A 5 A 6 A 2 A 12 P 6 A	50.5 59.0 63.0 64.0 53.0 51.0	66 71 78 93 71 67	9.0 7l 9.4 0.0 12.0 9.9	e e se se w nw	55 105 70 131 281 94	9	1 P 0 A	11 p 12:30 a	.31		

^{*}Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

September, 1944	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.53, 18th Minimum 29.13, 14th Mean semi-daily 30.029 Range 1.40	30.45 29.57 30.06 .88	30.65, 1924 28.41, 1938 1.99, 1938; .57, 1910
Air Temperature, in degrees F.		, , , ,
Highest 89, 2nd Lowest 33, 24th Mean 63.0 Range 56 Highest mean daily 77.0, 2nd Lowest mean daily 47.5, 23rd Mean maximum 73.9 Mean minimum 52.2 Greatest daily range 36, 11th Least daily range 6, 13th	87.7 33.2 61.7 54.5	97, 1895, 1929 24.5, 1914
Precipitation, in inches		
Precipitation 5.31 Snow	4.24	14.55, 1938; .52, 1914
Maximum precipitation in 24 hrs. 2.66, 14th Number of days with .01 or more11	10	16, 1933; 3, 1903
Wind, in miles		
Total movement 3152 Greatest daily movement 321, 15th Least daily movement 35, 19th Mean hourly velocity 4.4 Maximum velocity 55, 14th	3271 4.5 25.9	4,686, 1896; 1,414, 1894 80, 1938
Wind, direction		
Prevailing direction WSW North, days 0 Northeast, days 1 East, days 4 Southeast, days 6 South, days 0 Southwest, days 5 West, days 8 Northwest, days 6	WSW	
Weather		
Mean relative humidity, percent 76.2 Mean cloudiness, percent 56 Number of clear days 13 Number of fair days 7 Number of cloudy days 10 Number hours bright sunshine 207 Percent of possible hours of bright sunshine 55.5 Thunder and lightning 1st, 2nd, 21st First Frost Sept. 21	10 10 10 201 54.9	70, 1934; 27, 1908, 1914 19, 1932; 2, 1907, 1938 19, 1908; 3, 1880 20, 1934; 3, 1041 255, 1916; 106, 1934 Aug. 22, '94, '95; Oct. 13, '09

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1880 to 1943

The hurricane on the 14th and 15th was the outstanding feature of the weather during September. The storm was much less severe in Amherst than the hurricane of 1938 and comparatively little damage was done in this section. The highest wind velocity was 55 miles per hour at 10:45 P. M. (E. S. T.) on the 14th. The direction ranged from north to northeast. The barometer reached its lowest reading of 29.13 inches at 10:30 P. M. The total rainfall from the 12th to the 15th was 3.76 inches.

The mean temperature during the month was 63 degrees while the normal mean temperature for September is 61.7 degrees. The lowest temperature was 33 degrees on the 24th. There were no killing frosts during the month, although there were light frosts on low ground on the 24th and 25th. The domestic heating load during the month was 122 degree-days compared to a normal of 149 degree-days in September.

The rain fall during the month was 5.31 inches, the normal being 4.24 inches. The total precipitation since January 1 is 31.17 inches, while the normal precipitation for this period is 33.62 inches. There were 207 hours of bright sunshine which is 6 hours more than normal for September.

Dr. J. K. Shaw, research pomologist, makes the following comment: "While the hurricane on September 14 was by no means so severe as that of 1938, it ruined some shade and orchard trees and blew off from 20 to 50 percent of the McIntosh apples remaining on the trees. Later varieties suffered less but some of these fell. The taller farm crops still in the field were injured. Corn fields were much disturbed. Hurricane damage was worse in the eastern part of the state where a loss up to 75 percent of the McIntosh still on the trees was reported. Otherwise, the progress of vegetation was about normal. Most fruits ripened a little earlier than usual, perhaps due to abnormally warm weather during the summer."

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 670

October 1944

Meteorological Observations

FOR

OCTOBER

1944

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Те	mpera	ture		ty	s	Wind Precipitation						
	Max	imum	Min	Minimum		fumidi		20		es				
Day	Degrees	Time	Degrees	Time	Mcan	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 3 4 4 5 5 6 6 7 7 8 8 9 10 11 12 13 13 14 15 16 6 17 18 19 20 21 22 23 24 25 26 27 8 29 30 31	67 63 64 67 72 67 85 77 77 71 70 68 69 49 56 64 64 53 46 62 47 53 58 58 59	3 P 1 P 1 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2	38 39 33 329 35 57 59 49 39 35 50 31 27 37 35 34 29 41 34 29 41 42 36 43 42 35 30 30 43 43 42 43 43 43 43 43 43 43 44 43 43 44 43 43	7 A 6 A P 12 P 12 P 7 A A 12 P 12 P 12 P 7 A A 12 P 12 P 12 P 7 A A 12 P 12 P 12 P 7 A A 12 P 12 P 12 P 7 A A 12 P 12 P 12 P 7 A A 12 P 12 P 12 P 7 A A 12 P 12 P 12 P 12 P 12 P 12 P 12 P	52.5 51.0 48.5 48.0 53.5 62.0 63.0 57.0 59.5 50.5 50.5 54.5 41.0 41.5 55.5, 42.5 55.5 42.5 54.0 54.0 54.0 54.0 54.0 54.0 54.0 54	72 67 58 70 76 97 76 78 81 76 78 79 97 62 63 64 65 58 74 96 563 71 92 73 61 45 60 51 70	11.8 11.7 2.2 11.6 0.0 7.3 5.0 6.2 7.7 7.5 0.0 1.0 8.9 3.8 11.0 7.9 2.1 0.0 7.3 10.8 4.2 10.6 10.5 10.5	nw nw nw nw e se se sw se sw sw w w se nw nw nw w se e w nw nw w se se e w nw nw se	114 209 104 55 62 132 109 52 38 69 67 51 44 77 306 119 98 66 233 59 256 243 107 123 81 135 324 431 214	13 6 5 11 13 10 5	8 A 2:30 A 10 A 10:30 P 0 A	3 A	.26	

^{*}Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

	1	
October, 1944	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.66, 31st Minimum 29.50, 14th Mean semi-daily 30.064	30.53 29.42 30.06	30.68, 1929 29.00, 1926
Range 1.16	1.11	1.47, 1926; .76, 1899
Air Temperature, in degrees F.		
Highest 85, 7th Lowest 24, 31st Mean 50.00 Range 61 Highest mean daily 72, 7th Lowest mean daily 40, 15th Mean maximum 62.2 Mean minimum 37.8 Greatest daily range 38, 4th Least daily range 5, 21st	79.4 23.4 50.5 56.0	90.5, 1908 17, 1936 56.4, 1920; 43.2, 1890
Precipitation, in inches		
Precipitation 1.74 Snow Maximum precipitation in 24 hrs.	3.29	8.81, 1911; .01, 1924
Number of days with .01 or more5	9	15, 1913; 1, 1897, 1924
Wind, in miles		
Total movement 4396 Greatest daily movement 431, 29th Least daily movement 38, 9th Mean hourly velocity 5.9 Maximum velocity 32, 29th	5.4	5,467, 1910; 2,540, 1894 42, 1937
Wind, direction	27.0	T_, 1737
Prevailing direction SW North, days 0 Northeast, days 0 East. days 2 Southeast, days 8 South, days 2 Southwest, days 4 West, days 7 Northwest, days 8		
Weather		
Mean relative humidity, percent 71.6 Mean cloudiness, percent 53 Number of clear days 16 Number of fair days 9 Number of cloudy days 6 Number hours bright sunshine 225 Percent of possible hours of bright sunshine 660	48 10 10 11 175.0	67. 1932: 18. 1924 21. 1938; 1. 1911 17. 1924; 3. 1938 19. 1896; 3. 1924 232, '23, '38; 91, 1913
Thunder and lightning 7th, 8th, 9th First frost Oct. 4	Sept. 21	Aug. 22, '94, '95; Oct. 15, '39

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1943.

The weather during October was dry and clear, with the temperature very near normal. Rain fell on only five days, with a total fall of 1.74 inches, whereas the normal for October is 3.29 inches. The rainfall during the past three months—August, September and October—was 11.38 inches, while the normal rainfall for this period is 11.61 inches. There were 225 hours of bright sunshine compared to a normal of 175 hours.

The mean temperature for the month was 50.0 degrees. The normal mean temperature for October is 50.5 degrees. The highest temperature was 85 degrees on the 7th and the lowest, 24 degrees on the 31st. The domestic heating load was 472 degreedays, the normal for October being 456 degreedays. The total heating load for September and October this year was 594 degreedays, while the normal for these two months is 605 degreedays.

The total wind travel during the month was 4396 miles, the normal for October being 4074 miles. A maximum velocity of 32 miles per hour was recorded on the 29th.

The mean relative humidity was 71.6 percent, while the normal for October is 69.0 percent.

Correction: A typographical error on page 3 in the September bulletin gives September 21 as the date of the first frost in 1944. There was no frost at this station in September. The date "September 21" should appear in the "Normal" column.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series Bulletin No. 671

November 1944

Meteorological Observations

FOR

NOVEMBER

1944

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

> Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

	Temper ature					ty	v	Wind Precipitation										
	Max	imum	Minimum		Minimum		Minimum			Iumidi	f Hour nshine			ສ				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow				
1 2 3 4	66 66 68 64	4 P 3 P 4 P 3 P	45 42 39 46	3 A 12 P 1 A 7 A	55.5 54.0 53.5 55.0	84 86 88 91	8.7 8.0 8.1 4.5	se se se sw	99 50 36 56	11 5 7 6			Т					
2 3 4 5 6 7 8 9	54 39 46 50	1 P 2 P 2 P 11 A 3 P	36 33 33 28 25	12 P 9 A 7 A 12 P 3 A	45.0 36.0 39.5 39.0 34.0	62 76 56 66 83	10.0 0.1 9.3 3.8 2.1	nw nw	222 341 335 92 13	22 22 20 15 1	10:30 A	2 Р	.04	.25				
10 11 12 13	43 50 52 45 49	9 P 10 A 3 P 3 P	41 32 28 22	0 A 12 P 12 Р 7 А	45.5 42.0 36.5 35.5	97 72 60 63	0.0 0.3 9.9 9.9	W W W nw	11 174 253 67	1 25 20 17	3 A 7 A	6 р 8 а	.73 .02					
14 15 16 17 18	53 42 39 44 40 48	3 P 3 P 3 A 3 P 1 P 3 P	23 28 32 32 34 27	6 A 2 A 5 P 0 A 12 P 12 P	38.0 35.0 35.5 38.0 37.0 37.5	63 75 93 76 59 67	9.8 0.7 0.0 0.7 0.0 9.6	mw nw nw	60 25 129 239 131 104	13 4 15 14 10 13	11 P 4 A	12 P 9 P	.03 .90	.50				
20 21 22 23 24 25 26 27 28	43 42 38 40 39 38	7 P 0 A 1 P 3 P 2 P	23 36 29 24 19 28	7 A 11 P 12 P 12 P 7 A 12 P	33.0 39.0 33.5 32.0 29.0 33.0	84 79 90 66 78 59	1.7 0.0 0.0 7.0 4.7 3.7	n n nw sw nw	20 297 224 88 133 288	7 22 18 17 22 25 22	7 P 6 A	8 P 1 P	.01	Т				
26 27 28 29 30 31	52 46 50 33 39	2 P 3 P 9 P 11 A 12 P 1 P	23 24 29 28 30	7 A 1 A 12 P 1 A 12 P	37.5 35.0 39.5 30.5 34.5	68 95 73 81 85	9.3 0.0 5.1 0.0 0.0	w se nw nw s	131 76 134 124 278	22 11 20 9 28	6 A 1 A 8 A 0 A	12 P 1:30 A 12 P 2 P	.70 .02 .05 1.68					

^{*}Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

	1 1	
November, 1944	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.56, 2nd Minimum 28.87, 30th Mean semi-daily 30.005	30.59 29.30 30.05	30.87, 1932 28.73, 1904
Range 1.69	1.29	1.84, 1904; .89, 1943
Air Temperature, in degrees F.		
Highest 68, 3rd Lowest 19, 24th Mean 39.0 Range 49 Highest mean daily 25.5, 1st Lowest mean daily 29.0, 24th Mean maximum 47.3 Mean minimum 30.6 Greatest daily range 30, 14th Least daily range 5, 29th		75, 1924, 1938 —1, 1938 44.1, 1931; 33.6, 1901
 Precipitation, in inches 		
Precipitation 4.21 Snow 0.75 Maximum precipitation in 24 hrs.		8.64, 1927; .63, 1917 13.50, 1938
Number of days with .01 or more11	9	22, 1921; 2, 1904
Wind, in miles		# 0#0 100¢ 0 #00 1000
Total movement 4230 Greatest daily movement 341, 6th Least daily movement 11, 10th Mean hourly velocity 5.9 Maximum velocity 28, 30th	6.3	5,978, 1906; 2,589, 1889 44, 1938
Wind, direction	00.0	14, 1700
Prevailing direction WNW North, days	3 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
Weather	50.6	
Mean relative humidity, percent 75.8 Mean cloudiness, percent 66 Number of clear days	55.1 6 9 4 15 7 121	72, 1900, 1927; 34, 1917 15, 1903; 1, 1900, 1911, 1927 16, '12; 4, '89, '30, '32, '35 24, 1927; 9, 1905, 1917 182, 1903; 66, 1927
sunshine 43. Thunder and lightning 64	_	Oct. 10, '25; Nov. 27, '31
First snow	Nov. 6	Oct. 10, 23, NOV. 27, 31

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, everyt that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1943

REMARKS

The weather during November was near normal in most respects. The mean temperature for the month was 39 degrees which is one tenth of a degree higher than normal. The highest temperature was 68 degrees on the 3rd and the lowest was 19 on the 24th. The normal minimum for November is 12.9 degrees. In 1938 the temperature dropped to -4 degrees in November. The domestic heating load for November was 782 degree-days compared to a normal of 784 degree-days. The total heating load for the season to the end of November is 1376 degree-days, while the normal for the period is 1389. This represents 29 percent of the normal year's heating load.

The total precipitation during the month was 4.21 inches. Of this amount, 1.73 inches fell during the storm on the 29th and 30th. This storm which was quite severe on the coast gave only moderate wind velocities at this station. The highest velocity recorded was 28 miles per hour. It was accompanied with a very low barometer. The minimum was 28.87 inches on the 30th. The normal precipitation for November is 3.41 inches. The first snow of the season fell on November 8 when 0.25 of an inch fell. This is the normal date for the first snow. Again on the 18th, 0.50 of an inch of snow fell. The normal snowfall for November is 2.34 inches. There were 127 hours of bright runshine, the normal being 121 hours. The mean relative humidity was 75.8 percent, while the normal for November is 70.3 percent.

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorogical Series

Bulletin No. 672

December 1944

Meteorological Observations

FOR

DECEMBER

1944

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23' 31'' N. Longitude, 72° 31' 48'' W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Те	mpera	ture		ty	ø		Wind			Precipita	ition	
	Max	imum	Min	imum		Iumidi	r of Hours Sunshine			s 41				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Bright Su	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 4 5 5 6 6 7 8 9 100 111 12 13 114 115 116 117 118 119 200 221 223 224 225 229 300 31	33 27 26 37 49 49 44 41 40 31 33 38 42 20 29 33 32 38 38 37 18 31 19 28 32	11 A	27 12 11 14 18 18 20 29 30 30 30 22 28 22 17 18 29 31 20 1 -3 11 21 21 21 21 21 21 21 21 21 21 21 21	4 A A A A A A A A A A A A A A A A A A A	30.0 19.5 18.5 25.5 33.5 28.5 34.5 37.0 35.5 31.0 34.0 26.5 25.0 28.0 35.5 36.5 27.0 10.5 12.5 29.5 33.0 24.5 29.5 33.0 24.5 29.5 33.0 24.5 29.5 33.0 24.5 25.0 26.5 27.0 27.0 27.0 27.0 27.0 27.0 27.0 27.0	69 55 54 54 76 83 69 64 79 69 77 72 73 77 78 69 81 79 92 469 53 47 52 80	0.8 4.4 9.3 9.2 9.2 3.7 8.4 0.0 2.7 0.0 6.8 0.0 9.2 2.8 4.3 2.0 5.1 5.7 9.0 0.0 9.2 2.7 9.2 9.1 0.0	sw nw nw nw nw nw n ne w w w se se se sw nw nw se se se w w w se se se w nw nw nw se se w nw nw nw nw nw nw nw nw nw nw nw nw n	409 269 263 225 44 77 96 210 229 64 260 316 118 131 163 262 188 179 135 270 69 34 33 30 314 47 331 150 44 29	22 20 18 5 17 25 22 20 14 28 25 17 12 25 15 13 17 25 15 13 16 5 5 5 17 25 18 18 18 18 18 18 18 18 18 18 18 18 18	1 P 1 A 2 A 4 A 6 P 0 A 9 A	11:30p 2 A 8:30 p 1:30 p 10 p 12 p 4 A	.27 .05 .32 .49 .40 .32	.25 2.50 .50 2.00 4.00 T

^{*}Based on least time required to blow one mile.

FERNAND E. BARTLETT. Observer

December, 1944	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.71, 27th Minimum 29.17, 13th Mean semi-daily 30.084 Range 1.54	30.66 29.27 . 30.06 1.38	30.96, 188 28.85, 191 1.78, 1895; 1.01, 189
	1.00	1.70, 1093 , 1.01, 109
Air Temperature, in degrees F. Highest 49, 5th, 7th, 8th Lowest -6, 22nd Mean 25.3 Range 55 Highest mean daily 39, 8th Lowest mean daily 8.5, 22nd Mean maximum 34.4 Mean minimum 16.2 Greatest daily range 32, 20th Least daily range 6, 1st	54.3 -1.9 27.5 56.2	65.5, 190 -22.5, 191 36.9, 1891; 17.1, 191
Precipitation, in inches		
Precipitation 2.18 Snow 9.25 Maximum precipitation in 24 hrs.	3.39 8.50	7.77, 1901; .58, 194 26.50, '02; t, 1891, 194
Number of days with .01 or more	10	17, 1902; 4, 1892, 194
Wind, in miles		
Total movement 4936 Greatest daily movement 409, 1st Least daily movement 7, 6th Mean hourly velocity 6.6 Maximum velocity 48, 28th	4710 6.3 31.4	6,694, 1925; 3,239, 191 48, 1938, 194
Wind, direction		10, 1200, 171
Prevailing direction W North, days 1 Northeast, days 1 East, days 0 Southeast, days 7 South, days 1 Southwest, days 2 West, days 11 Northwest, days 8	WNW	
Weather		
Mean relative humidity, percent 66.7 Mean cloudiness, percent 54 Number of clear days 12 Number of fair days 6 Number of cloudy days 13 Number hours bright sunshine 145 Percent of possible hours of bright sunshine 51.2	69.4 54.9 8 9 14 128 45.2	71, 1929; 39, 1919 15, 1890; 2, 1922, 193, 16, '09; 4, '89 '30, '31, '36, '36 23, '33; 7, '09, '2, 172, 1896; 63, 193,

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1943

ANNUAL SUMMARY

Annual, 1944	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.73, Mar. 22 Minimum 28.87, Nov. 30 Mean semi-daily 30.031 Range 1.86	30.81 28.95 30.01	31.05, 1920, 1943 28.41, 1938
	1.85	2.47, 1938; 1.38, 1933
Air Temperature, in degrees F.		
Highest 99, Aug. 12 Lowest -6, Dec. 22 Mean 48.4 Range 105 Highest mean daily 84.0, Aug. 13, 14, 16	95.7 -12.2 47.4 107.8	104, 1911 -26, 190- 49.8, 1921; 44, 190-
-6, Dec. 22 Mean 48.4 Range 105 Highest mean daily 84.0, Aug. 13, 14, 16 Lowest mean daily 8.5, Dec. 22 Mean maximum 59.0 Mean minimum 37.7 Breatest daily range 48, Feb. 14 Least daily range 2, Feb. 27	57.8 36.5	
Precipitation, in inches		
Precipitation 39.30 Snow 31.50 Maximum precipitation in 24 hrs.	43.70 47.78	59.00, 1938; 30.68, 1908 89.00, 1893; 24.50, 1919
2.66, Sept. 14 Number of days with .01 or more 107	124	144, 1902; 96, 1924
Wind, in miles		
Total movement 48,557 Greatest daily movement 484, Mar. 1 Least daily movement 3, Jan. 18 Mean hourly velocity 55, Sept. 14	52,223	63,571, 1908; 36,257, 1894
Mean hourly velocity	5.8 39.5	80, 1938
Wind direction		
Prevailing direction W North, days 9 Northeast, days 6 East, days 13 Southeast, days 82 South, days 25 Southwest, days 37 West, days 76 Northwest, days 118	W	
Weather	C7 C	
Mean relative humidity, percent 68.3 Mean cloudiness, percent 56.5 Number of clear days 189 Number of fair days 79 Number of cloudy days 98 Number hours bright sunshine 2772 Percent of possible hours of bright		60, '98, '01, '02; 41, '08, '24 217, 1941; 59, 1927 182, 1912; 64, 1936 179, '01, '02; 71, 1910, 1941 3038, 1941; 1864, 1902
sunshine 62.2 Last snow Mar. 29 First snow Nov. 6 Last frost May 19 First frost Oct. 4	52.8 April 15 Nov. 6 May 14 Sept. 21	Mar. 14, '10; May 11, '07 Oct. 10, '25; Dec. 13, '41 Apr. 14, '43; June 8, '32 Aug. 22, '94, '95; Oct. 13, '09

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 673

January 1945

Meteorological Observations

FOR

JANUARY

1945

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Te	mpera	ture		ţ.	os.		Wind			Precipit	ation	
	Max	mum	Min	imum		Iumidi	r of Hours Sunshine			s.				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Bright Sur	Prevailing Direction	Total Movement	Max, Miles per Hour*	Ведан	Ended	Water	Snow
1 2	55 34	5 P	30 12	0 A 12 P	42.5 23.0	84	0,5 6,7	11	151 295	25 32 15	0.5	12 P	1.05	
3 4 5	26 38 27	3 P 3 P 0 A	12 7 21 11	8 A 1 A 12 P	16.5 29.5 19.0	52 71 43	3.6 0.2 9.3	se se	127 75 229	22	8	11 A	.06	1.00
2 3 4 5 6 7 8 9	17 19 32 32	3 P 12 P 2 P 1 P	4 6 19 6	8 A 7 A 0 A 12 P	10.5 12.5 25.5 19.0	40 80 82 73	9.2 0.0 3.0 3.0	mw. m mw.	261 159 77 117	18 17 17 18	4.7	8 r	.38	4.50
10 11 12 13	12 19 33 35	3 P 2 P 8 P 1 P	-4 -8 -5 16	12 P 3 A 2 A 12 P 8 A	4.0 5.5 14.0 25.5 16.5	45 62 71 69	9.3 6.2 0.0 0.1	W SW SC DW	190 189 135	18 5 20 14	3 р	5 г	.01	Т
14 15 16 17	20 31 23 24	12 P 3 P 12 P 3 A	13 6 17 14	8 \ 7 \ 1 \ 12 P	16.5 18.5 20.0 19.0	81 81 83 61	70 00 48	13// 11// 11/// 11///	285 184 253 337	20 13 22	6 x 5 x	12 p	.38	6,07) T 3,50
18 19	31 32 38 32 29	3 P 3 P 3 P 3 P	5 10 0 2 7	9 A 12 P 7 A 7 A	18.0 21.0 19.0 17.0 18.0	52 51 57 66 79	9.5 9.5 9.6 6.2 4.0	nw n nw s	145 80 48 38 43	17 14 8 4 8	2 p	8 P	.15	1.50
20 21 22 23 24 25 26 27 28 29 30	36 27 8 14	12 m 3 P 3 P	14 -3 -10 -4	1 A 12 P 7 A 5 A	25.0 12.0 -1.0 5.0 16.5	67 62 49 43	78 60 97 98	nw nw	158 288 316 260	25 45 32 28				
27 28 29 30 31	27 37 29 28 22	1 P 2 P 1 P 3 P 1 P	6 6 18 13 9	3 A 7 A 12 P 7 A 6 A	16 = 21.5 = 23.5 = 20.5 = 15.5	17 10 77 57 59	8.6 4.2 10 0 9 0	//. ///. ///. 26 ////.	336 112 164 232 150	25 20 25 25 25 22	0.3	11 A	.70	7.00

^{*}Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

January, 1945	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.44, 3rd Minimum 28.98, 1st Mean senii-daily 29.998 Range 1.46	30.70 29.20 30.07 1.50	31.00, 192; 28.55, 191; 2.18, 1913; 0.97, 1896
Air Temperature, in degrees F.		
Highest 55, 1st Lowest -10, 25th Mean 17.8 Range 65 Highest mean daily 42.5, 1st Lowest mean daily -1.0, 25th Mean maximum 28.0 Mean minimum 7.7 Greatest daily range 38, 12th, 20th Least daily range 6, 16th	51.1 -7.1 24.2 58.2	66.0, 193 -26.0, 190 34.2, 1913; 13.9, 191
Precipitation, in inches		
Precipitation 3.07 Snow 23.50	3.61 13.32	7.15, 1898; 1.07, 189 33, 1898, 1923; 1.50, '0
Maximum precipitation in 24 hrs. 1.05, 1st Number of days with .01 or more 8	11	15, '93, '20; 4, 190
Wind, in miles		
Total movement 5471 Greatest daily movement 337, 17th Least daily movement 29, 11th Mean hourly velocity 7.4 Maximum velocity 45, 24th	5055 6.8 32.7	7770, 1908; 2806, 180; 47, 193
Wind, direction	,,_,,	77, 19,0
Prevailing direction WNW North, days 1 Northeast, days 0 East, days 0 Southeast, days 4 South, days 1 Southwest, days 1 West, days 11 Northwest, days 13	WNW	
Weather		
Mean relative humidity, percent 62.6 Mean cloudiness, percent 51.8 Number of clear days 14 Number of fair days 10 Number of cloudy days 7 Number hours bright sunshine 177 Percent of possible hours of bright sunshine 60.2 Thunder and lightning 29th	70.1 55 9 13 137 46.7	78, 1932; 37, 1908 20, 1944; 2, 191- 18, 1800, 1926; 3, 1916, 194- 22, 1923, 1931; 4, 1920, 1936 214, 1920; 74, 1932

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1944.

REMARKS

January produced severe winter weather. The mean temperature was 17.8 degrees which is the coldest January since 1920. In that year the mean temperature in January was 15.2 degrees. In only four other years since this station was established in 1889 has January been colder than the present year. On six days the temperature dropped below zero and on only 8 days was the maximum above 32 degrees. The lowest temperature was -10 degrees on the 25th. The heating load was 1462 degree-days whereas the normal for January is 1265 degree-days. The total heating load for the season to the end of January is 4070 degree-days, the normal being 3817 degree-days. The heating load for the season to the end of January is nearly 7 percent higher than normal.

The snowfall during the month was 23.50 inches compared to a normal of 13.32 inches. The snowfall for the winter is 33.50 inches; the normal to the end of January is 24.16 inches. The heavy snowfall in January was equalled in 1904 and has been exceeded only 3 times since 1889. The total precipitation was 3.07 inches, the normal being 3.61 inches. There were 177 hours of bright sunshine, the normal being 137 hours. The highest wind velocity was 45 miles per hour on the 24th.

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 674

February 1945

Meteorological Observations

FOR

FEBRUARY

1945

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23' 31'' N. Longitude, 72° 31' 48'' W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

-		Те	mpera	ture		ty	w		Wind			Precipit	ation	
		imum		imum		Mean Relative Humidity	Number of Hours Bright Sunshine	ng n	nt	iles r*				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative	Number Bright	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8 9	23 21 32 33 33 37 39 32 44 38	3 P 3 P 3 P 2 P 4 P 4 P 3 P 4 A 3 P	5 11 5 3 6 -1 2 23 22 19	7 A 7 A 8 A 8 A 12 P 7 A 4 A 12 P 7 A 0 A	14.0 16.0 18.5 18.0 19.5 18.0 20.5 27.5 33.0 28.5	52 53 59 60 75 71 87 77	10.0 10.0 10.0 6.8 8.2 10.1 7.2 0.0 8.0 2.5	nw w nw sw nw se nw nw nw	226 351 196 48 179 31 80 213 212 24 154	28 28 18 7 18 7 12 20 22 9 20	9 P 0 A	12 P 8 A	.09 .21	1.00 3.00 7.00
11 12 13 14 15 16 17 18 19 20 21	43 39 25 26 38 42 35 29 29 30 39	3 P 3 P 4 P 3 P 11 A 1 P 11 A 3 P 3 P 4 P	28 23 14 21 18 30 17 4 5 -1 10	10 P 8 A 7 A 11 P 4 A 0 A 12 P 7 A 7 A 5 A 7 A	35.5 31.0 19.5 23.5 28.0 36.0 26.0 16.5 17.0 14.5 24.5	57 56 68 80 78 57 51 66 65 73	10.4 5.5 0.0 2.1 3.4 7.0 4.0 10.6 10.7 3.5 8.0	W W NW NW Se W W W W	124 94 177 65 232 200 121 133 59 86	18 13 13 11 28 28 20 12 7	6 а 8 р 0 а	4 P 12 P 8 A	.05	1.00 1.00 3.00
20 21 22 23 24 25 26 27 28	38 42 37 41 43 43 42	12 P 3 P 1 P 3 P 12 P 0 A 3 P	28 34 31 23 20 28 20	1 A 8 A 6 A 12 P 3 A 8 P 7 A	33.0 38.0 34.0 32.0 31.5 35.5 31.0	97 75 60 53 89 54 52	0.0 3.0 5.4 11.0 0.0 4.2 9.4	Se W W SW W Se	90 126 274 240 50 341 151	20 20 32 22 11 40 22	12:30 P 2 A 9:30 P	2 P 7 A 9:30 P 5:30 A 12 P	1.31 .02 .62 .05	0.50

^{*}Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

February, 1945	Normal	Extremes
Barometer, in inches (Readings reduced to sea level) Maximum	30.63 29.24 30.04 1.38	31.05, 192(28.56, 189)
Air Temperature, in degrees F.	1.30	1.89, '00, '08; .88, '13, '3'
Highest 44, 9th Lowest -1, 6th, 20th Mean 25.7 Range 45 Highest mean daily 38, 23rd Lowest mean daily 14, 1st	50.5 -7.4 23.7 58	65.0, 193 -23, 194
Lowest mean daily 14, 1st Mean maximum 35.5 Mean minimum 16.0 Greatest daily range 38, 6th Least daily range 5, 14th	32.8 14.5	
Precipitation, in inches		
Precipitation 3.33 Snow 16.50 Maximum precipitation in 24 hrs.	3.19 13.99	8.12, 1900; 0.62, 190 48.75, 1893; 0.50, 193
Number of days with .01 or more 12	10	15, '93, '20; 4, 190
Wind, in miles		
Total movement 4277 Greatest daily movement 351, 2nd Least daily movement 24, 10th Mean hourly velocity 6.4 Maximum velocity 32, 24th	7.1 31.7	6445, 1896; 3438, 189 48, 1934, 1937, 194
Wind, direction		
Prevailing direction W North, days 1 Northeast, days 0 East, days 2 Southeast, days 4 South, days 0 Southwest, days 2 West, days 11 Northwest, days 8	WNW	
Weather		
Mean relative humidity, percent 67.5 Mean cloudiness, percent 58 Number of clear days 12 Number of fair days 6 Number of cloudy days 10 Number hours bright sunshine 171 Percent of possible hours of bright sunshine 57.8 Thunder and lightning 57.8	50.9 10 8 10 158	66, '90, '27; 31, 190, 19, 1941; 2, 192; 16, 1920; 2, 193; 16, 1894; 2, 192; 221, 1924; 110, '27, '38

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1944.

REMARKS

The weather during February was warmer than normal with a mean temperature of 25.7 degrees. The normal for February is 23.7 degrees. It was quite cold during the first week of the month, about normal for two weeks, and then warmer than normal during the last week. The mean temperature for the three winter months of December, January, and February was 22.9 degrees while the normal for that period is 25.1 degrees. The domestic heating load for the month was 1100 degree-days, making the total for the season 5170 degree-days. The normal for February is 1165 degree-days and for the season to the end of February the normal is 4982 degree-days. The heating load to the end of February is therefore about 4 percent higher than normal. The snowfall during the month was 16.50 inches compared to a normal of 13.99 inches. The snowfall for the winter is now 50.00 inches, the normal for the period being 38.15 inches. The total precipitation for the month was 3.33 inches. The normal for February is 3.19 inches.

There were 171 hours of bright sunshine, the normal being 158 hours. Total wind movement was 4277 miles while the normal wind travel for February is 4776 miles. The maximum wind velocity was 32 miles per hour on the 24th. The prevailing direction was west. Mean relative humidity was 67.5 percent, the normal being 66.6 percent.

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 675

March 1945

Meteorological Observations

FOR

MARCH

1945

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Te	mpera	ture		ty	s		Wind			Precipita	tion	
	Max	imum	Min	imum		Humidity	f Hour nshine			s)				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative F	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1	47	2 р	25	7 A 5 A	36.0	44	11.2	nw	111	13	12:30a	2:30 A	.05	.50
3	51 54 43	4 P 3 P 3 P	28 36 27	3 A 12 Р	39.5 45.0 35.0	60 83 41	4.5 2.0 11.4	e se w	104 124 208	9 22 28	4 A	8 A	.07	
2 3 4 5 6 7 8	44 44 42 43	4 P 4 P 3 P	15 34 26 18	8 A 2 A 12 P 7 A	29.5 39.0 34.0 30.5	65 95 56 48	11.4 1.0 11.5 11.6	n e nw	102 86 213 83	15 11 18 8	5 A 12:30A	11:30 p 1 A	.45 .01	
9 10 11	46 40 41	3 P 1 P	22 28 23	7 A 7 A 7 A	34.0 34.0 33.5	59 81 48	11.6 1.8 11.7	w nw nw	59 22 149	12 4 20	6 Р	7 Р	.04	
12 13 14	39 50 59	1 P 4 P 3 P	21 30 26	6 A 6 A 7 A	30.0 40.0 42.5	81 57 51	0.0 11.8 11.0	se W se	175 147 117	18 25 17	2:30 р	10-р	.32	1.00
15 16 17	71 51 68	4 P 4 P 3 P	35 37 36	7 A 7 A 7 A	53.0 44.0 52.0	78 91 79	8.9 0.0 6.2	se nw se	97 31 62 174	6 6 13 18			Т	
18 19 20 21	66 52 78 51	3 P 4 P 3 P 0 A	37 31 39 37	0 A 5 A 3 A 12 P	51.5 41.5 58.5 44.0	45 87 82 94	12.0 0.7 6.2 0.0	SW SW 11	43 83 127	18 17 18	7:30 p 3:30 a	8:30 P 10:30 P	.07 .61	
22 23 24	37 46 61	0 A 4 P 3 P	33 31 36	8 P 7 A 5 A	35.0 38.5 48.5	83 72 48	0.0 5.1 12.3	nw w nw	260 296 256	18 22 32	1 Р	6:30 р	.21	1.00
21 22 23 24 25 26 27 28 29 30	67 73 77	4 P 3 P 1 P	29 33 35	5 A 6 A 6 A	48.0 53.0 56.0	55 58 79	12.4 12.4 5.1	nw se se	98 187 86	14 14 28	3 р	11 :30 г	.24 T	
	84 85 68	3 P 3 P 3 P	44 56 52	7 A 7 A 10 Р	64.0 70.5 60.0	75 55 54	8.2 11.0 12.6	se w nw	99 176 304	17 14 22				
31	61	4 P	49	5 A	55.0	87	0.4	e	90	12	8 A	9 р	.09	

^{*}Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

March, 1945	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.37, 5th Minimum 29.45, 31st Mean semi-daily 30.001 Range 0.92	30.57 29.24 30.00 1.34	31.05, 1943 28.47, 1914 2.10, 1914; 0.85, 1915
Air Temperature, in degrees F.		, , , , , , , , , , , , , , , , , , , ,
Highest 85, 29th Lowest 15, 5th Mean 44.4 Range 70 Highest mean daily 70.5, 29th	63.8 6.1 34.4 57.6	79.5, 1907, 1921 -7.5, 1906
Lowest mean daily29.5, 5thMean maximum56.2Mean minimum32.6Greatest daily range42, 27thLeast daily range4, 22nd	43.2 25.4	
Precipitation, in inches		
Precipitation 2.16 Snow 2.50 Maximum precipitation in 24 hrs45, 6th	3.70 7.47	7.89, 1942; 0.12, 1915 27, 1899; 0, 1921
Number of days with .01 or more11	11	17, 1890, 1913, 1936; 3, 1915
Wind, in miles		
Total movement4169Greatest daily movement304, 30thLeast daily movement22, 10thMean hourly velocity5.6Maximum velocity32, 24th	5944 7.7 31.2	8182, 1896; 3006, 1905 48, 1932, 1939, 1941, 1942
Wind, direction		
Prevailing direction WNW North, days 2 Northeast, days 0 East, days 3 Southeast, days 8 South, days 0 Southwest, days 2 West, days 6 Northwest, days 10	WNW	
Weather		
Mean relative humidity, percent 67.5 Mean cloudiness, percent 53 Number of clear days 17 Number of fair days 5 Number of cloudy days 9 Number lours bright sunshine 226 Percent of possible hours of bright sunshine 60.9 Thunder and lightning 60.9	64.2 51.8 11 10 10 199 53.6	68, 1901; 27, 1915 22, 1924; 3, 1901 17, 1900; 1, 1943 21, 1901; 1, 1915 292, 1924; 93, 1901

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1944.

REMARKS

Weather during March was abnormally warm, with less rain and snow than usual. The mean temperature during the month was 44.4 degrees, while the normal mean temperature for March is 34.4 degrees. This is the highest mean temperature ever recorded for March at Amherst since 1837. The maximum temperature of 85 degrees on March 29 is the highest reading ever recorded in March. The heating load during the month was 645 degree-days, the normal being 950 degree-days. The heating load for the season is now 5815 degree-days whereas the normal for that period is 5932 degree-days.

A total of 2.50 inches of snow fell during the month and the total precipitation was 2.16 inches. Normally we get 7.47 inches of snow in March with a total precipitation of 3.70 inches. The total snowfall for the winter to the end of March is 52.50 inches, the normal being 47.78 inches. There were 226 hours of bright sunshine, the normal for March being 199 hours.

Dr. J. K. Shaw, research pomologist, gives the following report: "While the past winter was colder than the average, there were no severe minima, and peach fruit buds survived the winter in numbers sufficient for a good crop. The heavy covering of snow protected low-growing plants and the ground thawed from below so that there was little frost in the ground when the snow cover melted. This took place in March with unusual rapidity but with no high river floods. The extremely high temperatures in March have advanced vegetation so that it is about as far along as it was a month later last year. Grass is green and leaves are developing on many shrubs. Forsythia is in full bloom at the last of the month. It is, so far, the earliest spring in many years.

"We cannot expect this unseasonably warm weather to continue. We usually get temperatures in April low enough to cause severe injury to vegetation when it is advanced as far as it is at the end of March. While a light frost would not now do great damage, temperatures in the low twenties would cause great injury. Last year the low temperatures in April brought about a situation indicating little danger from spring frosts. Warm weather of early May advanced vegetation very rapidly, apple trees blossomed a day or two earlier than the average, and a disastrous frost occurred on the morning of May 19. This year conditions are reversed and damaging spring frosts are probable. We can only hope that luck will change this year and that we may have cooler weather in April, slowing the progress of vegetation, but without minima low enough to cause injury."

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 676

April 1945

Meteorological Observations

FOR

APRIL

1945

C. I. GUNNESS

OBSERVATORY

Latitude, $42^{\circ}\ 23'\ 31''\ N$. Longitude, $72^{\circ}\ 31'\ 48''\ W$. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

April, 1945	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.69, 7th Minimum 29.57, 27th Mean semi-daily 30.050 Range 1.12	30.48 29.38 29.99	30.71, 1911 28.98, 1943
	1.11	1.52, 1930; .72, 1919
Air Temperature, in degrees F.	79. 1	90, 1941
Highest 86, 13th Lowest 26, 23rd Mean 51.8 Range 60 Highest mean daily 70, 13th Lowest mean daily 41, 5th, 6th, 23rd Mean maximum 63.6 Mean minimum 40.0 Greatest daily range 44, 10th Least daily range 9, 2nd, 4th	22.0 45.7 57.4	90, 1941 8.5, 1923 52, 1921; 41, 1943
Mean maximum63.6Mean minimum40.0Greatest daily range44, 10thLeast daily range9, 2nd, 4th	56.8 34.8	
Precipitation, in inches		
Precipitation 5.43 Snow 5.283, 25th Number of days with .01 or more 9	2.17	6.89, 1929; .55, 1941 11, 1891; 0, 1910, 1934
	11	18, 1909; 3, 1892
Wind, in miles		0200 1000 2052 1017
Total movement5369Greatest daily movement484, 22ndLeast daily movement70, 30thMean hourly velocity7.5Maximum velocity40, 5th	7.5 31.4	8208, 1908; 3853, 1917 40, 1935, 1938, 1940
Wind, direction		
Prevailing direction SW North, days 0 Northeast, days 0 East, days 1 Southeast, days 13 South, days 0 Southwest, days 0 West, days 8 Northwest, days 8 Routhwest, days 8	WNW	
Weather		
Mean relative humidity, percent 67.6 Mean cloudiness, percent 56 Number of clear days 16 Number of fair days 8 Number of cloudy days 6 Number hours bright sunshine 258 Percent of possible hours of bright sunshine 64.2 Thunder and lightning	61.6 51.8 11 9 10 220 54.7	75, 1901; 34, 1925, 1927 23, 1941; 3, 1898, 1902 18, 1915; 2, 1894, 1901 22, 1901; 1, 1941 329, 1941; 103, 1901
Last snow . March 22	Apr. 15	Mar. 14, 1910; May 11, 1907

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1944.

		Te	mpera	ture		ty	Yind Wind				Precipita	tion		
	Max	imum	Min	imum		Humidi	r of Hours Sunshine	20		sə .				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number o Bright Su	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 22 23 24 25 26 27 28 29 30 30 30 30 30 30 30 30 30 30 30 30 30	71 61 69 48 49 51 71 76 81 80 82 86 75 57 53 67 52 63 48 56 62 55 68 57 67 67 67 67 67 67 67 67 67 67 67 67 67	4 P P 6 P P 3 P P 4 P P 1 P	45 52 48 39 33 31 27 36 35 37 47 54 49 36 34 42 43 35 34 42 43 40 45	5 A A 12 P 12 P 12 P 12 P 12 P 12 P 12 P	58.0 56.5 58.5 58.5 58.5 41.0 41.0 43.5 59.0 63.5 69.0 62.0 46.5 54.5 55.5 54.5 54.5 54.5 55.5 54.5 54.5 54.5 55.5 54.5 55.5 54.5 69.0 62.0 46.5 54.5 55.5 55.5 55.5 56.5	53 84 73 87 77 44 57 66 65 79 78 80 55 46 64 70 54 42 61 93 76 63 76 81	12.7 0.0 11.7 0.0 4.5 13.0 13.0 13.1 13.1 8.5 8.0 9.0 10.7 13.3 1.9 0.0 12.6 13.5 7.0 10.0 9.9 9.0 13.7 6.9 0.0 12.3 10.0 13.7	w se se se se se w nw se se nw nw se nw nw w w w w w w w w w w w w w w w w	313 189 211 197 272 366 187 166 80 82 92 118 105 222 154 113 132 275 185 79 484 119 72 83 224 119 70 70	28 15 28 14 40 20 28 13 7 11 10 12 14 25 11 11 17 32 20 17 32 13 9 9 17 28 28 13 13 14 20 15 16 17 17 18 18 18 18 18 18 18 18 18 18 18 18 18	9 A 8:30 P 0 A 4:30 P 12:30 P 9 P 0 A 5:30 A	12 P 12 P 12 M 12 M 7:30 I 11:30 P 12 P 7 A	.41 .06 1.13 .50 .26 2.83 .05 .06	

^{*}Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

The warm weather which started in March continued through the first half of April but the latter half of the month has been near normal. The mean temperature for the month, however, was 51.8 degrees which is 6.8 degrees higher than normal. The high mean temperature of 51.8 degrees has been exceeded only once in April since 1889; that was in 1941 when the mean was 52.1 degrees. The highest temperature was 86 degrees on the 13th. The lowest temperature was 26 degrees on the 23rd, causing general damage to the fruit crop.

The domestic heating load was 406 degree-days compared to a normal of 579 degree-days. The heating load for the season to the end of April is 6221 degree-days, the normal for that period

being 6511 degree-days.

The rainfall during the month was 5.43 inches with no snow. The normal precipitation for April is 3.35 inches with 2.17 inches of snow. There were 258 hours of bright sunshine which is 38 hours more than normal.

Dr. J. K. Shaw makes the following report on the progress of vegetation: "The fear of cold injury expressed last month proved well founded. Temperatures appropriate to March came along in April and there has been severe damage to fruit crops. Apple bloom was not as heavy as expected but there was enough for a good crop. We had a narrow escape on April 6 and 7 when temperatures approached the danger point but caused little if any injury. Then on the morning of April 23 there was severe and widespread damage. There was less difference in orchards on different elevations than there was from the freeze of May 19, 1944. Injury was more general and more severe. From 80 to 90 percent of the flowers on all tree fruits in our orchards were killed. This seems to include all that were open at the time. Some of the unopened buds were killed and others survive. Ordinarily such buds do not produce much good fruit but being relieved of the competition from the earlier opening flowers they may mature good fruits. The continued low temperatures that prevailed through April are unfavorable to bee flight and to fertilization of the blossoms. From one to three weeks must pass before we can tell how many of these flowers will produce fruits. The little embryonic peaches have not swelled following bloom as they usually do. This may be only a delay due to continued cold weather or they may be ruined. In some parts of this state more injury is reported to have occurred on April 24 but in Amherst the temperature was about 4° higher and little if any additional injury took place. Apparently in Amherst orchards as much or more than in any other section of the state there is still a danger period of 2 or 3 weeks ahead of us.

"Open strawberry flowers were killed but these were too few to materially affect the crop. No distinct injury to bush fruits and

grapes has yet been determined.

"The apple blossom period was the earliest and most prolonged of any in the past 40 or more years and fully three weeks ahead of normal. Last year the blooming period of all tree fruits lasted scarcely over a week; this year it has continued for nearly a month and some apple blossoms had not opened at the end of April.

"Two successive years of cold damage to fruits has never before occurred in Massachusetts within the memory of the present generation. There may be a good many bushels of apples and other tree fruits in Massachusetts in 1945 but the crop must be reduced 50 percent and probably more—perhaps much more."

BLACKSBURG, VIRGINIA

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 677

May 1945

Meteorological Observations

FOR

MAY

1945

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Тe	mpera	iture		t y	90		Wind			Precipita	ition		
	Maximum		Minimum			lumidi	r of Hours Sunshine				so U				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number o Bright Su	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow	
1 2 3 4 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 22 23 31 24 25 27 28 29 30 31 31 31 31 31 31 31 31 31 31 31 31 31	55 66 61 77 80 64 66 70 58 71 71 80	4 P P 1 P P P P P P P P P P P P P P P P	43 38 42 46 39 40 41 39 36 31 39 48 41 51 55 45 36 46 40 37 41 41 41 41 41 41 41 41 41 41	12 P 5 A A 0 A A 5 A A 5 A A 12 P 1 A A 3 A 12 P 5 A A 12 P 5 A A 11 P 12 P 5 A 12 P 5 A 12 P 5 A 12 P 13 A 12 P 12 P 13 A 12 P 14 A 15 A 16 A 17 A 18 A 18 A 18 A 18 A 18 A 18 A 18 A 18	47.5 49.0 51.0 54.5 50.0 54.5 50.5 41.0 43.0 52.5 57.5 59.0 65.5 57.5 58.5 65.0 54.5 52.5 54.5 54.5 54.5 54.5 55.5 54.5 55.5 55.5 55.5 55.5 56.0 56.0 56.0 56	96 77 83 97 72 79 66 74 49 86 57 85 97 92 77 85 61 64 64 59 89 89 88 89 88 89 88 88 88 88 88 88 88	0.0 6.3 0.0 7.9 8.8 14.2 2.2 13.5 0.0 10.1 1.5 11.8 10.7 4.6 1.1 2.8 0.0 0.0 3.7 6.8 6.5 12.5 7.5 13.5 0.0 0.0 8.8	e se e nw se w w se nw n se nw w se nw w se nw m se nw w se nw w se nw w w se se se nw w w w w w w w w w w w w w w w w w w	97 66 133 162 152 120 47 88 206 79 341 89 62 88 175 86 42 92 145 125 92 145 124 358 167 65 71 57 140 215 301 391	15 9 18 14 18 15 8 18 22 15 32 10 20 12 18 12 6 10 18 17 12 17 12 18 8 7 8 7 8 7 17 17 17 17 17 17 17 17 17 17 17 17 1	1 :30 A 1 :30 A 2 P 1 P 11 :30 A 0 A 1 P 8 P	11:30 A 5 P 12 P 9 P 3:30 P 12 P 6 A 6 P 12:30 A 12 P 12:30 A 12 P 1 A 12:30 P 1 A	.25 .04 .50 .58 .04 .25 1.07 .46 .54 .61 .10 .14 1.04 .35 .04	.75 .25	

^{*}Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

May 1945	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum30.35, 27th	30.39	30.62, 1936
Minimum 29.30, 10th	29.51	29.10, 1938
Mean semi-daily	29.96	27113, 1733
Range1.05	0.88	
Air Temperature, in degrees F.		
Highest 81, 26th	86:1	94.5, 1896, 1911
Lowest 31, 11th	31.2	24.0, 1906
Mean	57.1	
Range 50	54.9	
Highest mean daily 65.5, 15th		
Lowest mean daily 41, 10th Mean maximum 64.5		
Mean minimum 43.7		
Greatest daily range 40, 26th		
Least daily range 6, 4th		
Precipitation, in inches		
Precipitation6.45	3.60	7.44, 1931; .48, 1903
Snow 1.00	3.00	
Maximum precipitation in 24 hrs.		
1.53, 10th, 11th		
Number of days with .01 or more20	12	20, 1901; 5, 1903
Wind, in miles		
Total movement 4379	4504	5946, 1907; 2180, 189-
Greatest daily movement 391, 31st		
Least daily movement 42, 17th		
Mean hourly velocity5.9 Maximum velocity35, 31st	6.1 28.8	45, 1933
· ·	28.8	75, 1955
Wind, direction	117	
Prevailing direction SSW	117	
North, days 1 Northeast, days 0		
Northeast, days		
East, days 2 Southeast, days 12		
South, days 1		
Southwest, days0		
West days 8		
West, days 8 Northwest, days 7		
Weather		
Mean relative humidity, percent73.2	60.7	50 100 100 20 20 20
Mean cloudiness, percent 65	52.1	70, 1901, 1902; 30, 1923
Number of clear days 8	11	21, 1944; 0, 1927 17, 1907, 1926; 5, 1933
Number of fair days = 11	11	20, 1927; 2, 194
Number of cloudy days 12	252	350, 1944; 137, 1927
Number of cloudy days Number hours bright sunshine Percent of possible hours of bright sunshine Thunder and lightning Last snow 12 190 41.9 42.9 41.9 41.9 41.9 41.9	200	000, 1277, 107, 1927
conchine 41.0	55.6	
Thurston and lightning 22nd		
I minder and nething		

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1944

REMARKS

The weather during May was cold and wet, delaying planting and in general producing a late spring in spite of the warm weather in late March and early April. The mean temperature for May was 54.1 degrees, the normal mean being 57.1 degrees. This is the coldest May since 1927 and in only five years has there been a colder May since this station was established in 1889. The lowest temperature was 31 degrees on the 11th. This was the last frost at the observatory although there were frosts on low ground in Amherst on the 25th. The heating load for May was 338 degree-days, the normal being 267 degree-days. The total heating load for the season is now 6559 degree-days, the normal to the end of May being 6778 degree-days.

The rainfall for the month was 6.45 inches compared to a normal of 3.60 inches. Rain fell on 20 days during the month. This heavy rainfall in May has been exceeded only twice since 1889, the last being in 1931 when 7.44 inches fell. An inch of snow fell on the 10th and 11th. There were 190 hours of bright sunshine,

the normal for May being 252 hours.

Dr. J. K. Shaw reports on the progress of vegetation as follows: "The weather during May continued cool with an excess of rainfall. These conditions assure a good hay crop but were not very favorable for fruit. Yet the blossoms not killed by the April freeze set rather well so that a small and spotty crop is in sight. Prospects are better than was feared a month ago. Plums are very light but pears and peaches seem to have set good crops. The apple crop over the state is variable, some orchards having a good prospect of a crop and in others a total failure is indicated. Of course, many orchards are between these extremes. The common estimate of half a crop seems reasonable. Frequent rainy periods have favored apple scab infection so that orchards not thoroughly sprayed may produce many scabby apples. Those well sprayed at proper times are quite free from scab. While frost killed the very early strawberry blossoms, conditions so far have favored a good crop from the limited plantings made last year. The soil is now well supplied with moisture, and if dry, sunny weather prevails during the ripening period, a good crop should be harvested. The month closes with a cold night, reminding us that it is still possible to have further frost injury."

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 678

June 1945

Meteorological Observations

FOR

JUNE

1945

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

JUNE, 1945	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.16, 15th Minimum 29.64, 3rd Mean semi-daily 29.942 Range 0.52	30.30 29.55 29.95 0.75	30.54, 1889 29.24, 1902
Air Temperature, in degrees F.		
Highest 92, 15th, 30th Lowest 37, 1st, 7th Mean 66.1 Range 55 Highest mean daily 80, 30th Lowest mean daily 50, 2nd Mean maximum 77.3 Mean minimum 54.8 Greatest daily range 33, 7th Least daily range 7, 19th	91.2 40.0 65.7 51.3	101, 1919 34, 1891
Precipitation, in inches		
Precipitation	3.75	9.68, 1922; 0.76, 1908
Number of days with .01 or more17	11	17, 1922; 4, 1908
Wind, in miles Total movement 2805 Greatest daily movement 30, 13th Least daily movement 30, 13th Mean hourly velocity 3.9 Maximum velocity 32, 15th	3585	4571, 1908; 1409, 1906
Mean hourly velocity 3.9 Maximum velocity 32, 15th	5.0 24.6	48, 1939
Wind, direction		
Prevailing direction NNW North, days 5 Northeast, days 0 East, days 4 Southeast, days 6 South, days 2 Southwest, days 1 West, days 3 Northwest, days 9	WSW	
Weather		
Mean relative humidity, percent Mean cloudiness, percent 58 Number of clear days 8 Number of fair days 15 Number of cloudy days 7 Number hours bright sunshine 242 Percent of possible hours of bright sunshine 53 Thunder and lightning	66.9 51.1 10 12 8 257 54.0	71, 1903; 28 1908 22, 1908, 1941, 1943; 1, 1927 23, 1912; 3, 1941 22, 1903; 1, 1923 362, 1908; 102, 1903
Last frost And Ighthing 17th, 18th, 25th, 26th April 24		

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1944.

		Те	mpera	ture		ty	s	Wind			Precipitation			
	Max	imum	Minimum			Iumidi	r of Hours Sunshine			99				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Bright Sur	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 3 4 4 5 6 7 8 9 10 112 13 114 115 116 117 118 119 20 21 22 23 22 4 25 26 27 28 9 30	64 61 57 62 57 65 70 72 78 81 81 78 90 92 87 88 88 88 72 75 75 79 82 84 84 89 90 92 89 92	5 P 12 M 1 2 P 1 3 P P 1 3 P P 1 2 P P 1 3 P P 1 1 A P P 1 3 5 P P P 1 1 A P P 1 5 5 P P P 1 A P P 1 2 P P 1 A P P 1 2 P P 1 A P 1 A P P 1 A P 1	37 39 46 44 45 37 43 45 55 58 52 60 66 54 62 58 59 61 63 68	5 A A A A A A A A A A A A A A A A A A A	50.5 50.0 51.5 54.0 50.5 53.5 57.5 62.0 70.0 69.5 77.5 68.5 77.5 68.5 71.0 69.0 76.0 69.0 76.0 69.0 76.0 69.0 76.0 69.0 76.0	46 71 89 64 88 67 58 64 64 88 98 67 76 85 85 85 85 85 64 72 88 67 72 66	10.0 3.0 3.1 4.3 5.0 9.0 9.3 12.6 7.2 1.0 6.6 13.0 9.0 10.6 6.0 7.2 8.5 5.0 2.1 14.0 9.2 12.0 9.0 11.6 11.6 12.6 12.6 12.6 12.6 13.0 14.3 14.3 14.3 14.3 14.3 14.3 14.3 14.3	w n n n e n nw nw se e se se se e sw se nw nw s nw nw se nw nw s	241 41 68 71 110 86 77 40 79 113 89 30 54 106 37 93 91 181 181 55 87 141 279 45 59	22 5 10 11 12 20 17 22 6 9 18 13 5 7 32 11 12 20 17 22 6 9 18 13 11 12 20 17 22 4 4 10 11 11 11 11 11 11 11 11 11 11 11 11	3:30 P 6 A	5 P 8 A A 10:30 P 2 A 12 P 6 A A 10 P 9 P 12 P 8 A A 10 P 6 :30 P 4 P	.06 .08 .22 .05 .21 .68 .01 .01 1.85 .13 2.25 .22 .09 .73 .58	

^{*}Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

REMARKS

The outstanding feature of June weather was the excessive rainfall. Rain fell on 17 days with a total of 7.67 inches. This rainfall for June has been exceeded only three times since the station was established in 1889: In June 1903 with a rainfall of 7.79 inches, in 1922 with 9.68 inches, and in 1938 with 8.45 inches. Only once was this June rainfall exceeded during the period from 1836 to 1888. In 1862 the June rainfall in Amherst was 11.69 inches. The normal rainfall for June is 3.75 inches. The normal total rainfall for April, May and June is 10.70 inches. This year the rainfall for these three months is 19.55 inches. This has never been reached since records have been taken since 1836. Normally rain falls on 34 days during the three spring months. This year rain fell on 46 days.

The mean temperature for the month was 66.1 degrees, the normal being 65.7 degrees. The highest temperature was 92 degrees on the 15th and 30th. The lowest temperature was 37 degrees on the 1st and 7th. The heating load during the month was 105 degree-days compared to a normal of 68 degree-days. This brings the total heating load for the season to 6664 degree-days, the normal being 6846 degree-days. The heating load for the season was therefore slightly less than 3 percent below normal. There were 242 hours of bright sunshine compared to a normal of 257 hours.

Dr. J. K. Shaw makes the following statement: "Excessive rainfall continued through June but with a change to hot, humid weather. The hay crop is heavy but frequent rains have delayed harvest and injured quality. Peaches and pears have set a good crop, but a severe local hailstorm on the afternoon of June 15 caused injury and broke a large percentage of our greenhouse glass, although there was little or no hail damage a mile or two away. The strawberry crop was good but suffered from rot because of wet weather. The raspberry crop will be good but blueberries will be short. Some varieties of apples set well from late bloom in spite of April freezes but, in our own orchards, suffered from hail damage. McIntosh, our main variety, failed to set well from the uninjured blossoms and the crop will be very small. Few of our mature trees will yield a bushel each. The apple crop in this vicinity will be the smallest in many years and of poor quality because of scab, frost russeting, and, in our own orchards, hail damage. Last month's estimate of half a crop was over optimistic, largely because of a poor set of McIntosh. Present indications suggest a quarter of a crop is a more reasonable expectation."

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 679

July, 1945

Meteorological Observations

FOR

JULY

1945

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Te	трега	ture		ty	•	Wind			Precipitation			
	Max	imum	Minimum			Humidity	Hour nshine			*				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative F	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7	92 82 77 79 84 83	2 P 1 A 4 P 2 P 3 P 1 P	63 60 53 50 60 58	5 A 12 P 12 P 5 A 5 A 12 P	77.5 71.0 65.0 64.5 72.0 70.5	73 80 62 65 69 66	12.3 5.0 15.2 14.2 11.7 9.0	se s nw se se nw	76 47 168 87 146 145	28 13 20 11 13 25	3:30 P 4 P	4 P 9:30 P	.58 .26	
8 9 10 11 12 13	81 83 86 79 69 76 77	4 P 3 P 2 P 1 P 5 P 4 P	52 55 57 56 51 46 56	4 A 2 A 3 A 12 P 3 A 5 A	66.5 69.0 71.5 67.5 60.0 61.0 66.5	61 61 76 84 49 53 67	15.0 14.0 9.0 5.2 14.0 13.5 10.0	w se s se w se se	90 74 112 200 222 101 162	18 12 13 17 22 10 14	8:30 P 2 P	9:30 P 3:30 P	.04 1.24	
14 15 16 17 18	82 76 79 75 83 80	2 P 11 A 5 P 1 P	61 68 64 61 66 70	7 A 12 P 12 P 5 A 6 A	71.5 72.0 71.5 68.0 74.5 75.0	76 89 81 77 82 86	9.5 0.0 8.0 4.3 8.0	se se nw nw se	116 227 94 56 75 93	12 18 6 6 8 9	11:30 P 0 A 9:30 P 1:30 A 9 A	12 P 7 P 11:30 P 6 A 6 P	.13 1.72 .06 .69 .11	
19 20 21 22 23 24	80 85 78 81 79	11 A 5 P 2 P 11 A 4 P 5 P	69 69 66 64 64	0 A 4 A 12 P 3 A 5 A	74.5 77.0 72.0 72.5 71.5	86 69 76 72 82	3.3 6.0 10.0 2.0 10.0 9.0	se nw e se e	115 121 71 47 59	12 10 13 6 5	4:30 P 4 A	12 M 6:30 P 6 A	.03 .27 .01	
21 22 23 24 25 26 27 28 29 30	87 88 81 77 77 84 81	2 P 1 P 3 P 2 P 5 P 3 P 4 P	65 68 57 53 65 66 63	2 A 12 P 12 P 4 A 8 A 6 A 12 P	76.0 78.0 69.0 65.0 71.0 75.0 72.0	79 73 57 75 89 76 76	7.0 9.2 10.0 7.0 3.8 10.9 8.9	se nw e e nw se e	108 95 123 46 64 53 77	9 10 10 5 5 6 7	6 A 6:30 P 0 A	6:30 A 12 P 12 P	.04 .91 .93	

^{*}Based on least time required to blow one mile.

FERNAND E. BARTLETT. Observer

July, 1945	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.37, 17th Minimum 29.67, 10th Mean semi-daily 30.024 Range 0.70	30.27 29.59 29.96 0.68	30.50, 1892 29.27, 1932 0.97, 1892; 0.47, 1938
Air Temperature, in degrees F.	0.00	0.57, 1052, 0.47, 1938
Highest 92, 1st Lowest 46, 12th Mean 70.6 Range 46 Highest mean daily 78, 26th Lowest mean daily 60, 11th Mean maximum 80.7 Mean minimum 60.5 Greatest daily range 30, 12th Least daily range 8, 15th	93.9 46.4 70.8 47.7	104, 1911 40, 1890, 1898 74.7, 1921; 66.3, 1891
Precipitation, in inches		
Precipitation 7.36 Snow 7.36 Maximum precipitation in 24 hrs.	4.10	14.51, 1897; 0.70, 1929
Number of days with .01 or more16	11	20, 1915; 4, 1924
Wind, in miles		
Total movement 3270 Greatest daily movement 227, 15th Least daily movement 46, 28th Mean hourly velocity 4.4	3422 4.6	5097, 1909; 1109, 1894
Maximum velocity 28, 1st	28.7	44, 1936
Wind, direction Prevailing direction SE North, days 0 Northeast, days 5 Southeast, days 15 South, days 2 Southwest days 0 West, days 2 Northwest, days 7	SW	
Weather		
Mean relative humidity, percent73.1Mean cloudiness, percent60Number of clear days9Number of fair days17	68.4 50.9 10 14	70, 1943; 31, 1924 22, 1923; 0, 1915 24, '09, '21; 5, '89, '23, '28.
Number of cloudy days 5 Number hours bright sunshine 275 Percent of possible hours of bright sunshine 59.5 Thunder and lightning 1st, 6th, 10th, 18th	7 268 58	'36 18, 1889; 0, 1910 371, 1910; 180, 1931

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938. except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1944.

REMARKS

The outstanding feature of July weather was the continuation of the heavy rainfall which started in April. Rain fell on 16 days in July and the total was 7.36 inches as compared to a normal of 4.10 inches. This heavy rainfall has been exceeded only three times in July since 1889: 10.52 inches fell in 1889; 14.51 in 1897, and 9.13 in 1915. Much of the rain occurred as heavy showers. The total rainfall since April 1 is now 26.91 inches, the normal for this period being 14.80 inches. The heavy rainfall has been disastrous for many crops and has made haying particularly difficult.

The mean temperature for the month was 70.6 degrees which is only 0.2 of a degree less than normal. In spite of the frequent rains, we had 275 hours of bright sunshine while the normal for July is 268 hours. There were no high winds and the average humidity was 73.1 percent.

Professor J. S. Bailey of the Department of Pomology makes the following comments: "July has been another month of hot, humid weather and excessive rainfall. A prospective good raspberry crop turned out to be very light because so many berries became too soft or molded on the bushes. The blueberry crop, both cultivated and wild, has been cut heavily by mummy berry disease. Apple scab has been the worst in years. McIntosh orchards that were not frequently and thoroughly sprayed are heavily infected. Scab carryover for next year will be heavy.

"Light crops and abundant water have resulted in rapid sizing of fruit with prospects of many oversized apples of poor keeping quality. Several very heavy rains have increased the difficulty of insect control by washing off arsenical sprays. Severe erosion during the heavy rains re-emphasizes the need for erosion control in many places. Late blight of potatoes appeared early and is very severe in many Connecticut Valley fields. Many other diseases of both fruits and vegetables have been unusually severe."

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 680

August, 1945

Meteorological Observations

FOR

AUGUST

1945

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Te	mpera	iture		ty			Wind			Precipitation			
	Maximum		Minimum			Humidity	r of Hours Sunshine			80 60					
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative F	Number of Bright Su	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow	
1 2 3 3 4 5 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 1 22 23 24 25 6 27 28 29 30 31	75 81 86 76 78 62 67 80 80 84 88 87 85 84 81 77 91 82 84 85 75 69 61 70 70 74 84 86 86 86 86 86 86 87 88 88 88 88 88 88 88 88 88 88 88 88	5 P 1 P 12 M 4 P 12 M 2 P 1 2 M 1 P 1 2 P 1 A P	60 64 55 55 50 52 58 57 57 53 56 66 67 57 47 50 52 54 53 61 62 62	6 5 A A A A A A A A A A A A A A A A A A	67.5 72.5 74.0 65.5 64.0 62.5 68.5 72.0 75.5 69.0 62.5 68.5 72.0 62.0 64.5 67.0 69.0 76.0 60.5 560.0 63.5 75.5 75.5 73.0 75.5 75.5 75.5	79 71 57 60 91 87 74 70 78 71 62 67 71 64 69 75 64 66 95 97 77 62 88 84	6.7 8.5 13.0 9.2 13.0 0.8 8.7 9.7 6.1 14.0 9.0 10.1 7.8 5.0 12.9 10.0 13.8 11.5 11.0 10.7 12.2 7.0 0.0 5.0 12.9 12.0 8.9 11.0 6.5	se se nw w w se se nw nw nw w w se se se nw nw nw nw nw w se s	67 81 170 159 139 51 146 48 37 40 36 120 157 213 58 64 60 59 51 100 101 53 88 121 125 90 97 53	6 9 18 20 17 13 13 10 10 5 3 7 7 11 20 7 8 5 6 6 5 11 11 28 7 8 18 18 18 18 18 18 18 18 18 18 18 18 1	10 A 12:30P 6:30 A 0 A 6:30 A	12 P 1 P 8:30 P 7:30 A 9 P	1.05 .25 .91 .09		

^{*}Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

August, 1945	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.27, 24th Minimum 29.66, 26th Mean semi-daily 30.004 Range 0.61	30.32 29.61 30.00 0.71	30.50, 1934 28.87, 1930
Air Temperature, in degrees F.		
Highest 91, 21st Lowest 47, 16th Mean 68.3 Range 44 Highest mean daily 76, 21st Lowest mean daily 56, 25th Mean maximum 79.8 Mean minimum 56.9 Greatest daily range 34, 20th Least daily range 9, 7th	91.6 43.4 68.6 48.1	100, 1918 37, 1894, 1908 34, 1940
Precipitation, in inches		
Precipitation 2.79 Snow — —	4.08	8.40, 1928; .31, 1894
Maximum precipitation in 24 hrs1.05, 6th Number of days with .01 or more6	11	16, 1892, 1933; 4, 1899
Wind, in miles		
Total movement2777Greatest daily movement213, 15thLeast daily movement36, 12thMean hourly velocity3.7Maximum velocity28, 23rd	3127 4.2 22.7	4,271, 1910; 1,920, 189- 40, 1941
Wind, direction	23.7	
Prevailing direction SW North, days 0 Northeast, days 0 East, days 1 Southeast, days 9 South, days 4 Sonthwest, days 2 West, days 7 Northwest, days 8	ł	
Weather		
Mean relative humidity, percent 72.6 Mean cloudiness, percent 50 Number of clear days 13 Number of fair days 14 Number of cloudy days 4 Number hours bright sunshine 263 Percent of possible hours of bright sunshine 61.3 Thunder and lightning 30th, 31st	49.7 9 13 9 237	67, 1901; 27, 192; 23, 1941; 0, 1915, 1926 25, 1912; 3, 1936 18, 1901, '28; 2, 1910, '2; 332, 1941; 152, 1915, 1926

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1944.

The weather during August was near normal in most respects. The mean temperature was 58.3 degrees, the normal being 68.8 degrees. The rainfall was less than normal. The total rainfall was 2.79 inches, the normal being 4.08 inches. There were 263 hours of bright sunshine as compared to a normal of 237 hours. The mean relative humidity was 72.6 percent, the normal being 70.2 percent. There were no high winds.

The following report is given by Dr. J. K. Shaw: "The excessive rainfall which has prevailed since spring came to an end with August, but the soil is still well supplied with water and we can be quite sure that the summer of 1945 will be one with no injury from dry weather. Hot, humid periods interspersed with cool weather prevailed during the month. The weather has been bad for potatoes for the vines grew well during the wet weather and were unable to stand the hot, sunny weather which followed. Consequently, we may expect a poor crop in this vicinity.

"The apple crop will be the smallest in many years, and of very poor quality in the College orchards because of injury by hail. Apple scab is more prevalent than usual because of 'scab weather' and the fact that many growers did not spray as thoroughly as usual on account of the lack of a crop. The peach crop is good both here and over the whole country. Pears will make a fair crop but plums are few.

Plenty of rain insured a good second crop of hay, which should be plentiful next winter although much of it will be of poor quality owing to wet weather which delayed harvest."

Dr. A. B. Beaumont makes the following comments on clovers: "This has been one of the best 'clover' years observed by local agronomists. Tame and wild clovers have grown luxuriantly. Red and alsike clovers have been abundant in new hay seedings, and red clover has reappeared in many old mowings. The rowen crop now being harvested contains a high proportion of clovers. Wild white clover and the giant white cultivated clover (Ladino) have grown especially well in fertilized pastures. Sweet clover and rabbit-foot clover have been much in evidence along roadsides and in waste places. Abundant rains during the early part of the season are thought to have been responsible for the abundance of clovers this year."

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series Bulletin No. 681

September 1945

Meteorological Observations

FOR

SEPTEMBER

1945

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Те	mpera	iture		ty	s		Wind			Precipit	ation	
	Max	imum	Min	imum		Humidity	f Hour nshine	ba		80_				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative I	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30	85 73 75 76 83 87 91 85 81 77 77 66 63 56 61 74 67 67 67 67 67 67 67 67 67 67 67 67 67	2 P 12 M 3 P 4 P P 3 P P 1 P P	666 582 455 517 600 588 633 668 560 57 611 47 47 47 47 47 47 47 47 47 47 47 47 47	3 A 12 P 12 P 6 A 7 A 6 A 3 A 2 A 12 P 12 P 12 P 4 A 3 A 12 P 4 A 12 P 4 A 12 P 5 A 12 P 7 A 12 P 8 A 12 P 8 A 12 P 8 A 13 A 14 A 15 A 16 A 17 A 18 A 18 A 18 A 18 A 18 A 18 A 18 A 18	75.5 63.5 60.5 67.0 72.0 73.5 74.5 74.5 63.5 61.5 54.5 54.5 54.5 54.5 54.5 54.5 63.5 74.0 62.5 54.5 63.5 64.5 64.5 64.5 64.5 64.5 64.5 64.5 64	80 83 63 71 72 73 72 90 87 66 73 95 82 67 82 76 81 81 86 84 78 76 93 81 64	5.5 2.0 12.0 10.6 9.0 10.8 9.3 6.3 4.0 0.7 10.0 10.2 0.1 0.0 6.5 0.0 9.4 10.7 3.0 0.0 6.7 0.0 6.7	se n nw se e se sw sw ne n n s nw nw n se se se sw nw nw n n se w n n se se nw w se nw nw nw nw nw se nw nw nw se nw nw se nw nw se nw nw nw se nw nw nw nw nw nw nw nw nw nw nw nw nw	162 125 66 34 653 663 664 7100 666 74 98 50 32 121 142 105 136 1105 88 178 136 41 117 1158 140 43 109 221 185	18 14 10 12 6 5 6 6 13 9 12 12 8 5 14 18 7 12 14 9 28 17 6 9 12 11 5 11 18 20	2 A 3 A 12:30 A	7 P 4:30 P 9:30 A 12 P 4 A	.19 .16 .20 .39 1.97 .19 .37	

^{*}Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

September, 1945	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.61, 30th Mininum 29.69, 11th Mean semi-daily 30.156	30.45 29.57 30.06	30.65, 1924 28.41, 1938
Range	.88	1.99, 1938; .57, 1910
Air Temperature, in degrees F.		
Highest 91, 7th, 8th Lowest 36, 30th Mean 64.6 Range 55 Highest mean daily 75.5, 1st, 7th Lowest mean daily 46.5, 30th Mean maximum 74.5 Mean minimum 54.6 Greatest daily range 33, 8th Least daily range 8, 11th	87.7 33.2 61.7 54.5	97, 1895, 1929 24.5, 1914
Precipitation, in inches		
Precipitation 3.57	4.24	14.55, 1938; .52, 1914
Maximum precipitation in 24 hrs. 1.97, 15th Number of days with .01 or more	10	16, 1933; 3, 1903
Wind, in miles		
Total movement 3118 Greatest daily movement 221, 29th Least daily movement 32, 14th	3271	4,686, 1896; 1,414, 1894
Mean hourly velocity 4.3 Maximum velocity 28, 21st	4.5 25.9	. 80, 1938
Wind, direction		
Prevailing direction WNW North, days 5 Northeast, days 1 East, days 2 Southeast, days 7 South, days 2 Southwest, days 3 West, days 3 Northwest, days 7	WSW	
Weather		
Mean relative humidity, percent 76.5 Mean cloudiness, percent 54 Number of clear days 10 Number of fair days 10 Number of cloudy days 10 Number hours bright sunshine 174 Percent of possible hours of bright sunshine 46.6	73.6 49.9 10 10 10 201 54.9	70, 1934; 27, 1908, 1914 19, 1932; 2, 1907, 1928 19, 1908; 3, 1889 20, 1934; 3, 1941 255, 1916; 106, 1934
Thunder and lightning — — First frost — —	Sept. 21	Oct. 13; Aug. 22

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1944.

The weather during the month of September was very pleasant, with higher temperatures than normal and slightly less rainfall. The total precipitation for the month was 3.57 inches, while the normal for September is 4.24 inches. The mean temperature was 64.6 degrees, the normal for September being 61.7 degrees. The highest temperature was 91 degrees on the 7th and 8th and the lowest, 36 degrees on the 30th. There has, therefore, been no frost in September this year. The normal date for the first frost is September 21st. The domestic heating load for the month was 110 degree-days compared to a normal of 149 degree-days. There were 174 hours of bright sunshine compared to a normal of 201 hours. There were no very high winds during the month, the highest being 28 miles per hour on the 21st.

Dr. J. K. Shaw makes the following comments: "Cool weather interspersed with hot, humid periods characterized the weather of September. The McIntosh apple harvest was completed. Not only was the crop the smallest in many years, but the quality was poor owing to unusual prevalence of scab in some orchards. Other later varieties are partially harvested. A few, mostly the later blooming varieties, will yield good crops, but the total crop in Massachusetts will not greatly exceed one-fourth of a normal crop. It will be less than that in this immediate vicinity. Probably the short McIntosh crop is due as much to cold, wet weather following bloom as to the spring frost during the remarkably early blooming period. These conditions interfered with adequate fertilization of the flowers. Enough McIntosh blossoms survived the frost to make a fair crop in many orchards if fertilization had been effective. Other tree fruits matured fair to good crops. The potato crop in this vicinity will be poor owing to the wet spring and unfavorable weather in August. The onion crop in the Valley is considerably below average."

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 682

October 1945

Meteorological Observations

FOR

OCTOBER

1945

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Day	Max	imum	Min	imum		==					1			
	grees					Iumic	r of Hours Sunshine			2				
1	Ă	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Bright Su	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
2 3 4 5 6 7 8	59 69 50 55 67 68	12 P 9 A 0 A 3 P 3 P 12 M	35 50 30 25 34 50	5 A 12 P 12 P 6 A 5 A 6 A	42.0 59.5 40.0 40.0 55.5 59.0	97 79 48 59 70 81	0.0 0.0 9.0 10.0 8.7 2.3	s sw w se se se	31 244 336 66 119 77	8 20 39 10 11 10	9:30 A 2 A	8:30 P 1 P	.35	
9 10 11	67 75 55 58 55	4 P 3 P 1 A 2 P 3 P	56 48 38 32 33	12 P 7 A 12 P 6 A 12 P	61.5 61.5 46.5 45.0 44.0	91 59 78 68 67	1.1 9.7 2.3 8.5 6.0	nw se w nw nw	54 121 153 46 54	11 14 18 8 5	5 P 6:30 A	5:30 p	.63	
12 13 14 15 16 17 18 19	59 57 49 50 51 65 77 81	3 P 1 P 1 P 2 P 5 P 3 P 3 P	31 41 32 32 37 28 37 41	3 A 12 P 6 A 1 A 12 P 7 A 7 A	45.0 49.0 40.5 41.0 44.0 46.5 57.0 61.0	90 63 84 72 77 70 72 67	4.7 3.0 1.0 4.8 0.0 7.2 7.5 8.4	se w nw nw s se sw se	67 188 49 97 127 84 50 85	15 18 10 13 22 11 5	8 A 5 P	12 м	.02	
20 21 22 23 24 25 26 27 28 29 30 31	68 71 73 62 48 51 50 54 59 64 71	3 P 3 P 2 P 0 A 3 P 12 M 2 P 3 P 4 P 2 P	44 36 41 43 40 41 40 38 31 27 45 42	12 P 7 A 5 A 12 P 6 A 4 A 11 P 1 A 7 A 7 A 12 P	56.0 53.5 57.0 52.5 44.0 46.0 45.0 42.5 43.0 54.5 56.5	67 61 81 76 83 95 87 57 68 79 59	9.4 9.2 4.4 0.0 0.0 0.0 5.4 5.0 1.9 7.3 2.2	se w se se nw nw se sw w nw se se se	141 63 136 187 190 62 66 302 120 72 68 175	15 8 17	10:30 p 0 a 2 a 1 p	12 P 5 A 12 M 4:30 P	.14 .25 .25 .06 T	

^{*}Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

October, 1945	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.54, 1st Minimum 29.55, 2nd, 9th Mean semi-daily 30.004 Range 99	30.53 29.42 30.06 1.11	30.68, 1929 29.00, 1926 1.47, 1926; .76, 1899
Air Temperature, in degrees F.	:	
Highest 81, 19th Lowest 25, 4th Mean 49.5 Range 56 Highest mean daily 61.5, 7th, 8th Lowest mean daily 40.0, 3rd, 4th Mean maximum 60.9 Mean ninimum 38.0 Greatest daily range 40, 18th, 19th Least daily range 8, 24th	79.4 23.4 50.5 56.0	90.5, 1908 17, 1936 56.4, 1920; 43.2, 1890
Precipitation, in inches		
Precipitation 2.18 Snow —	3.29	8.81, 1911; .01, 1924
Maximum precipitation in 24 hrs63, 9th Number of days with .01 or more11	9	15, 1913; 1, 1897, 1924
Wind, in miles		-
Total movement	4074	5,467, 1910; 2,540, 1894
Least daily movement 31, 1st Mean hourly velocity 4.9 Maximum velocity 39, 3rd	5.4 29.5	42, 1937
Wind, direction		
Prevailing direction SSW North, days 0 Northeast, days 0 East, days 13 Southeast, days 2 Southwest, days 2 Southwest, days 3 West, days 5 Northwest, days 8	W	
Weather	69.0	
Mean relative humidity, percent 73.3 Mean cloudiness, percent 52 Number of clear days 77 Number of fair days 10 Number of cloudy days 14 Number hours bright sunshine 139 Percent of possible hours of bright sunshine 40.8 Thunder and lightning 12th	48 10 10 11 175.0 51.3	67, 1932; 18, 1924 21, 1938; 1, 1911 17, 1924; 3, 1938 19, 1896; 3, 1924 232, '23, '38; 91, 1913
First frost Oct. 4	Sept. 21	Aug. 22, '94, '95 : Oct. 15. '39

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1944.

The weather was near normal in most respects during October. The mean temperature for the month was 49.5 degrees, the normal for October being 50.5 degrees. The highest temperature was 81 degrees on the 19th. The lowest was 25 degrees on the 4th, which is the first frost of the season. The normal date for the first frost is September 21st. The heating load during October was 482 degree-days, the normal for October being 456 degree-days.

The total rainfall during the month was 2.18 inches. Normal precipitation for October is 3.29 inches. The total precipitation since January 1 to the end of October is 44.01 inches which is 7.10 inches above normal. There were only 139 hours of bright sunshine during the month compared to a normal of 175 hours. There were high winds on the 3rd and 27th, with a maximum of 39 miles per hour on the 3rd.

Dr. J. K. Shaw states: "The month of October concludes the crop season of 1945. The summer was unusual in some respects. It was unseasonably warm in March and early April. Then came cold nights with seriously damaging frosts and generally cool weather that counteracted the effect of the early warm weather. The season as a whole was wet, especially in May; there were few, if any, periods when crops or lawns suffered from drought. These conditions favored injury from many fungus diseases, but there was a bumper hay crop. Apple scab was prevalent because the spring frosts and ensuing cool weather resulted in the smallest apple crop in many years, and many growers thought it not worth while to spray their orchards.

"Conditions that injure some crops are favorable for others. On the whole, the season of $1945~{\rm was}$ neither bad nor especially good."

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 683

November 1945

Meteorological Observations

FOR

NOVEMBER

1945

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23' 31'' N. Longitude, 72° 31' 48'' W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

	ļ	Те	mpera	ature		ty	s		Wind			Precipita	tion	
.	Max	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
Day	å	i.		Ë	M	RE	N B	Pr	T _o M	De D	Be	Ha H	≽	Sn
12345667891011121314155617892021222224252627282930	54 58 63 44 47 55 55 70 71 59 40 44 61 67 51 43 47 42 41 49 35 37 40 41 41 42 35 33 33	4 P 2 P 3 P 0 A 2 P 1 P 0 A 1 P 1 P 0 A 1 P 1 P 0 A 3 P 1 P 1 P 0 A 3 P 1 P 1 P 0 A 1 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P 1	31 32 44 38 28 24 30 30 31 40 32 32 32 32 33 30 30 40 32 32 32 32 40 32 40 32 40 32 40 40 40 40 40 40 40 40 40 40	11 P 6 A 12 P 7 P 12 P 6 A 10 A 4 A 12 P 8 A 0 A 12 P 11 P 0 A 12 P 7 A 12 P 7 A 12 P 7 A 12 P 7 A 12 P 7 A 12 P 7 A 12 P 7 A 12 P 9 A 11 P 11 P 12 P 13 P 14 A 15 A 16 A 16 A 17 A 18 A 18 A 18 A 18 A 18 A 18 A 18 A 18	42.5 45.0 53.5 41.0 37.5 39.5 41.5 62.5 38.0 40.5 52.0 53.5 41.5 35.0 45.5 38.5 36.0 32.5 32.0 29.5 33.5 32.0 29.5 33.5 32.0 29.5 33.5 32.0 29.5 33.5 32.0 29.5 33.5 32.0 29.5 33.5 33.5 33.5 33.5 33.5 33.5 33.5 3	63 81 87 86 59 75 76 74 85 99 87 75 47 87 77 67 71 87 72 63 75 75 88 75 75 88 75 75 75 75 75 75 75 75 75 75 75 75 75	8.7 0.7 0.0 0.0 8.0 7.3 3.6 0.0 0.0 0.0 0.0 3.6 5.0 3.2 7.4 6.1 6.8 8.3 0.0 0.0 0.5	nw se nw nw se se se se se nw nw nw se nw nw w se nw nw w se nw nw w se nw nw w se nw nw w se nw nw w se nw nw w se nw nw w se nw nw w se nw nw w se nw nw w se nw nw w se nw nw w se nw nw nw w se nw nw nw se nw nw nw se nw nw nw se nw nw nw	83 60 103 108 139 30 59 180 203 156 102 105 135 235 38 171 97 353 115 131 95 73 91 44 298 390	13 12 18 14 15 5 12 12 17 14 15 14 20 13 32 8 4 22 22 16 6 6 6 15 20 22	5:30 P 5 A 12 M 8 A 9 A 0 A 1 A 4 A 4:30 A 6 P 8 A 8 A 0 A 6 A	6 P 7 A 12:30 P 12:30 P 12 P 12 P 6 A 12 P 6 A	.01 .06 .01 .10 .20 .24 .27 .10 .01 .02 .03 .71 .55 .93	.25

^{*}Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

November, 1945	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.65, 11th Minimum 29.54, 19th Mean semi-daily 30.048 Range 1.11	30.59 29.30 30.05 1.29	30.87, 1932 28.73, 1904 1.84, 1904; .89, 1943
Air Temperature, in degrees F.		
Highest 71, 9th Lowest 16, 27th Mean 40,3 Range 55 Highest mean daily 62.5, 9th Lowest mean daily 27.5, 24th Mean maximum 48.8 Mean minimum 31.9 Greatest daily range 31, 6th Least daily range 4, 11th	66.0 12.9 38.9 53.0	75, 1924, 1938 -4, 1938 44.1, 1931; 33.6, 1901
Precipitation, in inches	2 11	0.61 1007 62 1017
Precipitation	3.41 2.34	8.64, 1927; .63, 1917 13.50, 1938
Number of days with .01 or more 17	9	22, 1921; 2, 1904
Wind, in miles		
Total movement4258Greatest daily movement395, 15thLeast daily movement29, 27thMean hourly velocity5.9Maximum velocity32, 15th	4531 6.3 30.3	5,978, 1906; 2,589, 1889 44, 1938
Wind, direction		
Prevailing direction WNW North, days 1 Northeast, days 0 East, days 0 Southeast, days 1 Southwest, days 1 Southwest, days 2 West, days 4 Northwest, days 11	WNW	
Weather		
Mean relative humidity, percent 76 Mean cloudiness, percent 64 Number of clear days 7 Number of fair days 8 Number of cloudy days 15 Number hours bright sunshine 95 Percent of possible hours of bright sunshine 32 Thunder and lightning 7 First snow Nov. 29	70.6 55.1 6 9 15 121 41.3 Nov. 6	72, 1900, 1927; 34, 1917 15, 1903; 1, 1900, 1911, 1927 16, '12; 4, '89, '30, '32, '35 24, 1927; 9, 1905, 1917 182, 1903; 66, 1927 Oct. 10, '25; Nov. 27, '31

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1944.

November weather was near normal in most respects. The snowstorm on the 29th was not very severe at Amherst, with a fall of only 3 inches, but was a severe storm in the eastern part of the state. It was the first snow of the season. It was the latest date for the first snow since the station was established. The normal date for the first snow is November 6 and the normal snowfall in November is 2.34 inches. The total precipitation for the month was 3.54 inches, normal being 3.41 inches. The mean temperature for the month was 40.3 degrees whereas the normal mean temperature for the month was 16 degrees on the 27th. The domestic heating load for the month was 740 degree-days, normal being 784 degree-days. The total heating load to the end of November is 1332 degree-days. The normal heating load through September, October, and November is 1389 degree-days. There were 95 hours of bright sunshine compared to a normal of 121 hours. There were no high winds, the highest velocity being 32 miles per hour on the 15th. The mean relative humidity was 76 percent, the normal being 70.6 percent.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorogical Series

Bulletin No. 684

December 1945

Meteorological Observations

FOR

DECEMBER

1945

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Те	mpera	ture		ty	s		Wind			Precipita	ation	
	Max	imum	Min	imum		Iumidi	r of Hours Sunshine							
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Bright Sur	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8 9	35 37 36 37 29	1 P 2 P 8 P 1 P 3 P	23 13 18 26 17	12 P 8 A 5 A 12 P 7 A	29.0 25.0 27.0 31.5 23.0	62 69 88 85 39	3.6 7.6 0.0 0.0 5.0	nw nw se nw	325 88 53 102 104	22 14 6 12 12	10:30 a	11 A	.01	0.3
6 7 8 9 10	37 41 49 46 36 30	3 P 2 P 1 P 2 P 0 A 2 A	22 34 31 24 28 12	2 A 0 A 11 P 8 A 11 P 12 P	29.5 37.5 40.0 35.0 32.0 21.0	92 99 68 89 67 49	0.0 0.0 8.0 6.2 0.0 8.0	nw nw w se w	172 234 185 81 116 354	14 13 17 13 12 28	5 A 3 A	12 P 4:30 P	.83 .70	0.2 T
12 13 14 15 16 17	21 26 25 17 19 21	1 P 1 P 1 P 6 P 2 P 3 P	7 9 10 4 3 1	7 A 8 A 12 P 6 A 3 A 5 A	14.0 17.5 17.5 10.5 11.0 11.0	39 44 91 94 69 61	8.0 5.0 1.6 1.0 4.0 6.6	W W W SW SW	325 163 9 0 102 123	32 20 2 0 20 17	2 а	8 A	.23	5.8
18 19 20 21 22 23	25 16 21 20 22 22	2 P 8 P 1 P 3 P 2 P 2 P 3 P	5 5 7 3 -1 -3	7 A 2 A 10 P 6 A 12 P 1 A	15.0 10.5 14.0 11.5 10.5 9.5	56 87 79 74 59 62	7.3 0.0 6.9 6.6 4.6 8.0	sw nw nw w w	54 114 124 66 44 212	8 15 17 15 28 28	11 A 0 A	12 p 2 a	.21 .03	5.3 0.2
24 25 26 27	26 34 45 36 35	3 P 12 P 3 P 0 A 3 P	9 4 34 20 14	8 A 3 A 0 A 12 P 8 A	17.5 19.0 39.5 28.0 24.5	60 96 87 61 71	8.2 0.0 2.0 4.3 7.0	w w w	208 24 99 238 21	15 7 14 28 6	7 р 0 а	12 P 5 A	.40 .51	Т
28 29 30 31	35 29 33 37	9 P 11 P 4 P	21 13 27	8 A 0 A 8 A 12 P	25.0 23.0 32.0	92 95 100	0.0 0.0 2.5 0.0	sw nw s w	117 43 26	14 9 9	4 A 0 A 9 A	12 P 12 P 4 P	.56 .23 .20	7.0

^{*}Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

December, 1945	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.43, 24th Minimum 29.26, 31st Mean semi-daily 30.004 Range 1.17	30.66 29.27 30.06 1.38	30.96, 1889 28.85, 1915 1.78, 1895; 1.01, 1892
Air Temperature, in degrees F.		, , , , , , , , , , , , , , , , , , , ,
Highest 49, 8tn Lowest -3, 23rd Mean 22.3 Range 52 Highest mean daily 40, 8th Lowest mean daily 9.5, 23rd Mean maximum 30.4 Mean minimum 14.2 Greatest daily range 30, 25th Least daily range 7, 7th	54.3 -1.9 27.5 56.2	65.5, 1908 -22.5, 1917 36.9, 1891; 17.1, 1917
Precipitation, in inches		
Precipitation 3.91 Snow 18.8 Maximum precipitation in 24 hrs.	3.39 8.50	7.77, 1901; .58, 1943 26.50, '02; t, 1891, 1943
Number of days with .01 or more 11	10	17 , 19 02; 4, 1892, 1943
Wind, in miles		
Total movement3926Greatest daily movement354, 11thLeast daily movement0, 15thMean hourly velocity5.3Maximum velocity32, 12th	4710 6.3 31.4	6,694, 1925; 3,239, 1918 48, 1938, 1943, 1944
Wind, direction		
Prevailing direction WNW North, days 0 Northeast, days 0 East, days 0 Southeast, days 2 South, days 1 Southwest, days 4 West, days 12 Northwest, days 12	WNW	
Weather		
Mean relative humidity, percent 73.7 Mean cloudiness, percent 56 Number of clear days 11 Number of fair days 7 Number of cloudy days 13 Number hours bright sunshine 122 Percent of possible hours of bright sunshine 43.1 Thunder and lightning	69.4 54.9 8 9 14 128	71, 1929; 39, 1919 15, 1890; 2, 1922, 1933 16, '09; 4, '80'30, '31, '36, '38 23, '33; 7, '09, '23 172, 1896; 63, 1933

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1944

ANNUAL SUMMARY

Annual, 1945	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum30.69, Apr. 2	30.81	31.05, 1920, 1943
Minimum28.98, Jan	28.95	28.41, 1938
Mean semi-daily30.017 Range1.7	30.01	2.47, 1938; 1.38, 1933
Air Temperature, in degrees F.		
Highest 92, June 15, 30; July	95.7	104, 1911
Lowest		-26, 1904 40.9 1031 44 1004
Mean47.9		49.8, 1921; 44, 1904
Range10. Highest mean daily80, June 30		
Lowest mean daily —1. Ian. 2.	5	
Lowest mean daily	57.8	
Mean minimum 37.	36.5	
Greatest daily range 44, Apr. 10 Least daily range 4, Mar. 22; Nov. 1		
Precipitation, in inches		
Precipitation51.4	43.70	59.00, 1938; 30.68, 1908
Snow65.30	47.78	89.00, 1893; 24.50, 1919
Snow65.30 Maximum precip. in 24 hrs. 2.83, Apr. 2.		
Number of days with .01 or more146	124	144, 1902; 96, 1924
Wind, in miles	F0.022	
Total movement 47,149 Greatest daily movement 484, Apr. 2	52,223	63,571, 1908; 36,257, 1894
Least daily movement		
Mean hourly velocity 5.5	5.8	
Mean hourly velocity5. Maximum velocity45, Jan. 24	39.5	80, 1938
Wind, direction		
Prevailing direction Worth, days 10	W	
North, days		
East, days	5	
Southeast, days10		
South, days1	ól	
Southwest, days20	2	
West, days8		
Northwest, days11		
Weather	(7.6	
Mean relative humidity, percent	67.6 51.7	60, '98, '01, '02; 41, '08, '24
Number of clear days 13	116	217, 1941; 59, 1927
Number of fair days12	1 123	182, 1912 : 64, 1936
Number of cloudy days 11.	126	179, '01, '02; 71, 1910, 1941
Number hours bright sunshine 233.	2,353	3038, 1941; 1864, 1902
Percent of possible hours of bright sunshine 52.	52.8	
Last snow May 1	1 April 15	Mar. 14. '10: May 11 '07
Last snow May 1 First snow Nov. 2	Nov. 6	Oct. 10, '25; Dec. 13, '41
Last frost Apr. 2 First frost Oct.	4 May 14	Mar. 14, '10; May 11, '07 Oct. 10, '25; Dec. 13, '41 Apr. 14, '43; June 8, '32 Aug. 22, '94, '95; Oct. 13, '09
	4 Sept. 21	1 A 22 701 705 . Ch. 4 12 700

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 685

January 1946

Meteorological Observations

FOR

JANUARY

1946

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Te	mpera	ture		ty	s		Wind			Precipita	tion	
	Max	imum	Min	imum		lumidi	Hour			ys.				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max, Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 22 23 24 25 26 26 27 28 29 20 20 20 20 20 20 20 20 20 20 20 20 20	27 21 29 39 46 50 49 44 43 35 42 39 36 24 31 18 26 32 33 33 28 29 35 36 31 17 25	0 A A 1 P 3 P 3 P 3 P 3 P 6 P 11 A 3 P 6 P 12 P 12 P 14 A A 12 P 10 A 3 P 14 P 2 P 10 A 3 P 12 P 11 P 12 P	17 5 1 23 31 41 40 28 27 32 32 32 32 31 11 0 17 1 -4 14 14 9 9 -7 -3 26 6 17	12 P 12 P 7 A 1 A O A 12 P 112 P 1 A O A 8 A 5 A P 8 A 12 P 8 A 12 P 8 A 12 P 7 A 12 P 7 A 12 P 7 A 12 P	22.0 13.0 15.0 31.0 38.5 45.5 44.5 36.0 37.5 37.0 21.0 9.0 21.0 21.0 23.0 21.0 21.0 50.0 13.5 14.0 21.0 21.0 10.5 13.0 30.5 24.0 14.0 20.5 11.5 21.0	56 50 70 79 90 94 65 88 62 50 86 49 49 66 70 46 47 92 68 59 46 47 92 67 70 89 68 89 89 89 89 89 89 89 89 89 8	6.0 9.1 7.7 4.0 0.2 0.0 6.0 0.0 3.8 8.0 0.0 7.0 5.3 3.0 9.5 5.7 3.4 4.0 1.8 6.3 9.8 6.3 9.8 6.3 9.8 6.0 6.0 6.0 6.0 6.0 6.0 6.0 6.0	W nw se se se se s nw w w se w nw nw w nw	281 218 39 47 120 198 55 57 17 206 175 70 341 242 150 117 9 57 375 151 196 48 39 226 254 111 56 270 48 33	20 18 6 7 9 12 7 10 6 25 25 25 12 14 40 22 18 20 2 32 32 20 15 25 25 25 25 25 25 25 25 25 2	11:30 A 1 P 0 A 3 A 2 A 7 A 1 A 5 A 0 A 5 P 0 A	12 M 12 P 1 A 8 A 2:30 A 2:30 A 12 P 4 A 12 P 2 P	.01 .40 .02 .15 .03 T T .03 .53 .10 .37 .40	0.66 T T 7.00 3.7 4.00

^{*}Based on least time required to blow one mile.

FERDINAND E. BARTLETT, Observer

January, 1946	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.70, 20th Minimum 29.41, 31st Mean semi-daily 30.118 Range 1.29	30.70 29.20 30.07 1.50	31.00, 1927 28.55, 1913 2.18, 1913; 0.97, 1896
Air Temperature, in degrees F.		
Highest 50, 6th Lowest -8, 28th Mean 23.2 Range 58 Highest mean daily 45.5, 6th Lowest mean daily 6.0, 20th Mean Maximum 31.9 Mean Minimum 14.4 Greatest daily range 35, 23rd Least daily range 7, 12th	51.1 -7.1 24.2 58.2	66.0, 1932 -26.0, 1904 34.2, 1913; 13.9, 1918
Precipitation, in inches		
Precipitation 2.72 Snow 20.3	3.61 13.32	7.15, 1898; 1.07, 1896 33, 1898, 1923; 1.50, '08
Maximum precip. in 24 hrs 0.77, 24th, 25th Number of days with .01 or more12	11	15, '93, '20 ; 4, 1901
Wind, in miles		
Total movement 4421 Greatest daily movement 375, 19th Least daily movement 9, 17th Mean hourly velocity 5.9 Maximum velocity 40, 13th	5055 6.8 32.7	7770, 1908; 2896, 1895 47, 1938
Wind, direction		,
Prevailing direction W North, days	WNW	
Weather		
Mean relative humidity, percent 66.6 Mean cloudiness, percent 58 Number of clear days 9 Number of fair days 13 Number of cloudy days 9 Number hours bright sunshine 143 Percent of possible hours of bright sunshine 48.6 Thunder and lightning -	70.1 55 9 13 137 46.7	78, 1932; 37, 1908 20, 1944; 2, 1914 18, 1890, 1926; 3, 1916, 1944 22, 1923,1931; 4, 1920, 1939 214, 1920; 74, 1932

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1945.

The weather during January was slightly colder than normal and there was more snow than usual. The mean temperature was 23.2 degrees while the normal mean temperature for January is 24.2 degrees. The lowest temperature was –8 degrees on the 28th and the highest, 50 degrees on the 6th. The average minimum temperature in January is –7.1 degrees. The domestic heating load for the month was 1296 degree-days as compared to a normal of 1265 degree-days. The heating load for the season to the end of January is 3952 degree-days, the normal for this period being 3817 degree-days.

The total snowfall for the month was 20.3 inches. Usually we get 13.32 inches of snow in January. The total precipitation for the month was 2.72 inches, the normal being 3.61 inches. There were 143 hours of bright sunshine as compared to a normal of 137 hours. There were high winds on January 13, the maximum being 40 miles per hour.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 686

February 1946

Meteorological Observations

FOR

FEBRUARY

1946

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Te	mpera	iture		ty	s	•	Wind			Precipita	tion	
	Maximum		Minimum		mum		r of Hours Sunshine			s				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Bright Su	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17	38	1 P 0 A 2 P 4 P 9 P 0 A 3 P 3 P 4 P 2 P 0 A 12 P	22 12 P 14 7 A 4 12 P -8 7 A -4 7 A 13 0 A 23 12 P 12 8 A 10 1 A 17 12 P 7 8 A 25 1 A 31 12 P 7 12 P 7 12 P	26.5 26.0 12.0 8.5 12.0 28.5 30.5 22.5 23.5 26.5 19.5 28.5 32.0 41.0 19.0 14.0 33.5	52 55 56 56 55 89 67 52 73 52 58 56 77 84 49 51	5.0 8.0 10.1 9.5 9.5 0.0 1.7 10.2 0.0 10.3 10.3 10.4 0.0 0.5 4.2 9.0 7.0	w se w nw w se nw nw se se se w se w	279 183 291 143 98 56 224 164 44 206 147 80 178 232 383 303 125	183 25 291 28 143 13 98 8 56 22 224 25 164 18 44 7 206 22 147 15 80 13 178 13 232 22 383 48 303 40	8 A 0 A 5 P 3 P	12 P 12 M	.49 .03	Т	
18 19 20 21 22 23 24	29 21 31 21 29 41 33	3 A 8 P 3 P 0 A 3 P 3 P 3 P	13 4 18 7 2 18 21	12 F 7 A 1 A 9 A 7 A 1 A	21.0 12.5 24.5 14.0 15.5 29.5 27.0	43 51 83 51 61 67 75	10.7 3.0 4.5 10.8 7.0 9.5 2.5	w nw w w se nw	297 82 329 499 130 38 64	32 50 40 17 6	8 P 0 A	12 P 1 P	.02 1.13	0.30 13.00 T
25 26 27 28	31 22 28 36	1 P 12 P 8 P 4 P	9 1 22 21	12 P 6 A 0 A 12 P	20.0 11.5 25.0 28.5	53 78 91 69	9.3 0.0 0.0 11.0	nw se nw nw	363 80 108 204	28 25 13 22	9 A 0 A 0 A	12 P 12 P 7 A	.30 .79 .45	3.00 1.50 0.50

^{*}Based on least time required to blow one mile.

FERDINAND E. BARTLETT, Observer

February, 1946	Normal	Extrem e s
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.61, 5th Minimum 29.23, 14th Mean semi-daily 30.019 Range 1.38	30.63 29.24 30.04 1.38	31.05, 1920 28.56, 1895 1.89, '00, '08; .88, '13, '31
Air Temperature, in degrees F.	1.00	1105, 00, 00, 100, 10, 01
Highest 51, 14th Lowest -8, 4th Mean 22.6 Range 59 Highest mean daily 41, 14th	50.5 -7.4 23.7 58	65.0, 1930 -23, 1943
Lowest mean daily 8.5, 4th Mean maximum 32.5 Mean minimum 12.7 Greatest daily range 33, 4th Least daily range 6, 27th	32.8 14.5	
Precipitation, in inches		
Precipitation 3.52 Snow 18.30 Maximum precipitation in 24 hours	3.19 13.99	8.12, 1900; 0.62, 1901 48.75, 1893; 0.50, 1937
Number of days with .01 or more9	10	15, '93, '20; 4, 1901
Wind, in miles	ļ	
Total movement5330Greatest daily movement499, 21stLeast daily movement38, 23rdMean hourly velocity7.9Maximum velocity50, 20th	7.1 31.7	6445, 1896; 3438, 1892 48, 1934, 1937, 1944
Wind, direction		
Prevailing direction W North, days 0 Northeast, days 0 East, days 0 Southeast, days 7 South, days 1 Southwest, days 0 West, days 10 Northwest, days 10		
Weather		
Mean relative humidity, percent Mean cloudiness, percent Number of clear days Number of fair days Number of cloudy days Number of hours bright sunshine Percent of possible hours of bright sunshine Thunder and lightning 63 63 63 64 65 65 66 67 68 68 68 68 68 68 68 68 68 68 68 68 68	50.9 10 8 10 158	66, '90, '27; 31, 1905 19, 1941; 2, 1927 16, 1920; 2, 1936 16, 1894; 2, 1926 221, 1924; 110, '27, '38

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1945.

February was a severe winter month. The mean temperature was 22.6 degrees, the normal being 23.7 degrees. This makes the month the coldest February since 1936. The lowest temperature was –8 degrees on the 4th and the highest, 51 degrees on the 14th. The heating load during the month was 1187 degreedays, the normal being 1165 degree-days. The total heating load from September to the end of February is 5139 degree-days as compared to a normal of 4982 degree-days. The snowfall during the month was 18.30 inches. The normal snowfall for February is 13.99 inches. The total snowfall for the season is now 60.4 inches. The normal snowfall to the end of February is 38.15 inches and the normal snowfall for the year is 47.78 inches.

There were 174 hours of bright sunshine. This is well in excess of the 158 hours which is normal for February. The relative humidity was 63 percent compared to a normal of 66.6 percent.

There were high winds on the 15th and 20th. On the former date α velocity of 48 miles per hour was recorded and on the 20th the maximum was 50 miles per hour.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 687

March 1946

Meteorological Observations

FOR

MARCH

1946

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

	Temperature					ity	e e		Wind			Precipita	tion	
		imum		imum		Mean Relative Humidity	Number of Hours Bright Sunshine	ing E	ent	iles or				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relativ	Number	Prevailing Direction	Total Movement	Max. Miles per Hour's	Began	Ended	Water	Snow
1 2 3	39 36 44	2 P 10 A 11 A	8 32 30	7 A 0 A 12 P	23.5 34.0 37.0	76 89 69	11.2 0.1 8.7	S HW W	51 113 146	10 14 39	2 P	7 r	.12 T	
4 5 6	58 50 55	5 P 4 P 5 P	28 37 34	6 A 7 A 7 A	43.0 43.5 44.5	55 70 65	9.5 5.0 9.1	s nw e	87 40 110	8 11 22	5 Р	5:30 р	Т	
7 8 9	48 51 56	2 р 3 р 11 л	37 31 32	7 A 7 A 12 P	42.5 41.0 44.0	90 83 59	1.2 9.7 4.8	se w sw	126 44 250	13 6 32	3 Р 7 Р 0 л	9 1 12 1 8 2	.29 .24 .22	
10 11 12	40 39 47	3 P 3 P 3 P	26 21 21	12 P	33.0 30.0 34.0	48 62 62	11.6 7.8 11.7	nw nw	327 138 147	32 13 13	5 Р	5:30 р	Т	T
13 14 15 16 17	70 75 58 61 57	3 P 4 P 0 A 5 P 3 P	35 39 33 29 30	7 A 7 A 12 P 6 A 6 A	52.5 57.0 45.5 45.0 43.5	57 62 88 71 67	9.5 8.0 0.0 11.9 7.2	se W n W	209 122 97 43 78	20 15 11 5 9	8 .1	11 :30 л	.03	
18 19 20 21 22 23 24 25 26 27 28 29 30 31	57 55 63 69 73 51 65 60 71 67 79 83 60 45	3 P 4 P 3 P 4 P 4 P 3 P 4 P 3 P 4 P 3 P 3 P 4 P 3 P 1 P	28 27 24 27 31 30 25 40 37 48 46 47 36 28	6 A 6 A 7 A 6 A 7 A 5 A 8 A 7 A 4 A 7 A 12 P 5 A	42.5 41.0 43.5 48.0 52.0 40.5 45.0 50.0 54.0 57.5 62.5 65.0 48.0 36.5	58 64 57 58 54 57 59 87 82 44 52 57 49	12.0 12.0 12.1 12.2 10.5 12.3 10.5 5.5 8.0 12.5 12.5 12.6 10.8	e e nw sw s nw se e s w w se nw se nw se e s mw se e nw nw se nw se nw se nw nw nw nw nw nw nw nw nw nw nw nw nw	136 105 88 77 128 342 92 116 125 222 206 156 262 254	13 9 13 8 11 28 11 17 18 32 22 17 20 20	3 A 8 P	10 A 10 P	.51 .19	

^{*} Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

March, 1946	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.46, 12th Minimum 29.44, 9th Mean semi-daily 30.073 Range 1.02	30.57 29.24 30.00 1.34	31.05, 194, 28.47, 191- 2.10, 1914; 0.85, 1913
Air Temperature, in degrees F.		
Highest 83, 29th Lowest 8, 1st Mean .44.5 Range .75 Highest mean daily 65, 29th	63.8 6.1 34.4 57.6	85, 29t1 -7.5, 1906
Mean 44.3 Range 75 Highest mean daily 65, 29th Lowest mean daily 23.5, 1st Mean maximum 57.8 Mean minimum 31.5 Greatest daily range 42, 21st, 22nd Least daily range 4, 2nd	43.2 25.4	
Precipitation, in inches		
Precipitation	3.70 7.47	7.89, 1942; 0.12, 1915 27, 1899; 0, 1921
Number of days with .01 or more	11	17, 1890, 1913, 1936; 3, 1915
Wind, in miles		
Total movement	5944	8182, 1896; 3006, 1905
Mean hourly velocity 6.0 Maximum velocity 39, 9th	7.7 31.2	48, 1932, 1939, 1941, 1942
Wind, direction		
Prevailing direction WSW North, days 1 Northeast, days 0 East, days 5 Southeast, days 5 South, days 5 Southwest, days 2 West, days 6 Northwest, days 8	WNW	
Weather		
Mean relative humidity, percent 64.2 Mean cloudiness, percent 36 Number of clear days 21 Number of fair days 7 Number of cloudy days 3 Number of hours bright sunshine 283 Percent of possible hours of bright sunshine 76.3 Thunder and lightning 76.3	64.2 51.8 11 10 10 199 53.6	68, 1901; 27, 1915 22, 1924; 3, 1901 17, 1900; 1, 1945 21, 1901; 1, 1915 292, 1924; 93, 1901

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1945.

The weather during March was unusually warm, with much sunshine, very little rainfall and only a trace of snow. The mean temperature for the month was 44.5 degrees which exceeds the all-time record for March set last year by one-tenth of a degree. The normal mean temperature for March is 34.4 degrees. The maximum temperature was 83 degrees on the 29th and the lowest, 8 degrees on the 1st. Last year the maximum temperature in March was 85 degrees which also occurred on the 29th, and this is the highest reading ever recorded in March. The college pond was clear of ice on March 20. The heating load during the month was 636 degree-days, while the normal for March is 950 degree-days. The total load for the season to the end of March is 5775 degree-days, the normal for this period being 5932 degree-days.

The total precipitation was 1.60 inches, all of which fell as rain except for a light snow squall on the 11th. Normally we get 7.47 inches of snow, with a total precipitation of 3.70 inches. The total snowfall for this winter to the end of March is 60.40 inches, which is the heaviest seasonal fall since the winter of 1925-26 when 61.00 inches was recorded. The normal for the same period

is 45.62 inches.

There were 283 hours of sunshine, compared to a normal of 199 hours; and 21 days were classified as clear. The mean relative humidity was 64.2 percent, while the normal for March is also 64.2 percent. High winds were recorded on the 3rd, 9th, 10th and 27th.

Dr. J. K. Shaw makes the following report on the progress of vegetation: "The temperature this month has been slightly higher than it was during the abnormally warm March of 1945. Vegetation has advanced to about the same stage and our comments of a year ago can now be repeated. 'The extremely high temperatures in March have advanced vegetation so that it is about as far along as it was a month later last year. Grass is green and leaves are developing on many shrubs. Forsythia is in full bloom at the last of the month. It is, so far, the earliest spring in many years.' The month closes with cooler weather; if this persists, it could delay bloom of fruit trees until danger of frost injury is decreased. It would require a temperature around 20 degrees or possibly a little lower to cause serious injury to fruit buds at the present stage of development. We can be quite sure that bloom will be earlier than normal and that there will be a decreasing probability of cold injury until late May or even early June. We can hope to escape cold injury this year, but we may as well admit that the fruit crop of 1946 is in great danger.

"Fruit trees came through the winter in good condition. No winter injury has been observed, and most fruit trees are full of flower buds. Few peach fruit buds were killed and the prospect is for a heavy bloom. The fruit crop of this year hangs in the balance. If it escapes cold injury, there will be a big crop. If

cold comes, it may be reduced even to destruction."

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 688

April 1946

Meteorological Observations

FOR

APRIL

1946

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

April, 1946	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.50, 1st Minimum 29.29, 26th Mean semi-daily 29.956 Range 1.21	30.48 29.38 29.99 1.11	30.71, 1911 28.98, 1943 1.52, 1930; .72, 1919
Air Temperature, in degrees F.		1102, 1700, 112, 1717
Highest 78, 23rd Lowest 19, 1st Mean 45.0 Range 59 Highest mean daily 59.5, 24th	79.4 22.0 45.7 57.4	90, 1941 8.5, 1923 52, 1921; 41, 1943
Lowest mean daily27.5, 1stMean maximum55.7Mean minimum34.2Greatest daily range43, 23rdLeast daily range7, 27th	56.8 34.8	
Precipitation, in inches		
Precipitation 2.16 Snow 1.50 Maximum precip. in 24 hrs	3.35 2.17	6.89, 1929; .55, 1941 11, 1891; 0, 1910, 1934
Number of days with .01 or more10	11	18, 1909; 3, 1892
Wind, in miles		
Total movement 5630 Greatest daily movement 408, 5th Least daily movement 63, 22nd	5404	8208, 1908; 3853, 1917
Mean hourly velocity 7.8 Maximum velocity 35, 2nd	7.5 31.4	40, 1935, 1938, 1940, 1945
Wind, direction		
Prevailing direction W North, days 2 Northeast, days 0 East, days 0 Southeast, days 7 South, days 1 Southwest, days 2 West, days 10 Northwest, days 8	WNW	
Weather		
Mean relative humidity, percent 64.6 Mean cloudiness, percent 53 Number of clear days 13 Number of fair days 9 Number of cloudy days 8 Number of hours bright sunshine 228 Percent of possible hours of bright sunshine 56.7 Thunder and lightning 2nd, 3rd Last snow April 8	61.6 51.8 11 9 10 220 54.7 Apr. 15	75, 1901; 34, 1925, 1927 23, 1941; 3, 1898, 1902 18, 1915; 2, 1894, 1901 22, 1901; 1, 1941 329, 1941; 103, 1901 Mar. 14, 1910; May 11, 1907

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1945.

====		Temperature			Temperature							Wind			Precipita	tion	
		Tempe	lature			idity	ours		1	1		rrecipita					
	Maximum		Minimum			Imm	f H nshi	bo.		S							
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow			
1 2 3 4 5 6 7	36 65 55 45 47 48 48	12 M 4 P 9 A 3 P 1 P 2 P 3 P	19 32 33 30 35 30 31	6 A 0 A 12 P 3 A 10 P 5 A 12 P	27.5 48.5 44.0 37.5 41.0 39.0 39.5	66 64 69 94 69 61 52	4.9 7.6 4.7 0.0 2.0 3.3 13.0	se w nw se w sw	117 158 289 177 408 257 292	12 35 32 20 28 20 20	2:30 P 0 A 0 A	12 P 12 P 1 A	.15 .53 .15	0.50 1.00			
8 9 10 11 12 13 14	49 49 47 50 49 49 64	3 P 3 P 3 P 1 P 5 P 2 P	31 24 33 32 28 29 33 34	12 P 6 A 0 A 12 P 6 A 5 A 5 A	39.5 36.5 41.0 39.5 39.0 39.0 41.0 49.0	52 68 88 59 52 56 65 55	9.6 3.6 9.4 10.0 8.6 7.0 13.0	nw se w w nw n	104 140 322 212 116 65 120	20 17 22 25 20 15 7	7:30 P	9:30 p	.07	Т			
15 16 17 18 19 20	63 51 61 66 58 54	6 P 0 A 4 P 1 P 5 P 10 A	42 39 28 28 40 40	0 A 9 P 4 A 6 A 5 A 12 P	52.5 45.0 44.5 47.0 49.0 47.0	76 55 40 58 47 64	0.0 13.2 13.4 8.6 10.6 7.2	se w w se w nw	230 390 95 204 241 330	18 32 14 22 22 32	11 A	8 Р	.10				
21 22 23 24 25	61 68 78 73 70	1 P 4 P 5 P 3 P 2 P	35 36 35 46 38	7 A 5 A 5 A 6 A 5 A	48.0 52.0 56.5 59.5 54.0	42 55 55 64 74	13.6 12.6 12.0 7.9 9.3	nw s w n se	168 63 80 118 96	18 13 13 14 11	3:30 \	6 л	.05				
26 27 28 29	54 43 53 48 70	0 A 7 A 4 P 11 A	46 36 36 36	12 P 12 P 0 A 3 A	50.0 39.5 44.5 42.0	97 91 52 80	0.0 0.0 11.3 0.0	nw nw w nw	182 178 259 78	18 18 20 7 12	1 A 6 A 3 P	2 P 12 M 9 P	.82 .11				
30 31	70	4 Р	41	0 а	55.5	70	11.6	se	141	12							

^{*} Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

The weather during April was dry and slightly cooler than normal. The precipitation during the month was only 2.16 inches, normal for April being 3.35 inches. The snow fall during the month was 1.50 inches, the normal snowfall for April being 2.17 inches.

The mean temperature during April was 45.0 degrees. This is only half a degree higher than the mean during March this year. The normal mean temperature for April is 45.7 degrees. The lowest temperature was 19 degrees on the 1st. The domestic heating load for the month was 602 degree-days compared to a normal of 579 degree-days. The heating load for the season is 6377, the normal to the end of April being 6511 degree-days.

There were 228 hours of bright sunshine which is 8 hours more than normal. Mean relative humidity was 64.6 percent, the normal being 61.6 percent.

Dr. J. K. Shaw makes the following report on the progress of vegetation:

"While March 1946 was very similar to March 1945, the month of April 1946 was distinctly cooler than April 1945. As a result, vegetation advanced very slowly. At the end of the month most apple varieties were in the pink stage of bloom and will be in full bloom with two or three warm days, probably about two weeks ahead of normal instead of three weeks or more as last year. On the morning of April 1 there was considerable cold injury to fruit buds. In our orchards no serious crop damage occurred. In other orchards about the state, there was more injury. Possibly some low lying orchards suffered enough to reduce or ruin the crop, but it seems very unlikely that the crop over the state was much reduced. We have had several narrow escapes since, but we believe that there has been no further damage.

"Practically all fruit trees in our orchards show extremely heavy bloom. Rarely and probably never have we seen our orchards so full of flowers in all our experience of nearly 40 years. The season of 1945 must have been very favorable for fruit bud formation. The crop loss was one reason, but there must have been others.

"While every day brings us nearer the time when we can feel assured of a good crop, there is still danger of cold injury. A minimum temperature of 27° or 28° would cause heavy damage. The next two weeks will be the critical period. While injurious cold may occur after the middle of May, it is not probable."

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 689

May 1948

Meteorological Observations

FOR

MAY

1946

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

	,	Гетре	rature			ity	ırs e		Wind		Pı	ecipitatio	n	
Day	Maxi	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 4 5 6 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 30 31	72 666 57 57 59 69 59 56 67 65 62 61 68 65 65 65 73 75 70 80 86 78 66 78 66 78 66 78 86 86 78 86 86 86 86 86 86 86 86 86 86 86 86 86	5 P 4 P 7 3 P 2 P 6 P 7 P 1 P 7 P 1 P 7 P 1 P 7 P 1 P 7 P 1 P 7 P 1 P 1	41 36 30 31 40 40 32 40 41 41 48 51 53 44 43 52 48 41 45 55 60 55 40 55	5 A 5 A 6 5 A 12 P 5 A 12 P 5 A 12 P 6 A 12 P 12 P 4 A A 12 P 11 P 12 P 4 A A 5 A	56.5 51.0 43.5 44.0 44.5 53.0 49.5 52.5 55.5 55.5 55.0 56.0 58.0 63.5 59.0 62.5 70.5 69.0 58.0 62.5 70.5 69.0 58.0 69.5	47 44 44 56 99 57 60 55 36 41 75 77 41 48 75 61 86 51 53 54 70 78 99 74 49 56	14.0 14.1 11.0 0.0 8.0 6.0 5.0 8.8 14.3 0.6 0.9 14.5 9.0 4.8 1.6 4.0 9.2 10.6 4.0 14.8 14.9 8.4 7.3 0.0 0.0 11.2 15.0 15.0	w nw nw se n w se sw se nw ne w se nw se nw ne w se se nw ne w se se nw n n nw se se se se nw n nw w sw	159 202 270 111 129 233 126 159 164 241 85 81 140 115 104 97 136 129 102 72 185 161 83 118 177 131 97 252 170 108 134	14 22 13 13 10 20 17 32 22 21 10 20 15 9 8 13 9 13 8 22 17 17 10 11 10 11 11 11 11 11 11 11	2 A 3 P 0 A 3 P 7:30 A 3 A 10:30 P 6:30 A 0 A 6 A 7 P 1:30 A 0 A	12 M 12 P 6 A 3:30 P 12 M 9:30 P 12 P 7:30 A 11 P 9 P 11 P 5.30 P	.35 .23 .17 .06 .14 .29 .13 .58 .06 .93 .26 .57	

^{*} Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

May 1946	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.41, 14th Minimum 29.62, 8th Mean semi-daily 29.969 Range 0.79	30.39 29.51 29.96 0.88	30.62, 1936 29.10, 1938
Air Temperature, in degrees F.		
Highest 86, 25th Lowest 30, 3rd Mean 55.8 Range 56 Highest mean daily 70.5, 25th Lowest mean daily 43.5 3rd Mean maximum 67.7 Mean minimum 43.9 Greatest daily range 42, 30th Least daily range 8, 28th	86.1 31.2 57.1 54.9	94.5, 1896, 1911 24.0, 1900
Precipitation, in inches		
Precipitation 5.41 Snow —— Maximum precip. in 24 hrs. 1.47, 27th	3.60	7.44, 1931; .48, 1903
Number of days with .01 or more 15	12	20, 1901, 1945; 5, 1903
Wind, in miles		
Total movement 4471 Greatest daily movement 252, 28th Least daily movement 72, 20th	4504	5946, 1907; 2180, 1894
Mean hourly velocity 6.0 Maximum velocity 32, 6th, 9th	6.1 28.8	45, 1935
Wind, direction		
Prevailing direction W North, days 3 Northeast, days 1 East, days 0 Southeast, days 9 South, days 1 Southwest, days 3 West, days 6 Northwest, days 8	W	
Weather		
Mean relative humidity, percent 54.8 Mean cloudiness, percent 55 Number of clear days 12 Number of fair days 10 Number of cloudy days 9 Number of hours bright sunshine 254 Percent of possible hours of	60.7 52.1 11 11 9 252	70, 1901, 1902; 30, 1923 21, 1944; 0, 1927 17, 1907, 1926; 5, 1923 20, 1927; 2, 1941 350, 1944; 137, 1927
bright sunshine 56.1 Thunder and lightning	55.6	
		•

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1945.

REMARKS

May weather was colder and wetter than normal but neither as wet nor as cold as May, 1945.

The mean temperature during the month was 55.8 degrees, the normal being 57.1 degrees. May of last year was 54.1 degrees. The lowest temperature was 30 degrees on the 3rd. The 3rd and 4th are the only days on which the temperature fell below freezing. The highest temperature was 86 degrees on the 25th. The domestic heating load was 298 degree-days, the normal being 267 degree-days. The heating load for the season is now 6675 degree-days, the normal for the year to the end of May being 6778 degree-days.

The total precipitation for the month was 5.41 inches as compared to a normal of 3.60 inches. The rainfall in May, 1945 was 6.45 inches. The precipitation since January 1st is now 15.41 inches, while the normal for this period is 17.45 inches. Rain fell on 15 days. Rain fell on 20 days in May, 1945. There were 254 hours of bright sunshine which is two hours higher than normal. Relative humidity was 54.8 percent, the normal being 60.7 percent.

Dr. J. K. Shaw comments as follows on the progress of vegetation: "The early part of May this year was cooler than May, 1945 and the relatively slower development of vegetation of April continued. Full bloom of apples occurred during the first week in May, about two weeks ahead of normal. As in 1945, the blooming period was much longer than usual. At the end of May, development was fully back to normal. All tree fruits produced a remarkably heavy bloom.

"Cold injury during April varied in different orchards from none to severe. There were three mornings in May when the temperature was dangerously low. These were May 3, 4 and 9. On May 3 there was some further injury to apple blossoms. We can, at the end of the month, feel fairly sure that the fruit crop is past danger from cold. Contrary to common belief, peaches are more resistant to spring cold than apples and a good crop is probable; there was little or no killing from mid-winter cold. The apple crop will be much larger than that of 1945 and may be up to normal. The set of fruit of some varieties is rather poor and there may be further loss as the 'June drop' is not yet completed.

"Rainfall in May was heavy and a good crop of strawberries is assured if wet weather during June does not cause the berries to rot. The blueberry crop was somewhat reduced by cold."

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 690

June 1946

Meteorological Observations

FOR

JUNE

1946

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

JUNE, 1946	Normai	Extremes
Barometer, in inches (Readings reduced to sea level) Maximum 30.33, 16th Minimum 29.66, 18th Mean semi-daily 30.023 Range 0.67	30.30 29.55 29.95 0.75	30.54, 1889 29.24, 1902
Air Temperature, in degrees F.		
Highest 92, 27th Lowest 39, 3rd Mean 65.0 Range 53 Highest mean daily 79.5, 27th Lowest mean daily 46,5, 2nd Mean maximum 78.0 Mean minimum 52.0	91.2 40.0 65.7 51.3	101, 1919 34, 1891
Greatest daily range 37, 23rd Least daily range 7, 2nd	•	
Precipitation, in inches		
Precipitation 3.30 Snow	3.75	9,68, 1922; 0.76, 1908
Maximum precip, in 24 hrs. 1.26, 2nd Number of days with .01 or more 10	11	17, 1922, 1945; 4, 1908
Wind, in miles		
Total movement 3362 Greatest daily movemnt 308, 9th	3585	4571, 1908; 1409, 1906
Least daily movement 42, 23rd Mean hourly velocity 4.7 Maximum velocity 47, 11th	5.0 24.6	48, 1939
Wind, direction		
Prevailing direction S North, days 2 Northeast, days 1 East, days 0 Southeast, days 12 South, days 4 Southwest, days 66 West, days 3 Northwest, days 2	WSW	
Weather		
Mean relative humidity, percent 66.3 Mean cloudiness, percent 42 Number of clear days 17 Number of fair days 10 Number of cloudy days 33 Number of hours bright sunshine 331 Percent of possible hours of bright sunshine 72.4 Thunder and lightning 8th	66.9 51.1 10 12 8 257 54.0	71, 1903; 28, 1908 22, 1908, 1941, 1943; 1, 1927 23, 1912; 3, 1941 22, 1903; 1, 1923 362, 1908; 102, 1903

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1945.

		Tempe	rature	9	dity		dity		Hours hine		Wind		P	ecipitatio	n	
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hour Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow		
1 2 3 4 5 6 6 7 8 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31	666 50 73 69 67 72 72 79 90 72 72 78 82 80 73 73 77 78 85 89 91 91 92 88 88 83	0 A 6 P 3 P 1 P 2 P 12 M 3 P 1 S P 1	45 43 39 49 44 41 52 61 47 43 51 46 47 51 55 45 46 55 45 46 60 57 60 60 60 60 60 60 60 60 60 60 60 60 60	1 P 12 P 4 A A 12 P 5 A A 5 A A 11 P 12 P 12 P 12 P 12 P 12 P 12 P	55.5 46.5 56.0 59.0 55.5 56.5 75.5 57.5 57.5 57.5 64.5 71.0 65.5 63.0 65.5 65.0 65.5 72.5 76.5 72.5 76.5 77.0 79.5 76.0 74.5 75.5	844 92 633 61 72 600 67 79 444 433 722 774 71 566 60 552 78 47 59 80 67 63 59 58 67 63 67 68 79 78 80	0.0 0.0 13.6 14.1 13.3 10.0 7.0 15.3 15.4 8.3 7.4 15.3 11.3 4.5 15.3 11.3 15.3 11.8 8.0 11.0 15.3 15.3 15.3 15.3 15.3 15.3 15.3 15.3	ne n sw se nw se se se se se se se se se se se se se	110 83 52 80 100 83 92 99 308 142 159 133 113 80 60 116 166 185 85 128 123 130 42 70 82 109 144 85 83 120	12 10 5 8 22 15 13 40 28 13 47 7 7 10 11 22 10 11 7 15 6 8 9 13 12 12 13 11 11 11 11 11 11 11 11 11 11 11 11	8:30 A 3 A 1 P 7:30 P 4 P 4 A	3 P 12 P 4 A 5:30 P 8:30 P 5:30 P 8 A	.17 1.26 .07 .12 .47 .58 .27			

^{*} Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

REMARKS

June weather averaged near normal, the rainy first half of the month being offset by the hot, dry spell the last half of the month.

The mean temperature for the month was 65 degrees, the normal being 65.7 degrees. The maximum temperature was 92 degrees on the 27th, with the daily maximums for the last eight days ranging from 83 to 92. The minimum temperature was 39 degrees on the 3rd, the normal minimum for June being 40 degrees. The domestic heating load for June was 100 degreedays, the normal for this month being 68 degree-days. The total heating load for the past season is 6775 degree-days as compared to the normal of 6846 degree-days.

The total rainfall for the month was 3.30 inches, while the normal is 3.75 inches. The total precipitation since the first of the year is now 18.71 inches, compared to the normal of 21.20 inches for the same period.

There were 17 clear days this month, whereas during the month of June last year precipitation fell on 17 days. There were 331 hours of bright sunshine compared to a normal of 257 hours. The maximum wind velocity on the 11th was 47 miles an hour.

Dr. J. K. Shaw makes the following comment on the progress of vegetation: "The first part of the month of June was characterized by excessive rainfall and much cloudy, humid weather. This, together with the heavy carry-over of scabby leaves, resulted in one of the worst apple scab years for a long time; there will be a lot of scabby McIntosh this fall. The set of apples was fair and a crop nearly or quite up to average is indicated at present. In some orchards on frosty locations the crop of apples will be light or none. The crop will not come up to the promise made by the remarkably heavy bloom. Peaches generally escaped blossom killing and there should be a good crop. Weather for strawberries was favorable, plenty of rain during the growing period and drier weather during harvest. Available supplies are restricted by the reduced area devoted to this crop. The month closes with a hot period with little rainfall. Conditions have favored a good hay crop and harvesting conditions have been good so far."

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 691

July, 1946

Meteorological Observations

FOR

JULY

1946

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

July, 1946	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum	30.27 29.59	30.50, 1892 29.27, 1932
Mean semi-daily 30 040 Range 0.50	29.96 0.68	0.97, 1892; 0.47, 1938
Air Temperature, in degrees F.		
Highest 95, 19th	93.9	104, 1911
Lowest 46, 16th Mean 70.2	46.4 70.8	40, 1890, 1898 74.7, 1921; 66.3, 1891
Range	47.7	74.7, 1921, 00.0, 1091
Highest mean daily 81.5, 20th		
Lowest mean daily 62.0, 16th		1
Lowest mean daily 62.0, 16th Mean maximum 82.3		
Mean minimum 58.0		The state of the s
Mean minimum 58.0 Greatest daily range 39, 18th, 19th Least daily range 10, 22nd, 23rd, 31st		
Precipitation, in inches		
Precipitation5.30	4.10	14.51, 1897; 0.70, 1929
Snow	1.10	14.31, 1897 , 0.70, 1929
Maximum precip. in 24 hrs 3.46, 23rd		
Number of days with .01 or more	11	20, 1915; 4, 1924
Wind, in miles		
Total movement 3229	3422	5097, 1909; 1109, 1894
Greatest daily movement 204, 15th Least daily movement 48, 28th		
Mean hourly velocity 4.3	4.6	
Maximum velocity 28, 30th	28.7	44, 1936
Wind, direction		
Prevailing direction SW North, days 1	SW	
Northeast, days 0 East, days 1		
Southeast, days11		
South, days3		
Southwest, days2		
West, days 2 Northwest, days 11		
Weather Many relative housidity assured (0.2)	60.1	
Mean relative humidity, percent	68.4 50.9	70, 1943; 31, 1924
Number of clear days 21	10	22, 1923; 0, 1915
Number of fair days4	14	24, '09, '21; 5, '89, '23, '28,
Number of cloudy days6	7	36 18, 1889; 0, 1910
Number of hours bright sunshine319	268	371, 1910; 180, 1931
Percent of possible hours of bright		
sunshine 69	58	
Thunder and lightning30th		

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1945.

		Tempe	ratur	e		lity	urs		Wind		Pr	ecipitatio	on	:
		imum		imum		Humid	Mean Relative Humidity Number of Hours Bright Sunshine		ent	c. Miles Hour*				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative	Number Bright St	Prevailing Direction	Total Movement	Max. M per Hou	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 22 23 24 25 26 27 28 29 30 31 31 31 31 31 31 31 31 31 31 31 31 31	83 82 87 78 83 87 89	1 P 5 P 6 4 P 2 P 7 1 P 8 P 9 P 9 P 9 P 9 P 9 P 9 P 9 P 9 P 9	70 63 57 51 52 54 61 63 66 67 63 66 71 58 64 67 71 58 64 63 66 67 63 66 67 63 66 67 63 66 67 67 63 66 66 67 67 67 67 67 67 67 67 67 67 67	4 A A 12 P 12 P 5 A 6 A 12 P 6 A 6 A 6 A 12 P 12 P 12 P 12 P 12 P 12 P 12 P 12	76.0 72.5 69.0 66.5 68.0 70.0 74.0 77.5 70.5 70.0 72.5 63.0 62.0 65.5 69.5 75.5 81.5 66.0 73.0 72.5 63.0 63.0 63.0 63.0 63.0 63.0 65.5 66.0 71.5 75.0 66.0	78 70 71 57 63 60 63 89 63 77 80 57 64 52 55 62 56 61 67 94 99 97 85 68 55 67 77	7.6 11.2 15.2 15.2 15.2 12.6 14.6 12.9 3.8 11.6 5.8 13.2 14.5 11.0 0.0 0.8 0.0 10.6 9.2 12.3 14.6 14.5 13.2 9.7 2.0	se nw nw se se se se se w nw nw se se w nw nw se se w nw nw se se w nw nw nw se se nw	196 174 104 77 74 106 151 73 134 90 68 96 75 106 204 88 55 69 95 122 99 66 134 112 124 117 63 48 101 95 113	17 14 11 11 11 11 12 7 10 12 7 5 9 20 25 8 7 9 11 14 6 17 11 11 8 13 12 11 11 11 11 11 11 11 11 11 11 11 11	12:30 A 5 P 0 A 3 A 5:30 P	11 P 11 P 6:30 P 4:30 A	.54 .49 3.46 T .02	

^{*} Based on least time required to blow one mile.

JOHN D. HILCHEY, Observer

REMARKS

The unusual feature of July weather was the heavy rain on the 23rd. A total of 3.46 inches fell on that day. There was no rain from the 2nd to the 21st but a total of 4.49 inches fell in the storm from the 21st to the 23rd. The total rainfall for the month was 5.30 inches, normal for July being 4.10 inches. The total precipitation from January 1 to July 31 was 24.01 inches. The normal precipitation for this period is 25.30 inches. The mean temperature for July was 70.2 degrees, normal being 70.8 degrees. Rain fell on only 6 days and there were 319 hours of bright sunshine compared to a normal of 268 hours. Relative humidity was 69.2 percent. There were no high winds during the month.

Dr. J. K. Shaw, research pomologist, makes the following statement: "The dry period which began the latter part of June continued until July 22 when heavy showers relieved the situation. The strawberry crop was far enough along to mature a good crop before conditions became severe, but raspberries suffered severely. Lawns became brown where the soil was not retentive of moisture. The dry weather must have delayed the growth of apples and possibly other tree fruits but they will resume growth with a more abundant water supply. The bad scab situation caused by so much moist, rainy weather in the spring continues, and susceptible varieties will show the effects even with a continued vigorous spray program. Other diseases and insects of tree fruits have not yet caused unusual damage.

"The peach crop will be good except in orchards in frost pockets. The apple crop is spotty, spring frosts having reduced or destroyed the crop in some orchards while others will produce a nearly normal crop. It is reported that the crop in Massachusetts will be about three times that of the very short crop of 1945 but still considerably below normal.

"Cultivated blueberries suffered from spring frosts and the dry weather, during which the berries did not grow well, and when the showers came there was much cracking in varieties susceptible to such injury. The crop will be below normal."

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 692

August, 1946

Meteorological Observations

FOR

AUGUST

1946

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Тетре	rature	2		lity	ırs e		Wind		F	Precipitati	on	
13	Max	Lime	Degrees	Lime	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max, Miles per Hour*	Began	Ended	Water	Snow
Day	Ď	Ţ	ñ	E	M	RE	Na	7.2	F.Y.	M	ğ	편	×	S
1 23 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20	69 66 79 82 85 86 68 84 89 79 77 63 65 78 77 85 81 65	11 A 1 P 1 P 1 P 3 P 0 A 5 P 5 P 4 P 4 P 4 P 4 P 2 P 3 A 2 P	58 55 52 58 61 56 59 58 66 59 53 53 56 62 62 62 52	8 P 11 P 5 A 5 A 12 P 6 A 8 P 5 A 12 P 12 P 4 A 1 A 2 A 6 A 0 A 6 A	63.5 60.5 65.5 70.0 71.0 64.0 71.5 73.5 72.5 69.0 65.0 60.5 65.0 73.5 73.5 72.5 69.5 69.5 73.5	79 80 64 84 75 71 93 73 74 69 93 87 87 82 74	0.8 0.5 14.0 6.0 11.3 9.3 0.0 14.2 9.7 2.8 13.2 8.9 0.5 2.2 9.0 3.8 7.7 9.5 0.0 10.7	n n w sc sw w nw nw s se w nw nw w s se w nw nw w w w w w w w w w nw nw nw nw n	167 199 128 62 109 34 108 119 67 155 117 39 50 160 83 99 112 78	14 18 15 14 12 9 10 14 10 13 6 10 14 15 18 10 14 10 13 16 10 11 11 11 11 11 11 11 11 11 11 11 11	1:30 P 12:30 A 3:30 P 6 A 11:30 P 0 A 9:30 A 11:30 A 10:30 A 7 P 10 P 9 A	7:30 P 1 A 6:30 A 12 P 6:30 P 11 A 11:30 P 9:30 P 10:30 P 2:30 P	.08 .02 .04 .01 .73 .10 .54 .15 .55 .04	
21 22 23 24 25 26 27 28	78 81 74 75 76 75	3 P 4 P 2 P 3 P 4 P	51 59 60 50 46 49	6 A 6 A 7 A 6 A 6 A	64.5 70.0 67.0 62.5 61.0	72 76 87 71 75 76	11.0 9.5 7.0 13.0 11.6 13.2	SW S HW W H	61 51 54 47 71 89	9 7 14 7 9	3 г	3:30 р	.88	
27 28	72 83	4 P	57 52	4 A 4 A	64.5 67.5	94 78	0.0	n s	115 56	9 10	11:30 л	12 м	.02	
29 30 31	83 70 75	3 P 4 P	55 46 45	12 P 6 A 6 A	69.0 58.0 60.0	80 67 73	6.0 13.0 10.8	S 11147 SW	124 131 55	18 14 6	4 Р	8 г	.36	

^{*} Based on least time required to blow one mile.

John D. Hilchey, Observer

August, 1946	Normal	Extrem e s
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.35, 15th Minimum 29.66, 10th Mean semi-daily 29.992 Range 0.69	30.32 29.61 30.00 0.71	30.50, 1934 28.87, 1930
Air Temperature, in degrees F.		
Highest 89, 9th Lowest 45, 31st Mean 66.1 Range 44 Highest mean daily 73.5, 9th, 17th Lowest mean daily 58, 13th, 30th Mean maximum 76.7 Mean minimum 55.5 Greatest daily range 31, 9th, 28th Least daily range 8, 7th	91.6 43.4 68.6 48.1	100, 1918 34, 1940 62.4, 1903
Precipitation, in inches		
Precipitation 4.00 Snow ———	4.08	8.40, 1928; .31, 1894
Maximum precip. in 24 hrs 0.88, 23rd Number of days with .01 or more	11	16, 1892, 1933; 4, 1899
Wind, in miles		
Total movement 2863 Greatest daily movement 199, 2nd Least daily movement 34, 6th Mean hourly velocity 3.8 Maximum velocity 18, 2nd	3127 4.2 22.7	4,271, 1910; 1,920, 1894 40, 1941
Wind, direction	22.7	70, 1941
Prevailing direction W North, days 4 Northeast, days 0 East, days 0 Southeast, days 3 South, days 6 Southwest, days 3 West, days 6 Northwest, days 9	SW	
Weather		
Mean relative humidity, percent 79.2 Mean cloudiness, percent 67 Number of clear days 11 Number of fair days 10 Number of cloudy days 10 Number of hours bright sunshine 239 Percent of possible hours of bright sunshine 55.7 Thunder and lightning 23rd, 26th, 27th, 29th First frost	70.2 49.7 9 13 9 237 55.2 Sept. 21	67, 1901; 27, 1923 23, 1941; 0, 1915, 1929 25, 1912; 3, 1939 18, 1901, '28; 2, 1910, '23 332, 1941; 152, 1915, 1929 Aug. 22, '94, '95; Oct. 13, '09

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1945

REMĀRKS

August weather was colder than usual, with the total precipitation for the month only .08 inch less than normal.

The mean temperature was 66.1 degrees, while the normal for August is 68.6 degrees. This is the coldest August since 1934 when the mean temperature was 65.0 degrees. The record at this station for August is 62.4 degrees in 1903. The maximum temperature for the month was 89 degrees on the 9th, while the lowest was 45 degrees on the 31st.

The total precipitation was 4.00 inches compared to the normal for August of 4.08 inches. Rain fell on 15 days. There were 239 hours of sunshine, while the normal for August is 237 hours. There were no high winds during the month.

Dr. J. K. Shaw, research pomologist, gives the following report: "Rainfall continued to be ample during August and the effects of the dry period of June and July have disappeared. Vegetation has progressed normally. Temperatures seem rather low for August but no frosts have been reported. The apple crop will be larger than seemed possible at the time of the spring frosts though some low-lying orchards have a small crop or no crop at all. Scab on susceptible varieties will be severe. Conditions have not been favorable for good color development but abundant sunshine and cool nights may still favor color in later varieties. The peach crop is up to or above average."

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series Bulletin No. 693

September 1946

Meteorological Observations

FOR

SEPTEMBER

1946

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

> Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

	,	Гетре	rature			lity	ars e		Wind		P	recipitati	on	
	Maxi	imum	Min	imum		Humic	r of Hours Sunshine	مخ	4	s *				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Hunidity	Number Bright St	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 3 4 4 5 5 6 6 7 7 8 9 100 11 122 13 124 125 126 227 28 29 30	80 75 72 73 74 75 82 81 76 83 73 665 70 75 79 83 84 86 86 71 70 74 70 80 77 72 66	2 P 11 A A 4 P P 2 P P 4 P P 1 5 P P 3 P P 1 A P P 2 P P 4 P P 3 P P 4 P P 3 P P 4 P P 3 P P 4 P P 4 P P P A P P P A P P P P P P	45 52 46 42 44 52 57 61 60 69 52 44 40 37 42 42 42 55 55 55 59 53 63 63 64 46 53 57 41	6 A 12 P 12 P 6 A 6 A 1 A 12 P 12 P 6 A 6 A 12 P 6 A 6 A 12 P 6 A 6 A 12 P 6 A 12 P 7 A 12 P 7 A 11 P	62.5 63.5 59.0 57.5 59.0 63.5 69.5 71.0 68.0 76.0 62.5 53.5 58.5 60.5 69.5 72.5 69.5 69.0 62.0 62.0 62.0 63.0 64.5 53.5	73 77 66 67 68 76 83 81 97 93 77 68 75 74 82 77 71 90 85 89 93 72 89 85 88	13.0 9.3 10.5 11.2 11.8 11.5 9.4 4.5 6.0 1.5 6.0 12.5 12.6 12.5 12.6 12.5 12.6 12.5 12.6 11.9 9.1 11.2 6.0 3.1 0.0 8.8 11.8 11.9 11.0	sw nw nw nw ne s se e se se se sw nw me ne ne ne ne nw w sw n sw sw n e e nw w w se se nw	77 117 85 92 92 77 100 79 54 148 165 91 113 76 23 40 30 48 32 41 176 47 113 77 48 37 52 85 173	10 12 9 9 11 9 8 8 6 18 20 10 17 11 14 4 4 10 6 6 6 8 9 9 11 12 12 14 14 14 16 16 17 17 18 18 18 18 18 18 18 18 18 18 18 18 18	8 P 0 A 12:30 A 2:30 P 0 A 12:30 A	12 P 3:30 A 2 A 12 P 3 A 12 P 12:30 A	.84 .37 T .13 .13 T .140 .01 T 1.85	

^{*} Based on least time required to blow one mile.

John D. Hilchey, Observer

September, 1946	Normal	Extremes
Barometer, in inches (Readings reduced to sea level) Maximum 30.48, 13th Minimum 29.50, 30th Mean semi-daily 30.136 Kange 0.98	30.45 29.57 30.06 .88	30.65, 1924 28.41, 1938 1.99, 1938; .57, 1910
Air Temperature, in degrees F.		
Highest 86, 19th, 20th Lowest 37, 14th Mean 63.9 Range 49 Highest mean daily 76, 10th Lowest mean daily 52.5, 13th Mean maximum 76.1 Mean minimum 51.7 Greatest daily range 37, 16th Least daily range 6, 24th	87.7 33.2 61.7 54.5	97, 1895, 1929 24.5, 1914
Precipitation, in inches		
Precipitation 4.88	4.24	14, 55, 1938; .52, 1914
Snow Maximum precip, in 24 hrs. 1.85, 30th Number of days with .01 or more	10	16, 1933; 3, 1903
Wind, in miles		
Total movement2429Greatest daily movement173, 30thLeast daily movement23, 16thMean hourly velocity3.4Maximum velocity22, 30th	3271 4.5 25.9	4 686, 1896; 1,414, 1894 80, 1938
Wind, direction		
Prevailing direction W North, days 2 Northeast, days 3 East, days 3 Southeast, days 5 South, days 1 Southwest, days 5 West, days 4 Northwest, days 7	WSW	
Weather		
Mean relative humidity, percent 78.4 Mean cloudiness, percent 44 Number of clear days 18 Number of fair days 6 Number of cloudy days 6 Number of hours bright sunshine 242 Percent of possible hours of bright sunshine 64.9 Thunder and lightning 30th First frost	10 10 10 201 54.9	70, 1934; 27, 1908, 1914 19, 1932; 2, 1907, 1928 19, 1908; 3, 1889 20, 1934; 3, 1941 255, 1916; 106, 1934 Oct. 13; Aug. 22

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1945.

REMARKS

September weather was warmer than usual, with a little more sunshine, humidity and rainfall than normal.

The mean temperature for the month was 63.9 degrees compared to the normal for September of 61.7 degrees. The highest temperature was 86 degrees on the 19th and 20th. The lowest was 37 degrees on the 14th and consequently no frosts occurred in September this year. Also no frosts were recorded in September last year. The normal date for first frost at this station is September 21.

The domestic heating load for the month was 90 degree-days compared to the normal for September of 149 degree-days.

The total rainfall for the month was 4.88 inches, while the normal is 4.24 inches. Most of the precipitation fell in three storms: 1.21 inches on the 8th and 9th, 1.40 inches on the 24th, and 1.85 inches on the 30th. There were 242 hours of bright sunshine while the normal is 201 hours. The mean relative humidity was 78.4 percent, the normal for September being 73.6 percent.

Dr. J. K. Shaw, research pomologist, reports as follows: "There is nothing extraordinary to report about the weather and crops in September. Progress was normal. A good peach crop was harvested from our orchards and others in good locations. Our apple crop is poor in the orchards on lower elevations but good in the better located orchards. McIntosh suffered from the early freezes and probably cool weather following bloom interfered with the set of fruit, but not as much as it did in 1945. The unseasonably warm weather in March, followed by cold which killed many fruit buds, threatened ruin but still most orchards came through with fair to good crops. Injury was severe to total in orchards in frosty locations."

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 694

October 1946

Meteorological Observations

FOR

OCTOBER

1946

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, C7 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

	,	Гетре	rature	:		dity			Wind		Р	recipitati	on	
Day	Max	Time m	Degrees ui W	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 3 4 4 5 6 7 7 8 8 9 10 11 12 13 14 15 166 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31	50 48 69 77 81 84 84 84 59 66 60 75 61 66 69 71 61 65 66 67 67 67 67 67 67 67 67 67 67 67 67	1 P 2 P 4 P P 3 P P 4 P P 3 P P P 3 P P P P P	39 36 32 38 447 45 41 445 438 53 37 445 42 38 45 37 34 47 40 32 34 8 56 60 55	11 A 12 P 7 A 6 A 12 P 15 A 12 P 12 P 12 P 12 P 12 P 12 P 12 P 12 A 12 P 12 P 12 A 12 P 12 P 12 A 12 P 12 P 12 A 12 P 12 P 13 A 14 A 15 A 16 A 17 A 18 A 18 A 18 A 18 A 18 A 18 A 18 A 18	44.5 42.0 50.5 57.5 64.5 50.0 54.0 64.0 53.0 54.5 51.5 51.5 51.5 51.5 51.5 51.5 51.5	67 69 74 823 835 728 728 738 564 74 74 74 863 756 776 989 774 74 74 757	0.2 1.3 11.7 11.6 8.3 7.9 11.5 0.6 8.7 0.6 11.2 9.3 2.6 10.3 8.2 9.3 11.0 6.7 7.6 10.8 10.8 6.4 6.7 2.3 5.0 9.3 5.1	sw sw sw sw se ne sw w se nw se se se sw w mw se se nw sw se nw sw sw sw sw sw sw se ne sw sw sw se nw nw se nw nw se nw nw na nc na nc na nc nc nc nc nc nc nc nc nc nc nc nc nc	283 168 44 27 35 24 33 93 47 51 32 189 225 104 32 83 134 123 179 164 28 24 27 30 113 102 30 54 172 105 219	25 13 6 2 3 2 4 15 4 15 14 15 15 14 15 15 18 18 18 18 18 18 18 18 18 18 18 18 18	0 A 3 P 0 A 1 P	7:30 A 12 P 1 A 5 P	.11 TTTTTT T .555 .07 T .58	

^{*} Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

October, 1946	Normal	Extremes
Barometer, in inches (Readings reduced to sea level) Maximum 30.56, 28th Minimum 29.46, 18th Mcan semi-daily 30.114 Range 1.10	30.53 29.42 30.06 1.11	30.68, 1929 29.00, 1926 1.47, 1926, .76, 1899
Air Temperature, in degrees F.		
Highest 84, 6th, 7th Lowest 28, 22nd Mean 54,7 Range 56 Highest mean daily 70, 30th Lowest mean daily 41,5, 20th Mean maximum 67,4 Mean minimum 41,9 Greatest daily range 39, 4th, 7th, 23rd Least daily range 10, 26th	79.4 23.4 50.5 56.0	90.5, 1908 17, 1936 56.4, 1920; 43.2, 1890
Precipitation, in inches		
Precipitation1.51	3.29	8.81, 1911; .01, 1924
Snow — Maximum precip. in 24 hrs 0.62, 12th, 13th Number of days with .01 or more	9	15, 1913; 1, 1897, 1924
Wind, in miles		
Total movement . 2914 Greatest daily movement . 283, 1st Least daily movement . 24, 6th, 24th Mean hourly velocity . 3.9 Maximum velocity . 28, 19th, 31st	4074 54	5,467, 1910; 2,540, 1894
Wind, direction	29.5	42, 1937
Prevailing direction SW North, days 0 Northeast, days 1 East, days 0 Southeast, days 7 South, days 2 Southwest, days 10 West, days 6 Northwest, days 5	W	
Weather		
Mean relative humidity, percent Mean cloudiness, percent 49 Number of clear days Number of fair days Number of cloudy days Number of hours bright sunshine Percent of possible hours of bright sunshine Thunder and lightning First frost Cot. 22	69.0 48 10 10 11 175.0 51.3 Sept. 21	67, 1932; 18, 1924 21, 1938; 1, 1911 17, 1924; 3, 1938 19, 1896; 3, 1924 232, '23, '28, 91, 1913 Aug. 22, '94, '95; Oct. 15, '79

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1945

REMARKS

The weather during October was warm and dry, and had very little wind. The mean temperature was 54.7 degrees compared to a normal of 50.5 degrees. During the first half of the month the temperature was near normal but was well above normal in the second half. While the mean temperature was considerably above normal, this record was exceeded in 1900, 1913, 1920 and 1938. In 1920 the mean temperature for October was 56.4 degrees. The highest temperature in October this year was 84 degrees on the 6th and 7th. The lowest temperature was 28 degrees on the 22nd. This was the first frost of the season, although there was light frost earlier in nearby locations. At the end of the month there has not been a killing frost. The normal date for the first frost is September 21.

The domestic heating load for the month was 326 degree-days. The total heating load for September and October is 416 degree-days while the normal for this period is 605 degree-days.

The total precipitation for the month was 1.51 inches, which is 1.78 inches less than the normal for October.

The total wind movement was only 2914 miles for the month, while the normal is 4074 miles. Only twice since 1889 has there been less wind in October.

Dr. J. K. Shaw, research pomologist, gives the following final report on the crop season of 1946: "The crop season of 1946 may be said to end with October. In many ways it was remarkably like that of 1945. The weather was unseasonably warm in late winter and early spring, then cooler with killing temperatures followed by seasonable temperatures for the rest of the summer. Rainfall was low in March and early April both years and again in June and July, followed by more adequate amounts ending with dry Octobers. For three successive years the fruit crop has been injured more or less severely by spring killing of fruit buds. The writer remembers only one spring in the previous 38 years when severe killing occurred in Massachusetts. Three frosty years in succession sets a record that will not be equaled for a long time. If we get frost damage in 1947 it will be most remarkable.

"Apple scab caused much damage to scab susceptible varieties in both years because of wet weather in May and June. The carryover of the disease on fallen leaves is heavy. Unless we have a dry May and June in 1947, a most thorough and skillful control program will be necessary to control the disease."

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 695

November 1946

Meteorological Observations

FOR

NOVEMBER

1946

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

	,	Гетре	rature			ty	S		Wind		I	Precipitat	ion	
	Maxi	imum	Mini	imum		fumidi	f Hours nshine			S				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hour Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 29 20 20 20 20 20 20 20 20 20 20 20 20 20	64 54 56 68 49 52 59 50 54 54 54 58 59 49 45 50 45 50 45 50 45 50 45 49 49 49 49 49 49 49 49 49 49 49 49 49	3 P 4 P 7 1 P 2 P 7 3 P 1 2 P 7 3 P 1 2 P 7 3 P 1 2 P 7 3 P 1 P 2 P 7 3 P 1 P 2 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P 1	41 47 48 49 39 32 41 44 40 33 50 38 32 28 20 40 23 27 24 27 24 38 35 27 24 38 27 27 24 27 25 28 25 26 27 27 27 28 28 28 27 27 28 28 28 28 28 28 28 28 28 28 28 28 28	7 A 4 A 3 A 12 P 10 P 7 A 12 P 7 A 12 P 7 A 12 P 7 A 12 P 7 A 12 P 7 A 12 P 7 A 12 P 7 A 12 P 7 A 12 P 7 A 12 P	52.5 50.5 52.0 58.5 44.0 42.0 50.0 47.0 43.5 54.0 36.5 35.0 46.5 32.5 34.0 31.0 31.5 50.0 31.0 31.0 31.5 50.0 30.0	61 69 89 74 41 47 60 92 68 76 87 57 51 76 93 83 83 65 54 52 74	8.1 0.0 0.0 3.0 8.4 10.1 3.5 5.2 0.0 7.5 9.8 9.8 9.8 9.0 7.7 9.6 2.2 9.6 2.2 9.3 0.3 0.3 0.3 0.3 0.3 0.3 0.3 0	w e w w se se w w nw se	142 69 95 154 397 219 110 40 133 88 150 264 14 134 191 61 116 151 225 56 44 34 34 101 116 125 139 164 255 253 61	20 7 5 5 5 22 30 20 18 5 17 9 13 25 18 6 6 7 7 22 22 13 13 13 13 14 25 18 25 18 25 18 25 18 25 18 25 18 25 18 25 18 25 25 25 25 25 25 25 25 25 25	2:30 A 5 P 1 A 8:30 A	8:30 p 10 p 3 A	.34 T .10 .13	Т

^{*} Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

A-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1	T T	
November, 1946	Normal	Extremes
Barometer, in inches (Readings reduced to sea level) Maximum 30.66, 18th Minimum 29.60, 8th Mean semi-daily 30.081 Range 1.06	30.59 29.30 30.05 1.29	30.87, 1932 28.73, 1904 1.84, 1904; .89, 1943
Air Temperature, in degrees F.		
Highest 68, 4th Lowest 19, 19th Mean 43.2 Range 49 Highest mean daily 58.5, 4th Lowest mean daily 30.0, 30th Mean maximum 51.7 Mean minimum 34.7 Greatest daily range 30, 16th, 19th Least daily range 6, 8th	66.0 12.9 38.9 53.0	75, 1924, 1938 -4, 1938 44.1, 1931; 33.6, 1901
Precipitation, in inches		
Precipitation	3.41 2.34	8.64, 1927; .63, 1917 13.50, 1938
Number of days with .01 or more	9	22, 1921; 2, 1904
Wind, in miles		
Total movement 4358 Greatest daily movement 397, 5th Least daily movement 34, 21st Mean hourly velocity 6.1 Maximum velocity 30, 5th	4531 6.3 30.3	5,978, 1906; 2,589, 1889 44, 1938
Wind, direction	30.3	44, 1900
Prevailing direction WSW North, days 0 Northeast, days 0 East, days 3 Southeast, days 6 South, days 1 Southwest, days 3 West, days 10 Northwest, days 7	WNW	
Weather		
Mean relative humidity, percent 68.8 Mean cloudiness, percent 57 Number of clear days 11 Number of fair days 55 Number of cloudy days 14 Number of hours bright sunshine 126 Percent of possible hours of bright sunshine 43 Thunder and lightning — First frost 13th	70.6 55.1 6 9 15 121 41.3 Nov. 6	72, 1900, 1927; 34, 1917 15, 1903; 1, 1900, 1911, 1927 16, '12; 4. '89, '30, '32, '35 24,1927; 9, 1905, 1917 182, 1903; 66, 1927 Oct. 10, '25; Dec. 13, 1941

Note—The first **column** in the above summary gives observations made during the month. The **second column** gives the averages based on observations made from 1889 to 1938, **except that** humidity records are based on observations made from 1929 to 1938. The **third column** gives extremes observed from 1889 to 1945

REMARKS

The weather during November was warm and dry. The mean temperature for the month was 43.2 degrees while the normal mean temperature for November is 38.9 degrees. This high mean temperature for November has been exceeded only three times since 1889, in 1927, 1931 and 1935. The highest temperature was 68 degrees on the 4th and the lowest 19 degrees on the 19th. The first frost of the season was on the 13th. The domestic heating load was 654 degree-days compared to a normal of 784 degree-days. The total heating load to the end of November is 1070 degree-days while the normal for that period is 1389 degree-days.

The total precipitation during the month was .70 inches. Only once has there been less precipitation in November since 1889. This was in 1917 when the total precipitation for November was .63 inches. The normal for November is 3.41 inches. No snow fell during the month whereas the normal snowfall for November is 2.34 inches. The normal date for the first snow is November 6.

There were 126 hours of bright sunshine, the normal being 121 hours.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series Bulletin No. 696

December 1946

Meteorological Observations

FOR

DECEMBER

1946

C. I. GUNNESS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft Height of wind instruments, 67 feet. Time used, E. S. T.

> Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Tempe	rature			ity	ırs		Wind			Precipita	tion	
	Max	imum	Mini	imum		Humid	of Hours	bo	<u> </u>	s e				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hour Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 28 29 30 31 31 31 31 31 31 31 31 31 31 31 31 31	53 34 35 39 40 40 48 45 56 52 57 30 33 32 29 35 37 37 35 35 37 37 32 32 32 32 32 32 32 32 32 32 32 32 32	11 A A B A B A A B A A B A A B A B A A B A B A A B B A B	33 12 11 23 27 26 37 41 26 23 27 22 20 16 17 20 30 31 31 27 22 20 16 17 20 30 7 21 21 21 21 21 21 21 21 21 21 21 21 21	0 A P 12 P 12 P 12 P 12 P 12 P 12 P 12 P	43.0 23.0 23.0 31.0 33.5 33.0 52.0 52.0 42.0 26.5 22.5 27.5 29.0 22.5 22.5 22.5 21.0 15.0 18.0 24.0 24.0 24.0 20.5 13.5	73 50 48 57 68 74 84 86 73 47 85 55 42 45 86 73 84 67 65 65 65 67 88	0.0 6.0 9.3 2.0 0.1 6.0 0.0 1.3 0.0 5.8 2.5 4.7 9.0 0.0 0.0 1.2 4.8 6.4 6.0 0.0 9.2 1.9	w nw nw se s w se w nw nw nw nw nw nw nw nw nw nw nw nw n	193 435 126 142 75 84 61 39 58 133 289 242 298 242 298 32 158 340 209 167 258 312 100 140 87 257 18	22 32 10 11 7 7 7 7 8 13 28 28 28 28 20 18 8 11 7 7 32 20 12 33 21 12 32 32 4	11 A 7:30 A 2 A 7 P 0 A	2:30 P 8 A 1 P 12 P 4 P 12 P 4:30 P	.16 .03 .11 .50 1.39	1.0 5.0 0.5 0.5 T

^{*} Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

December, 1946	Normal	Extremes
Barometer, in inches (Readings reduced to sea level) Maximum 30.59, 31st Minimum 29.30, 13th Mean semi-daily 30.033 Range 1.29	29.27 30.06	30.96, 1889 28.85, 1915 1.78, 1895; 1.01, 1892
Air Temperature, in degrees F.		
Highest 63, 10th Lowest 1, 31st Mean 29.9 Range 62 Highest mean daily 52, 10th Lowest mean daily 13.5, 31st Mean maximum 39.0 Mean minimum 20.8 Greatest daily range 30, 11th, 13th Least daily range 6, 21st, 28th	54.3 -1.9 27.5 56.2	65.5, 1908 -22.5, 1917 36.9, 1891; 17.1, 1917
Precipitation, in inches		
Precipitation 3.51 Snow 10.50 Maximum precip. in 24 hrs. 1.89, 20th, 21st	3.39 8.50	7.77, 1901; .58, 1943 26.50, '02; t, 1891, 1943
Number of days with .01 or more8	10	17, 1902; 4, 1892, 1943
Wind, in miles	4740	((01,100), 0.300,1010
Total movement 4937 Greatest daily movement 435, 2nd Least daily movement 18, 31st Mean hourly velocity 6.6 Maximum velocity 35, 26th	6.3 31.4	6,694, 1925; 3,239, 1918 48, 1938, 1943, 1944
Wind, direction		
Prevailing direction WNW North, days		
Weather		
Mean relative humidity, percent 67.8 Mean cloudiness, percent	69.4 54.9 8 9 14 128	71, 1929; 39, 1919 15, 1890; 2, 1922, 1933 16, '09; 4, '89, '30, '31, '36, '38 23, '33; 7, '09, '23 172, 1896; 63, 1933
Thunder and lightning — — — — — — — — — — — — — — — — — — —	45.2	

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1945.

ANNUAL SUMMARY

Annual, 1946	Normal	Extremes
Barometer, in inches (Readings reduced to sea level) Maximum 30.70, Jan. 20 Minimum 29.23, Feb. 14 Mean semi-daily 30.046 Range 1.47	30.81 28.95 30.01 1.85	31.05, 1920, 1943 28.41, 1938 2.47, 1938; 1.38, 1933
Air Temperature, in degrees F.		
Highest 95, July 19 Lowest -8, Jan. 28, Feb. 4 Mean 48.7 Range 103 Highest mean daily 81.5, July 20 Lowest mean daily .6.0, Jan. 20	95.7 -12.2 47.4 107.8	104, 1911 -26, 1904 49.8, 1921; 44, 1904
Mean maximum 59.7 Mean minimum 37.6 Greatest daily range 43, Apr. 23 Least daily range 4, Mar. 2	57.8 36.5	
Precipitation, in inches		
Precipitation 38.61 Snow 50.60 Maximum precip. in 24 hrs. 3.46, July 23rd	43.70 47.78	59.00, 1938; 30.68, 1908 89.00, 1893; 24.50, 1919
Number of days with .01 or more111	124	146, 1945; 96, 1924
Wind, in miles		
Total movement 48,381 Greatest daily movement 499, Feb. 21 Least daily movement 9, Jan. 17 Mean hourly velocity 5.5 Maximum velocity 50. Feb. 20	52,223 5.8 39.5	63.571, 1908; 36,257, 1894
Wind, direction		
Prevailing direction WSW North, days 23 Northeast, days 7 East, days 12 Southeast, days 81 South, days 32 Southwest, days 39 West, days 78 Northwest, days 93	W	
Weather		
Mean relative humidity, percent 68.1 Mean cloudiness, percent 51.6 Number of clear days 166 Number of fair days 97 Number of cloudy days 102 Number of hours bright sunshine 2642 Percent of possible hours of bright sunshine 59.3 Last snow Apr. 8 First snow Dec. 17 Last frost May 4 First frost Coc. 22	May 14	60, '98, '01, '02; 41, '08, '24 217, 1941; 59, 1927 182, 1912; 64, 1936 179, '01, '02; 71, 1910, 1941 3038, 1941; 1864, 1902 Mar. 14,'10; May 11,'07,'45 Oct. 10, '25; Dec. 13, '41 Apr. 14, '43; June 8, '32 Aug. 22,'94,'95; Oct. 13,'09

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 697

January, 1947

Meteorological Observations

FOR

JANUARY

1947

M. J. MARKUSON

OBSERVATORY

Latitude, 42° 23' 31'' N. Longitude, 72° 31' 48'' W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Temperature 2							Wind			Precipitation				
	Maximum		aximum Minimum		laximum Minimum			Humid	r of Hours Sunshine	bo .	<u>+</u>	*es				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number Bright St	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow		
1 2 3 4 5 6 7 8	19 30 34 32	4 P 12 P 9 P 0 A	10 15 30 14	8 A 0 A 0 A 12 P	14.5 22.5 32.0 23.0	73 94 98 73	0.2 0.0 0.0 9.2	n n n	70 237 145 120	25 28 4 13	7:30 P 0 A 8:30 A	12 P 2 P 6 P	.65 .24 .17	2.5 0.5		
6	32 31 37	1 P 3 P 8 P	8 10 5	7 а 12 р 5 а	20.0 20.5 21.0	69 64 74	5.0 7.0	se nw	62 159 97	8 14 12				T		
8	37 24	8 P 1 A 0 A	24 7	5 A 12 P 12 P	30.5 15.5	64 51	1.6 7.6 9.3	se w	283 235	25 28			Т	T		
10 11 12 13 14	26 42 43 25 32	1 P 3 P 4 A 3 P	-1 19 16 9 24	7 A 0 A 12 P 7 A	12.5 30.5 29.5 17.0	67 65 45 57	5.0 0.2 7.0 5.3	w s s n n	118 184 221 109	18 13 22 11	8:30 A	12 в	.14			
15 16 17 18	36 38 40 42	2 P 4 P 3 P 1 P	32 34 27 23	0 A 9 A 8 A 12 P	28.0 34.0 36.0 33.5 32.5	91 97 97 50 48	0.0 0.0 0.0 5.6 0.0	n nw nw nw w	50 7 40 181 206	6 4 8 28 20	8:30 A 2 A	8 P	.31			
19 20 21 22 23	29 45 45 17 24	6 P 12 P 0 A 0 A	10 25 17 9	8 A 0 A 12 P 9 A 8 A	19.5 35.0 31.0 13.0 17.5	63 97 66 52 54	4.0 0.0 4.0 8.6 2.8	nw nw w	60 35 261 332 83	13 5 32 28 8	5:30 A	бр	.84 T	Т		
23 24	45 45	12 P 2 P 11 A	24 26	0 A 11 P	34.5 35.5	67	3.5	se	130 133	9		i	Т	T		
25 26 27 28	38 47 53	12 P 3 P 3 P	20 28 26	8 A 12 P 7 A	29.0 37.5 39.5	81 87 67	2.2 6.0 9.1	n nw n nw	11 65 57	2 7 17	11:30 p 0 a	12 р 7:30 л	.01 .44			
29 30 31	35 44 55	0 A 12 P 1 P	28 30 34	5 A 0 A 12 P	31.5 37.0 44.5	70 92 69	0.0 0.0 5.9	n n nw	50 28 105	10 5 25	6 р 0 а	12 p 6 a	.08 .49			

^{*} Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

January, 1947	Normal	Extremes
Barometer, in inches (Readings reduced to sea level) Maximum 30.89, 1st Minimum 29.19, 20th Mean semi-daily 30.037 Range 1.70	30.70 29.20 30.07 1.50	31.00, 1927 28.55, 1913 2.18, 1913; 0.97, 1896
Air Temperature, in degrees F.		
Highest 55, 31st Lowest -1, 10th Mean 27.7 Range 56 Highest mean daily 44.5, 31st Lowest mean daily 12.5, 10th Mean maximum 36.2 Mean minimum 19.2 Greatest daily range 32, 7th Least daily range 4, 3rd, 15th, 16th	51.1 -7.1 24.2 58.2	66.0, 1932 -26.0, 1904 34.2, 1913; 13.9, 1918
Precipitation, in inches		
Precipitation 3.37 Snow 3.0 Maximum precip. in 24 hrs. 0.84, 20th	3.61 13.32	7.15, 1898; 1.07, 1896 33, 1898, 1923; 1.50, '08
Number of days with .01 or more 10	11	15, '93, '20; 4, 1901
Wind, in miles		
Total movement3874Greatest daily movement332, 22ndLeast daily movement7, 15thMean hourly velocity5.2Maximum velocity32, 21st	5055 6.8 32.7	7770, 1908; 2896, 1895 47, 1938
Wind, direction		, 2200
Prevailing direction NW North, days 10 Northeast, days 0 East, days 0 Southeast, days 3 South, days 3 Southwest, days 0 West, days 5 Northwest, days 10	WNW	
Weather		
Mean relative humidity, percent 71.3 Mean cloudiness, percent 56 Number of clear days 7 Number of fair days 9 Number of cloudy days 15 Number of hours bright sunshine 112 Percent of possible hours of bright sunshine 38.1 Thunder and lightning	70.1 55 9 9 13 137 46.7	78, 1932; 37, 1908 20, 1944; 2, 1914 18, 1890, 1926; 3, 1916, 1944 22, 1923, 1931; 4, 1920, 1939 214, 1920; 74, 1932

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1946.

REMARKS

The weather during January was much warmer than normal with very little snow. The mean temperature was 27.7 degrees, which is 3.5 degrees higher than the normal for January. This was the warmest January since 1937 which had a mean temperature of 31.7 degrees. Last year the mean temperature for January was 23.2 degrees. Thus far, this has been the warmest winter since 1931-1932. The highest temperature this January was 55 degrees on the 31st while the lowest was -1 degree on the 10th and this is the lowest reading so far this winter.

The domestic heating load for January was 1157 degree-days as compared to the normal of 1265 degree-days. The total heating load from September 1 to January 31 is now 3315 degree-days as compared to the normal of 3817 degree-days for this same period.

The snowfall for January was 3.0 inches while the normal for January is 13.32 inches. The total snowfall for this winter has now reached 13.5 inches. The normal for this same period is 24.16 inches. The total precipitation for January was 3.37 inches which is slightly under the normal of 3.61 inches. There were 112 hours of bright sunshine as compared to a normal of 137 hours. High winds occurred on the 21st when a maximum of 32 miles per hour was reached.

MABS. STATE COLLEGE 3 46 AMHERST, MASS.

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series Bulletin No. 698

February, 1947

Meteorological Observations

FOR

FEBRUARY

1947

M. J. MARKUSON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft Height of wind instruments, 67 feet. Time used, E. S. T.

> Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Tempe	rature			ity	ırs e		Wind		I	Precipita	tion			
	Maximum		Minimum		Minimum			Humid	of Ho anshin	ρū		s.				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max, Miles per Hour*	Began	Ended	Water	Snow		
1 2 3 4 5 6 7 8 9 10 11 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 28 29 29 20 20 20 21 21 22 23 24 24 25 26 27 27 28 28 28 28 28 28 28 28 28 28 28 28 28	28- 37 50 19 19 28 22 33 36 48 40 43 35 48 27 21 21 29 33 37 35 31 32 36 37 37 37 37 37 37 37 37 37 37 37 37 37	3 P 7 P 1 A A 3 P 12M 12M 2 P 3 P 4 P 1 A P 1 A P 2 P 2 P 2 P 3 P 2 P 3 P 2 P 3 P 2 P 3 P 3	22. 19 13 19 10 6 6 15 10 4 19 18 11 21 31 26 19 12 4 8 8 11 5 19 19 19 19 19 19 19 19 19 19 19 19 19	12 P 17 A A 12 P 12 P 12 P 7 A A 12 P 13 A 0 A 17 A 12 P 17 A 12 P 17 A 12 P 12 P 17 A 12 P 17 A 12 P 17 A 12 P 17 A 18 A 18 A 18 A 18 A 18 A 18 A 18 A 18	28.5 23.5 25.0 34.5 14.5 12.5 22.5 22.5 34.5 37.0 30.5 33.5 19.5 14.5 20.0 19.0 28.0 28.0 28.0 26.0	50 555 488 564 776 564 5776 561 767 768 769 769 769 769 769 769 769 769 769 769	10.0 5.8 9.0 0.5 6.1 2.7 1.6 0.0 4.6 7.3 10.0 10.4 4.7 6.7 0.0 0.8 2.1 10.5 10.7 6.5 5.6 7.0 1.8 9.6 7.0 1.8 1.8 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9	nw w s se nw nw nw nw nw nw nw n nw s s nw nw nw nw nw nw nw nw nw nw nw nw nw	236 153 143 124 293 135 169 239 373 412 157 49 86 65 209 2435 109 260 201 327 306 169 139 27 298 294 262	20 18 15 25 22 13 5 17 32 32 25 15 8 9 7 7 25 28 11 28 18 12 20 25 28 18 19 20 20 20 20 20 20 20 20 20 20 20 20 20	1 P 6 A 4 A 4 A 4 A A 4 A A A A A A A A A A	10 P 10 A 3 P 5 P	.04 .59	1.5 1.0 7.0 1.0 7.5 T		

^{*} Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

February, 1947	Normal	Extremes
Barometer, in inches (Readings reduced to sea level) Maximum 30.44, 12th	30.63	31.05, 1920
Maximum 30.44, 12th Minimum 29.15, 22nd Mean semi-daily 29.729 Range 1.29	29,24 30,04 1.38	28.56, 1895 1.89, '00, '08; .88, '13, '31
Air Temperature, in degrees F.		
Highest 50, 4th Lowest 4, 9th, 20th Mean 24.0 Range 36 Highest mean daily 37, 16th Lowest mean daily 12.5, 21st	50.5 -7.4 23.7 58	65.0, 1930 -23, 1943
Mean maximum 32.1 Mean minimum 15.8 Greatest daily range 31, 4th Least daily range 6, 27th	32.8 14.5	
Precipitation, in inches		
Precipitation	3.19 13.99	8.12, 1900; 0.62, 1901 48.75, 1893; 0.50, 1937
Number of days with .01 or more6	10	15, '93, '20; 4, 1901
Wind, in miles		
Total movement 5862 Greatest daily movement 435, 17th Least daily movement 15, 7th Mean hourly velocity 8.7 Maximum velocity 32, 9th, 10th	4776 7.1	6445, 1896; 3438, 1892
	31.7	50, 1946
Wind, direction	******	
Prevailing direction WNW North, days 2 Northeast, days 0 East, days 0 Southeast, days 1 South, days 4 Southwest, days 1 West, days 3 Northwest, days 17	WNW	
Weather		
Mean relative humidity, percent 58.9 Mean cloudiness, percent 58 Number of clear days .8 Number of fair days .9 Number of cloudy days .11 Number of hours of bright sunshine Percent of possible hours of bright sunshine Thunder and lightning 48.6 4th	50.9 10 8	66, '90, '27; 31, 1905 19, 1941; 2, 1927 16, 1920; 2, 1936 16, 1894; 2, 1920 221, 1924; 110, '27, '38

The weather for February was near normal. The mean temperature for the month was 24.0 degrees as compared to the normal of 23.7 degrees. The highest temperature during the month was 50 degrees on the 4th and the lowest, 4 degrees on the 9th and 20th. The lowest temperature for the winter still stands at -1 degree on Jaunary 10. The domestic heating load for the month was 1148 degree-days, while the normal for February is 1165 degree-days. The total heating load for the period September 1 to February 28 inclusive is now 4463 degree-days compared to the normal of 4982 degree-days for this period.

A total of 18 inches of snow was recorded during the month, with 16.5 inches of this amount falling in two storms. The total snowfall for this winter season is now 31.50 inches, while the normal for the period is 38.15 inches.

There were only six days of precipitation during the month. The total hours of sunshine was 144, while the normal is 158 hours. The total wind movement for the month was higher than normal.

MASS. STATE COLLEGE AMHERST, MASS. 3 46

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 699

March 1947

Meteorological Observations

FOR

MARCH

1947

M. J. MARKUSON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

	7	rempe	rature			ity	ırs e		Wind F			Precipita	Precipitation		
	Maxi	mum	Mini	mum		Humid	of Hou	bs: _	1	s *					
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow	
1 22 33 44 55 66 77 88 9 10 11 122 133 144 115 116 117 118 119 220 221 222 233 224 225 227 227 227 227 227 227 227 227 227	37 37 37 32 38 42 40 39 41 44 43 50 53 53 39 36 34 45 50 50 53 53 53 53 53 53 53 53 53 53 53 53 54 55 55 55 56 57 57 57 57 57 57 57 57 57 57 57 57 57	3 P P 1 A A 1 P P 1 P P 1 P P 1 P P 1 P P 1 P P 1 P P 1 P P 1 P P 1 P P 1 P	11 28 28 24 25 32 28 30 27 28 31 29 23 36 31 24 26 22 19 21 31 29 26 37 27 18 19 19	7 A A 12 P 7 A A 7 A A 7 A A 7 A A 6 A A 12 P 7 A A 6 A A 12 P 7 A A 6 A A 12 P 7 A A 6 A A 12 P 7 A A 6 A A 6 A A 1 A A 12 P 7 A A 6 A A 6 A A 12 P 7 A A 6 A A 12 P 7 A A 6 A A 12 P 7 A A 6 A A 13 P 7 A A 6 A A 14 P 7 A A 6 A A 15 A A 15 A A 16 A A 17 P 8 A 17 P 8 A 18 P	24.0 32.5 32.5 28.0 31.5 37.0 34.0 29.5 34.0 39.5 38.0 44.5 42.0 31.5 31.0 28.0 32.0 35.5 38.5 40.0 32.0 35.5 38.5 40.0 32.0 35.0 36.0 37.0 38.5 38.5 38.6 38.6 38.6 38.6 38.6 38.6 38.6 38.6	74 90 73 68 63 63 61 63 56 65 79 61 64 93 54 53 54 63 64 77 48 48	8.4 0.0 0.0 1.6 7.8 6.2 9.1 11.8 0.0 10.8 10.9 1.1 9.0 11.8 8.0 10.9 1.1 9.0 11.8 0.0 10.9 10.0 10	w nw sw w n n n n n s s n w nw w sw sw sw n n n n	55 185 368 211 243 205 176 100 218 215 126 70 111 174 240 218 139 242 157 98 140 94 115 61 1299 458 324	6 30 45 13 25 20 15 13 18 14 10 10 10 15 20 20 20 20 22 20 21 31 31 31 31 31 31 31 31 31 31 31 31 31	0 A 0 A 8 P 0 A 9 A 7 A	12 P 8 A	1.72 .14 .19 .95	3.00 T T T	
28 29 30 31	37 45 35 45	3 P 11 A 3 P 4 P	15 23 27 19	7 A 5 A 12 P 6 A	26.0 34.0 31.0 32.0	59 65 51 46	12.3 2.2 8.0 12.6	se w nw	153 155 332 241	20 20 32 18				Т	

^{*} Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

	1	1
March, 1947	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)	24.55	21.05
Maximum	30.57 29.24	31.05, 1943 28.47, 1914
Minimum 28.63, 3rd Mean semi-daily 29.840 Range 2.13	30.00 1.34	
•	1.54	2.10, 1914; 0.85, 1915
Air Temperature, in degrees F.		
Highest	63.8 6.1	85, 1929 -7.5, 1906
Lowest	34.4	7.07, 1.77
Range 44	57.6	
Range 44 Highest mean daily 46, 24th Lowest mean daily 22.5, 26th		
Mean maximum 42.61	43.2	
Mean minimum 25.3	25.4	
Mean minimum 25,3 Greatest daily range 30, 13th Least daily range 8, 4th, 30th		
Precipitation, in inches		
	2.70	7.90 1012 . 0.12 1017
Precipitation 3.29 Snow 3.00 Maximum precip. in 24 hrs. 1.72, 2nd	3.70 7.47	7.89, 1942; 0.12, 1915 27, 1899; 0, 1921
Maximum precip. in 24 hrs		
Number of days with .01 or more6	11	17, 1890, 1913, 1936 : 3, 1915
Wind, in miles		
Total movement 5923 Greatest daily movement 458, 26th Least daily movement 55, 1st Mean hourly velocity 7.9 Maximum velocity 45, 3rd	5944	8182, 1896; 3006, 1905
Maximum velocity	7.7 31.2	48, 1932, 1939, 1941, 1942
Wind, direction	0=	
Prevailing direction WNW North, days 7 Northeast, days 0 East. days 0 Southeast, days 2 South, days 3	WNW	
South, days 3 Southwest, days 3 West, days 7 Northwest, days 9		
Weather		
Mean relative humidity, percent 62.9 Mean cloudiness, percent 54 Number of clear days 13 Number of fair days 9 Number of cloudy days 9 Number of hours bright sunshine 199 Percent of possible hours of bright	64.2 51.8 11 10 10 199	68, 1901; 27, 1915 22, 1924; 3, 1901 17, 1900; 1, 1943 21, 1901; 1, 1915 292, 1924; 93, 1901
sunshine 53.6 Thunder and lightning -	53.6	

The weather during March was a little colder than normal with mean temperature of 34° compared to the normal mean of 34.4° . Last year we had the warmest March on record with mean temperature of 44.5° .

The mean temperature for the winter season now stands at 31.8° compared to the normal for the same period of 29.7° . Highest temperature of 55° this month occurred on the 24th; the normal high is 63.8° . The lowest temperature of 11° came on the 1st; the normal low is 6.1° . Record high was 85° on the 29th in 1945° and the record low is -7.5° as of March 24, 1906.

Degree-days for March were 962 against a normal of 950. The heating period, September 1 to March 31, is 5425 degree-days which is somewhat below the normal of 5932 degree-days.

There was less snow than usual when only 3 inches fell. The normal is 7.47 inches. Total snow of 34.5 inches this winter is considerably below normal of 45.62 inches. The normal for the entire snow season is 47.79 inches.

Precipitation is less than normal for March when 3.29 inches fell. The March normal is 3.70 inches.

There were 199 hours of bright sunshine which was the same as the normal.

Humidity was 62.9 percent; normal is 64.2 percent.

Two very low barometric readings developed in March when 28.63 inches occurred at 6 A.M. on the 3rd and 28.88 inches at 1 P.M. on the 25th. The hurricane of 1938 had a low barometer recording of 28.41 inches. Both lows this month were accompanied by high winds. Maximum velocity of 45 miles per hour was recorded on the 3rd.

Alle El, LACO.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series Bulletin No. 700

April 1947

Meteorological Observations

FOR

APRIL

1947

M. I. MARKUSON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

> Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

The large decision The lar			Тетре	rature	:		ity	ırs e	Wind Precipitation						
1	ay					ean	ean elative Humid	umber of Hours ight Sunshine	evailing	ovement	ax. Miles r Hour*	gan	nded	ater	Snow
2	Q	ğ	E	ă	Ë	M	Σĕ	z g	45	I Z	M	Be	편	3	Sn
28 50 4 P 28 5 A 39.0 38 13.9 W 265 32 1.50 P 7 111 28 50 4 P 28 5 A 39.0 38 13.9 W 245 28 29 63 3 P 33 5 A 48.0 48 5.5 se 227 18 30 74 2 P 49 0 A 61.5 77 5.4 se 249 15 6 A 12 P 77	2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 20 21 22 23 24 25 26 27 28	46 51 54 40 74 55 54 46 63 73 73 50 62 45 48 56 51 56 8 51 56 63 50	12 M 4 P 4 P 6 A 7 P 7 P 7 P 7 P 7 P 7 P 7 P 7 P 7 P 7	35 32 23 35 37 37 33 34 448 37 35 36 33 29 35 31 28 410 38 40 38 36 36 36 36 36 36 36 36 36 36 36 36 36	10 P 7 A 7 A 6 A 12 P 6 A 12 P 6 A 12 P 6 A 12 P 6 A 12 P 8 A 12 P 8 A 12 P 8 A 12 P 8 A 12 P 8 A 12 P	40.5 41.5 38.5 55.5 46.0 43.5 40.5 40.5 40.5 40.5 40.5 42.0 33.0 33.0 33.0 45.5 47.0 45.5 47.0 45.5 47.0 45.5 47.0 45.5 47.0 45.5 47.0 45.5 47.0 47.0 47.0 47.0 47.0 47.0 47.0 47.0	90 39 47 76 77 60 47 75 68 52 76 55 55 55 55 55 64 62 53 64 53 64 53 64 53 64 53 64 53 64 53 64 53 64 53 64 53 64 53 54 54 54 54 54 54 54 54 54 54 54 54 54	3.2 0.0 12.8 12.8 0.0 6.0 0.7 13.0 6.7 13.1 7.0 6.7 13.3 0.0 0.0 4.7 8.4 0.0 0.0 13.6 0.0 0.0 13.7 6.0 0.0 13.7 6.0 0.0 13.7 0.0 0.0 13.7 0.0 0.0 13.7 0.0 0.0 13.7 0.0 0.0 13.7 0.0 0.0 13.7 0.0 0.0 13.7 0.0 0.0 13.7 0.0 0.0 13.7 0.0 0.0 13.7 0.0 0.0 13.7 0.0 0.0 13.7 0.0 0.0 13.7 0.0 0.0 13.7 0.0 0.0 13.7 0.0 0.0 0.0 13.7 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0	s nw nw w s w nw n n sw sw w nw e w e w sw n n n se sw sw w w w w w w w w w w w w w w w	129 148 265 124 153 135 355 193 83 125 353 220 209 157 162 153 227 207 163 88 156 134 245	18 13 25 10 10 18 25 13 12 17 22 18 14 28 18 7 13 12 13 28 17 17 32 28	3 A 10 A 6 A 7 P 10:30A 6 A 0 A	8 P 10:30A 7 A 8:30 P 8:30 P 12 P 11 A	1.23 T .04 .03 .57 .52 .41.	1.25

^{*} Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

	1	1
April, 1947	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.51, 22nd Minimum 29.51, 27th Mean semi-daily 30.100	30.48 29.38 29.99	30.71, 1911 28.98, 1943
Range	1.11	1.52, 1930; .72, 1919
Air Temperature, in degrees F.		
Highest 74, 6th, 30th Lowest 23, 4th Mean 44.7 Range 51 Highest mean daily 61.5, 30th	79.4 22.0 45.7 57.4	90, 1941 8.5, 1923 52, 1921; 41, 1943
Mean maximum 31.3 Greatest daily range 37, 6th Least daily range 4, 20th	56.8 34.8	
Precipitation, in inches		
Precipitation 4.59 Snow 2.25 Maximum precip. in 24 hrs. 1.23, 5th	3.35 2.17	6.89, 1929; .55, 1941 11, 1891; 0, 1910, 1934
Number of days with .01 or more 10	11	18, 1909; 3, 1892
Wind, in miles		
Total movement5714Greatest daily movement355, 7thLeast daily movement83, 9thMean hourly velocity7.9Maximum velocity32, 27th	5404 7.5	8208, 1908; 3853, 1917
	31.4	40, 1935, 1938, 1940. 1945
Wind, direction Prevailing direction W North, days 6 Northeast, days 0 East, days 2 Southeast, days 3 South, days 2 Southwest, days 6 West, days 7 Northwest, days 4	WNW	
Weather		
Mean relative lumidity, percent Mean cloudiness, percent Number of clear days Number of fair days Number of cloudy days Number of thours bright sunshine Percent of possible hours of bright	61.6 51.8 11 9 10 220	75, 1901; 34, 1925, 1927 23, 1941; 3, 1898, 1902 18, 1915; 2, 1894, 1901 22, 1901; 1, 1941 329, 1941; 103, 1901
sunshine 45.5 Thunder and lightning 30th	54.7 Apr. 15	Mar. 14, 1910; May 11, 1907

The weather during April had about an inch more precipitation and a slightly lower mean temperature than normal. Precipitation totalled 4.59 inches which was fairly well distributed over 10 days. The normal precipitation is 3.35 inches. Mean temperature was 44.7 degrees while the normal mean temperature is 45.7 degrees. The highest temperature of 74 degrees occurred on the 6th and 30th; the normal is 79.4 degrees. The lowest temperature of 23 degrees was recorded on the 4th; the normal low is 22.0 degrees.

The snowfall totalled 2.25 inches compared to a normal of 2.17 inches. The total for the season is now 36.75 inches which is below the normal of 47.79 inches. The total hours of sunshine is 183 and this is somewhat below the normal of 220 hours for April.

The domestic heating load for the month was 608 degree-days which is above the normal of 579 degree-days. The season's total from September 1 to the end of April is now 6033 degree-days which is considerably less than the normal of 6511 degree-days.

Dr. J. K. Shaw, research pomologist, gives the following report: "The past winter was mild with only one sub-zero temperature recorded. Peach fruit buds came through with little killing and a full bloom is in prospect. There should be little or no winter injury to fruit plants. The progress of vegetation to May 1st is about normal and from two to four weeks later than it was in the last two years. Fruit buds are swelling and a few days of warm weather will bring them into bloom. There will be ample bloom on tree fruits but probably not equal to the heavy bloom of last year. Some leaf area of apple trees is exposed and the wet period at the beginning of May is favorable for infection from the heavy carryover of apple scab from last season. Thus the stage is set for another season of severe injury unless adequate preventive measures are taken."

AMHERST, MASS. 3 46

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 701

May 1947

Meteorological Observations

FOR

MAY

1947

F. J. SIEVERS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Ter	npera	iture		lity	L S		Wind		Pı	ecipitation	
	Max	imum	Min	imum		fumid	d Houn		-	S.			
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water
1 2 3 4 5	58 43 48 55 61	0 A 4 P 3 P 7 P 2 P 3 P	40 38 43 45 47	12 P 7 A 0 A 4 A 6 A	49.0 40.5 45.5 50.0 54.0	91 90 86 94 85	0.0 0.0 0.0 0.0 3.0	n ne n n	115 123 134 120 86	14 13 9 13 10	1:30 P	9:30 P 3 P 11 P 5 P 10:30 P	1.00 1.00 .14 .26
6 7 8 9	67 57 46 48	10 A 2 P 5 P	47 40 34 34	7 A 12 P 12 P 4 A	57.0 48.5 40.0 41.0	68 72 50 46	5.5 1.0 8.0 5.0	nw s nw nw	271 107 349 208	25 20 28 29	12:30 р	4 P	.16
10 11 12 13 14 15 16	60 71 81 87 62 65 66 73	5 P 5 P 1 P 0 A 5 P 1 P	28 32 49 54 37 31 38 42	5 A 6 A 0 A 4 A 12 P 5 A 5 A	44.0 51.5 65.0 70.5 49.5 48.0 52.0 57.5	38 35 37 52 64 48 55 48	14.4 14.4 14.5 7.0 3.0 14.6 7.0 11.5	w w sw sw e se se	127 127 160 143 185 105 179 110	12.5 12.0 20 25 12.5 10.5 15 20	10:30 р	12 p	.06
18 19	58 76	8 P	49 54	6 A	53.5	93	0.0	n s	98	10	1 A	6 P	.27
20 21 22 23 24 25	82 61 64 80 87	4 P 4 P 0 A 7 A 5 P 3 P	53 50 49 45 53	2 A 5 A 10 P 12 P 5 A 5 A	65.0 67.5 55.5 56.5 62.5 70.0	66 78 81 88 60 66	9.0 5.7 0.8 0.0 14.5	s s n n	185 129 149 69 102	10 28 12 18 11 16	12:30 A 9 A 9 A	4 A 12 M 10 A	.04 .20 .10
26 27	77 76 71	12 м 5 р 5 р	62 57 49	6 A 7 A 12 P	69.5 66.5 60.0	71 49 38	11.5 7.5 14.9 15.0	e nw nw	108 223 195	17 14 17	2 P 0 A	12 P 12:30 A	2.10
28 29 30 31	81 86 66 68	3 P 3 P 3 P 3 P	48 63 46 37	1 A 5 A 12 P 5 A	64.5 74.5 56.0 52.5	62 75 44 43	7.4 8.6 10.3 10.9	se se nw nw	90 198 259 191	11 22 25 22	8 P	11:30 р	.15

^{*} Based on least time required to blow one mile.

FERNAND E. BARTLETT. Observer

May, 1947	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30,32,10th Minimum 29,50, 6th Mean semi-daily 29,968 Range 0,82	30,39 29,51 29,96 0,88	30.62, 1936 29.10, 1938
Air Temperature, in degrees F.		
Highest 87, 13th, 24th Lowest 28, 10th Mean 56, 1 Range 59 Highest mean daily 74, 5, 29th Lowest mean daily 40, 0 Mean maximum 67, 1 Mean minimum 45, 0 Greatest daily range 39, 11th Least daily range 5, 2nd, 3rd	86.1 31.2 57.1 54.9	94.5, 1896, 1911 24.0 1900
Precipitation, in inches		
Precipitation	3.60	7.44, 1931; .48 1903
Snow	12	20, 1901, 1945; 5, 1903
Wind, in miles		
Total movement4762Greatest daily n.ovement349, 8thLeast daily movement69, 23rdMean hourly velocity6,4Maximum velocity29, 9th	4504 6,1 28,8	5946, 1907; 2180, 1894 45, 1935
Wind, direction		
Prevailing direction NW North, days 8 Northeast, days 1 East, days 2 Southeast, days 4 South, days 5 Southwest, days 2 West, days 2 Northwest, days 7	W	
Weather		
Mean relative humidity, percent 63.6 Mean cloudiness, percent 52.0 Number of clear days 110 Number of fair days 110 Number of cloudy days 100 Number of hours bright sunshine 225 Percent of possible hours of bright sunshine 500 Thunder and lightning 25th, 29th	60.7 52.1 11 11 252 55.6	70, 1901, 1902; 30, 192 3 21, 1944; 0, 1927 17, 1907, 1926; 5, 1923 20, 1927; 2, 1944 350, 1944; 137, 192 7

The weather for May was slightly colder than normal, with a little more precipitation than usual.

The mean temperature for the month was 56.1 degrees, while the normal for May is 57.1 degrees. The highest temperature during the month was 87 degrees on the 13th and 24th; the lowest, 28 degrees on the 10th. The domestic heating load for the month was 277 degree-days, compared to the normal of 267 degree-days for May. The heating load for the season to the end of May is now 6310 degree-days, while the normal for this period is 6778 degree-days.

The total precipitation for the month was 4.63 inches, most of which fell in two storms: 1.00 inch on the 3rd and 2.10 inches on the 25th. The normal precipitation for May is 3.60 inches. Thunder and lightning occurred on the 25th and 29th. Rain fell on 14 days. In May, 1945 rain fell on 20 days. There were 225 hours of bright sunshine. The normal for May is 252 hours.

Dr. J. K. Shaw of the Pomology Department makes the following comments: "For the fourth successive year the Massachusetts apple crop has been injured by spring frosts. record of such a condition as far back as records go. previous two years, the program of vegetation was retarded; fruit bloom was a week or ten days later than normal instead of three or four weeks early. Several cold mornings in late April and in May killed many apple blossoms, some in the early cluster bud stage and some later. Bloom was less than the heavy bloom of last year. The apple crop in low lying orchards will be much reduced but we may hope that the total crop will be considerably more than last Much wet weather favored scab development and only thoroughly sprayed orchards of susceptible varieties can escape severe scab injury. Severe frost damage is reported from more southern areas and it looks as though the 1947 eastern apple crop will be below normal.

"There was little or no injury to peaches and pears, and full crops are in prospect. Early strawberry bloom was killed, thus retarding the crop, but present indications are for a good crop." MASI. STATE CONLEGE AMBLEST, MASS. 3 46

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 702

June 1947

Meteorological Observations

FOR

IUNE

1947

F. J. SIEVERS

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Ter	npera	iture	idity	rs		Wind		I	Precipitatio	n	
	Max	cimum	Min	imum		Humic		50	=	s *			
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water
1 2 3 4 5	77 79 68 73	1 P 2 P 11 A 5 P 2 P	44 57 50 42 46	3 A 5 A 12 P 5 A 5 A	60.5 68.0 59.0 57.5 58.5	72 66 86 46 54	15.1 6.6 10.7 15.1 15.2	s s ne s	196 194 228 128 200	17 17 20 13 17	12:30 A	2:30 р	т .79 т
6 7 8 9 10	86 66 53 74 82	3 P 0 A 3 P 4 P 3 P	53 52 46 47 52	1 A 12 P 12 P 1 A 3 A	69.5 59.0 49.5 60.5 67.0	59 75 94 68 69	15.2 0.0 0.0 15.2 15.2	ne n nw se	187 123 104 170 83	18 11 15 20 10	4:30 A 2 P 0 A 0 A	5:30 A 12 P 10 P 4:30 A	.02 .06 1.06 .03
11 12 13 14 15	90 80 79 71 76 71	3 P 1 P 4 P 8 P 12 M	59 56 49 53 60	5 A 12 P 5 A 8 A 12 P	74.5 68.0 64.0 62.0 68.0	75 54 49 93 56	9.6 15.3 15.3 0.3 8.6	se nw se se nw	111 200 91 71 168	18 25 9 20 17	5:30 A 3 A	12 P 3 P	.51 .11
16 17 18 19 20 21	76 70 69 68 74	6 P 3 P 3 P 1 P 1 P 5 P	52 46 56 48 48 46	12 P 5 A 4 A 4 A 3 A 4 A	61.5 61.0 63.0 58.5 58.0 60.0	74 56 80 65 73 65	15.3 10.0 8.4 15.3 3.5 11.0	nw sw s n ne nw	141 60 117 134 71 82	18 10 7 22 7 15	10 р	10:30 р	10. T
22 23 24 25 26	82 85 67 77 85	5 P 2 P 11 A 4 P 3 P	46 47 53 53 60	5 A 6 A 5 A 1 A 4 A	64.0 66.0 60.0 65.0 72.5	64 58 82 90 57	9.0 15.3 1.0 8.8 8.5	n w s n s	72 79 91 118 136	25 12 15 17	6 P 8 A 0 A	10 P 12 P 4:30 P	.01 T .44 .18
27 28 29 30 31	87 91 88 90	5 P 4 P 1 P 3 P	56 56 69 69	5 A 5 A 2 A 4 A	71.5 73.5 78.5 79.5	73 60 74 68	10.7 12.0 7.9 11.0	W S S S	67 53 103 120	11 9 11 11			T T

^{*} Based on least time required to blow one mile.

Goon L. Chin, Observer

June, 1947	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum .30,70, 1st Minimum .29,49, 16th Mean semi-daily .29,923 Range .1,21	30.30 29.55 29.95 0.75	30.54, 1889 29.24, 1902
Air Temperature, in degrees F.		
Highest 91, 28th Lowest 42, 4th Mean 64.6 Range 49 Highest mean daily 79.5, 30th Lowest mean daily 49.5, 8th Mean maximum 76.8 Mean minimum 52.4 Greatest daily range 38, 23rd Least daily range 7, 8th	91.2 40.0 65.7 51.3	101, 1919 34, 1891
Precipitation, in inches		
Precipitation	3.75	9.68, 1922; 0.76, 1908
Snow. —	11	17, 1922, 1945; 4, 1908
Wind, in miles		
Total movement3698Greatest daily movement228, 3rdLeast daily movement53, 28thMean hourly velocity5.1Maximum velocity25, 12th, 22nd	3585 5.0 24.6	4571, 1908; 1409, 1906 48, 1939
Wind, direction		
Prevailing direction SSW North, days 3 Northeast, days 3 East, days 0 Southeast, days 4 South, days 11 Southwest, days 1 West, days 2 Northwest, days 6	WSW	
Weather		
Mean relative humidity, percent. 68.5 Mean cloudiness, percent 31.3 Number of clear days. 15 Number of fair days. 10 Number of cloudy days .5 Number of hours bright sunshine 305 Percent of possible hours of bright sunshine 66.7 Thunder and lightning, 3rd, 14th, 19th, 22nd	66.9 51.1 10 12 8 257 54.0	71, 1903; 28, 1908 22,1908,1941,1943; 1, 1927 23, 1912; 3, 1941 22, 1903; 1, 1923 362, 1908; 102, 1903

June weather was colder than normal, with slightly less rain than usual.

The mean temperature for the month was 64.6 degrees, while the normal for June is 65.7 degrees. The maximum temperature was 91 on the 28th and the minimum, 42 on the 4th. The domestic heating load for June was 88 degree-days compared with the normal for June of 68 degree-days. The total load for the heating season for the period September 1 to June 30 is now 6398 degree-days. The normal for this period is 6846 degree-days.

The total precipitation for the month was 3.22 inches, while the normal for June is 3.75 inches. There were 15 clear days, 10 partly cloudy, and 5 cloudy.

Dr. J. K. Shaw, research pomologist, gives the following report on the progress of vegetation:

"The weather during June continued to be cool and wet, retarding the progress of vegetation. These conditions favored scab development and only orehards intensively sprayed escaped serious infection. Scab spores matured early but infection in our own orchards was much less than last year, probably because of a better spray program. MeIntosh set poorly and the crop in our orchards will be the smallest in many years. Conditions for pollination seemed to be fairly good and it is difficult to find reasons for the poor set. Frost was doubtless a factor and the excessive leaf scab of last year may have weakened the trees. Most other varieties have set at least fairly well though the early blooming Gravenstein had many blossoms killed.

Peaches and pears promise full crops but the blueberry crop will be below normal. Strawberries are yielding well."

MASS. STATE COLLEGE 3 46 AMHERST. MASS.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series Bulletin No. 703

July 1947

Meteorological Observations

FOR

JULY

1947

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

> Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Tei	npera	iture		ity	<u>x</u>		Wind		F	Precipitation	n
	Max	ximum	Min	Jinimum		Humid	Humid of Hounshine		4 5				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water
1 2 3 4 5	90 78 80 79	3 P 4 P 4 P	68 59 52 53	12 P 12 P 5 A 5 A	79.0 68.5 -66.0	70 45 56 55	8.0 15.3 14.7 13.5	n n w	133 214 73 142	32 28 11 20	8:30 P 0 A	12 P 12:30 A	.05
6 7	86 88 85	4 P 4 P 2 P 2 P	51 62 64	5 A 5 A 4 A 12 P 5 A	66.0 68.5 75.0 74.5 63.5	57 69 65	13.0 11.5 15.2	ne sw se	60 130 190	13	S:3() P	9:30 р	.25
8 9	70 81	1 P 2 P	57 65	5 A	73.0	79	0.0 5.9	ne se	$\frac{164}{67}$	7 9	8:30 A	12:30 P	.06
10 11	85 86	4 P 2 P	70 70	0 A 5 A	77.5 78.0	74 77	2.0 7.0	se se	110 120	10	3 A	4 A	.03
12 13 14	84 85 90	3 P 12 M 11 A	60 62 68	12 P 0 A 3 A 3 A	72.0 72.5 76.0 77.5	74 67 74 80	8.0 15.0 15.0	S W D W S W	120 161 120 126	17 15 6 17	6:30 P	7:30 P	.08 .01
15 16 17	87 84 82	12 м ⁹ л 1 р	71 72	2 A	77.5	87 82	7.6 3.2 5.2	, W.	126 126 135	25	4:30 A 3:00 A	9 P 5 P	.59 .07
17 18 19	87 83	1 Р 12 М 3 Р	71 68	2 A 2 A 12 P	79.0 79.0 75.5	81	7.0		149 153	20 17	1.00 A 1 P 5 A	5:30 P 12:30 P	.08
	80 81	5 P	63	12 P 5 A 5 A	76.5 69.5	70	11.5	nw se	155 140	15		1201	
22	75 82	1 P 3 P	66	$\begin{bmatrix} 3 & A \\ 0 & A \end{bmatrix}$	70.5 71.0	74	0.0	se	144 203	28 11	— 3 Р	 4:30 P	1.00
20 21 22 23 24 25 26 27 28	82 85 88	5 P 3 P 1 P	53 58 58	6 A 6 A 6 A	67.5 71.5 73.0	64 63 67	14.8 14.7 10.5	W.	96 64 86	12 10			
29	87 80 86	2 P 9 A 2 P 3 P	64 69 64	4 A 1 A 5 A	70.5 74.5 75.0 75.5	71 88 72 73	14.5 3.0 10.2	s s n	63	9	5 A	3 P	.16
30 31	91	3 P 1 P	60 64	5 A 12 P	75.5 77.0	73 74	6.0 10.4	S W W	61 138	8 14			

^{*} Based on least time required to blow one mile.

THEODORE A. CALIANOS, Observer

Normal	Extremes
30.27 29.59 29.96	30,50, 1892 29,27, 1932
0.68	0.97, 1892; 0.47, 1938
93.9 46.4 70.8 47.7	104, 1911 40 1890 1898 74.7, 1921; 66.3 1891
4,10	14.51, 1897; 0.70, 1929 20, 1915; 4, 1924
3422	5097, 1909; 1109, 1894
4.6 28.7	44, 1936
SW	
68.1 50.9 10 14 7 268	70, 1943; 31, 1924 22, 1923; 0, 1915 24, '09, '21; 4, 1946 18, 1889; 0, 1910 371, 1910; 180, 1931
	30,27 29,59 29,59 0 68 93,9 46,4 70,8 47.7 4.10 11 3422 4.6 28.7 8W

July was both hotter and drier than normal. The mean of 73.2 degrees equals 1935 and has been exceeded only by 1921 and 1911. The normal maximum temperature, however, was not reached with 91 degrees on the 30th. Precipitation was recorded on 14 days but most of it fell in two storms on the 16th and the 22d, leaving the month 1.37 inches short and the year 1.51 inches below normal at the month's end. There was more wind movement than normal and more than in any year since 1932. Hours of bright sunshine were more than normal. These things all combined to the benefit of the tobacco harvest but lawns, gardens and field crops dependent on July moisture showed the effects of greater than normal temperatures and the deficiency in normal precipitation.

Dr. J. K. Shaw, research pomelogist, makes the following statement: "The progress of vegetation during July was generally normal. All fruit crops progressed as usual and consequently there is nothing special to be put on record. Our own apple crop, especially McIntosh, will be short while that of the state is reported to be above that of last year. It will be spotty, some orchards yielding very well and others poorly. The apple crop in the Appalachian region was badly injured by spring frosts, while there was little injury further north. In general the fruit crop of the country seems to be at least up to normal and we may expect a good supply in 1947."

MASS. STATE COLLEGE AMHERST, MASS.

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series — Bulletin No. 704

August, 1947

Meteorological Observations

FOR

AUGUST

1947

H. N. STAPLETON

OBSERVATORY

Latitude, 42 ° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

> Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Tei	mper	ature		ity	rs		Wind		F	Precipitation			
	Ma	ximum	Mir	nimum		Humid	of Hours inshine	50	1	*es					
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hour Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Honr*	Began	Ended	Water		
1 2 3 3 4 5 6 7 8 9 9 10 11 11 12 13 14 15 16 17 18 19 20 21 22 23 22 24 22 25 26 27 28 29 20 20 20 20 20 20 20 20 20 20 20 20 20	74 77 82 83 88 89 90 86 87 89 93 94 95 73 72 80 88 87 77 87 90 94 88 77 87 88 88 77 87 88 88 88 77 88 88	5 P 3 P 3 P 3 P 4 P 2 P 4 P 5 P 1 A 2 P 5 P 3 P 2 P 2 P 2 P 2 P	52 48 53 61 62 69 67 60 53 62 70 74 71 60 59 67 62 69 67 67 68 67 68 67 68 68 68 68 68 68 68 68 68 68 68 68 68	5 A 5 A 5 A 6 A 5 A 3 A 12 P 5 A 12 P 5 A 12 P 2 A 12 P 2 A 7 A 7 A 7 A 6 A 3 A 7 A 6 A 6 A 6 A 6 A 6 A 7 A 12 P 6 A 7 A 7 A 7 A 8 A 8 A 8 A 8 A 8 A 8 A 8 A 8 A 8 A 8	63.0 62.5 67.5 72.0 74.0 73.5 70.5 77.5 82.0 84.5 66.5 76.5 73.5 78.5 66.5 73.5 78.5 66.5 78.5 78.5 78.5 78.5 78.5 78.5 78.5 78	70 73 73 75 79 81 91 84 87 77 74 73 86 81 77 93 86 81 85 88 88 88 88 88 88 88 88 88 88 88 88	15.5 14.5 10.2 8.8 8.0 11.0 5.7 5.1 2.1 10.3 14.1 8.7 8.5 8.0 0.2 10.0 0.2 10.0 0.4 3.1 10.8 8.0 13.6 4.0 7.5 4.0 7.5 4.0 13.6 13.6 13.6 13.6 14.0 15.7 16.0 16.0 16.0 16.0 16.0 16.0 16.0 16.0	ne w sc s s sw s sw sw w nc ne sw me n nw w sw sw ne e w nc sc sc s sw sw mc ne sc sc s sc sc sw sw max sw sw me e sc sc sc sc sc sc sc sc sc sc sc sc sc	157 85 71 81 73 82 93 93 58 115 57 94 67 67 118 96 124 88 81 67 63 72 81 106 110 110 110 110 110 110 110 110 11	10 8 8 5 6 6 5 9 13 12 13 7 7 7 7 8 12 11 12 8 13 13 13 13 13 13 13 13 13 13	7:30 P 6:30 A 3:45 A 11:30 A 8:30 P	8:30 A 8:30 P 4:00 P 8:15 A	.06 .02 .50 .14		

^{*} Based on least time required to blow one mile.

THEODORE A. CALIANOS, Observer

August, 1947	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.35, 17th, 18th Minimum 29.80, 1st Mean semi-daily 30.097 Range 0.55	30.32 29.61 30.00 0.71	30.50, 1934 28.87, 1930
Air Temperature, in degrees F.		
Highest 95, 14th, 15th, 25th Lowest 48, 2nd Mean 73.0 Range 47 Highest mean daily 84.5, 14th Lowest mean daily .62.5, 2nd Aican maximum 84.5 Mean minimum 61.5 Greatest daily range 35, 11th Least daily range 13, 9th, 16th, 17th	91.6 42.4 68.6 48.1	100, 1918 34, 1940 62.4, 1903
Precipitation, in inches		
Pre ipitation	4.08	8.40, 1928; .31, 1894
Snow. — — — — — — — Maximum precip. in 24 hrs. — 0.95—26th Number of days with .01 or more — — 6	, II	16, 1892, 1933; 4, 1899
Wind, in miles		
Total movement2820Greatest daily movement160, 26thLeast daily movement52, 28thMean hourly velocity3,8Maximum velocity25, 29th, 31st	3127 4.2 22.7	4,271, 1910; 1,920, 1894 40, 1941
Wind, direction		
Prevailing direction SSW North, days 1 Northeast, days 6 East, days 1 Southeast, days 5 South, days 4 Southwest, days 6 West, days 6 Northwest, days 2	SW	
Weather		
Mean relative humidity, percent 80.4 Mean cloudiness, percent 33 Number of clear days 12 Number of fair days 11 Number of cloudy days 8 Number of hours bright sunshine 243 Percent of possible hours of	70.2 49.7 13 0 237	67, 1901, 1946; 27, 1923 23, 1941; 0, 1915, 1929 25, 1912; 3, 1939 18, 1901, '28; 2, 1910, '23 332, 1941; 152, 1915, 1929
bright sunshine 50,6 Thunder and lightning 26th First frost	55,2 Sept. 21	Aug. 22, 194,195; Oct. 13,109

The weather for the month of August provided a most uncomfortable combination of heat and humidity with few breaks, and recorded below normal precipitation. It was the warmest August since 1939 and recorded the highest mean relative humidity in the past 18 years. While the high of 95 degrees was recorded on but 3 days, a temperature of 90 degrees or more was recorded on 8 days of the month. The highest maximum, 100 degrees on August 7, 1918, was never reached but only 1937 and 1939 have recorded warmer Augusts since 1837, the beginning of weather records at Amherst. Of the 1.69 inches of precipitation for the month, 1.45 inches was recorded in 2 storms on the 16th and 26th, and measurable precipitation occurred on 6 days. The cumulative deficiency in precipitation is shown by the 7.64 inches recorded for June, July, and August against a normal of 11.93 inches for this period. There were no high winds during the month.

Dr. J. K. Shaw, research pomologist, makes the following statement: "The weather during August has been characterized by a lack of rainfall. While the drought this late in the season has not been severe enough to seriously injure the fruit crop, it may have slowed the rate of growth of the fruit. Probably the apple harvest will be a little later than average, due mostly to late bloom. The apple crop in the University orchards will be below average due principally to a short McIntosh crop. Reports indicate that the crop over the State will be larger than the small crop of 1946. Peaches are yielding well and other tree fruits are good. The blueberry crop proved to be somewhat better than anticipated."

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 705

September 1947

Meteorological Observations

FOR

SEPTEMBER

1947

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

3 81 3 9 60 6 0 70.5 70 13.0 w 66 7 7 7 7 7 7 7 7			Tei	npera	ature		ity	rs	Wind Precipitation						
1 81 2 P 53 6 A 67.0 72 13.2 w 52 13 2 82 2 P 63 4 A 72.5 95 3.6 se 97 20 5:30 A 11 A 1.21 3 81 3 P 60 6 A 70.5 70 13.0 w 66 7 4 85 3 P 56 6 A 70.5 80 13.0 nw 40 10 5 77 3 P 59 6 A 68.0 86 6.9 s 33 5 6 81 3 P 65 6 A 73.0 88 6.5 s 72 8 7 87 3 P 68 0 A 77.5 83 12.0 se 96 13 9 83 3 P 53 7 A 69.5 74 12.8 sw 78 13 10 86 3 P 53 7 A 69.5 74 12.8 sw 78 1		Maximum		Minimum			Humid	of Houn	20		S				
2 82 2 P 63 4 A 72.5 95 3.6 sc 97 20 5:30 A 11 A 1.21 3 81 3 P 60 6 A 70.5 70 13.0 w 66 7 4 85 3 P 56 6 A 70.5 80 13.0 nw 40 10 5 77 3 P 59 6 A 68.0 86 6.9 s 33 5 6 81 3 P 65 6 A 73.0 88 6.5 s 72 8 7 87 3 P 68 0 A 77.5 83 7.5 s 72 9 8 82 3 P 62 12 P 72.0 83 12.0 sc 96 13 9 83 3 P 59 6 A 71.0 82 7.5 s 46 9 10 86 3 P 53 7 A 69.5 74 12.8 sw 78 13 11 86 4 P 69 3 A 77.5 77 10.1 s 133 15 12 89 1 P 70 6 A 79.5 74 12.2 s 181 20 13 89 11 A 70 6 A 79.5 79 12.0 sc 116 8 14 81 3 P 66 12 P 73.5 92 3.2 sc 116 8 11 83 2 P 64 2 A 73.5 90 3.0 sc 67 17 17 77 3 P 44 6 A 60.5 66 12.4 nw 57 7 18 79 3 P 47 6 A 63.0 65 10.0 sc 73 11 19 76 1 P 49 12 P 62.5 84 0.0 sc 153 15 20 65 3 P 41 5 A 53.0 58 12.3 s 99 8 21 72 12 P 47 0 A 59.5 84 0.0 sc 153 15 22 74 5 A 43 12 P 58.5 86 0.9 n 234 25 60 A 24 61 1 P 34 5 A 47.5 73 4.4 s 63 15 25 60 3 P 42 12 P 51.0 77 5.0 n 137 20 26 59 3 P 35 5 A 47.5 73 4.4 s 63 15 27 58 3 P 29 6 A 43.5 58 12.0 n 90 20 28 60 3 P 29 6 A 44.5 62 11.9 sc 70 10	Day	Degrees	Time	Degrees	Time	Mean	Mean Relative	!	Prevailing Direction	Total Movemer	Max. Mil per Hour	Began	Ended	Water	
29 67 2 P 30 6 A 48.5 72 9.4 sc 125 40 30 53 0 A 37 12 P 45.0 61 6.4 nw 282 25 5 A 6 A .0	2 3 4 5. 6 7 8 9 10 11 12 13 14 15 16 17 18 20 21 22 23 24 25 26 27 28 29 29 29 20 20 20 20 20 20 20 20 20 20 20 20 20	82 81 85 77 81 87 88 88 88 88 88 88 88 88 77 76 65 72 65 74 60 65 66 67	2 P 3 P 3 P 3 P 3 P 1 P 1 P 3 P 1 2 P 1 2 P 3 P 1 2 P 3 P 1 2 P 3 P 2 P 3 P 2 P 2 P 2 P 2 P 2 P 2 P	63 60 56 59 65 68 62 59 70 66 64 49 44 47 49 41 42 35 29 30	4 A 6 A 6 A 6 A 12 P 6 A A 12 P 6 A A 12 P 6 A A 12 P 6 A A 12 P 6 A A 12 P 5 A 6 A 12 P 5 A 6 A 6 A 6 A	72.5 70.5 70.5 70.5 68.0 77.5 72.0 71.0 57.5 79.5 79.5 73.5 61.0 60.5 63.0 59.5 58.5 46.5 47.0 43.5 44.5	95 70 80 86 88 83 83 82 74 77 74 66 65 58 84 85 87 77 55 58 62 72	3.6 13.0 6.9 6.5 7.5 12.0 6.2 12.0 3.2 3.0 6.2 12.4 10.0 0.0 0.9 12.3 0.0 9.5 12.0 9.5 12.0 9.5 12.0 9.5 12.0 9.5 12.0 9.5 12.0 9.5 12.0 9.5 12.0 12.0 12.0 12.0 12.0 12.0 12.0 12.0	se w nw s s s se se se se se nw nw se s s se se se nw nw se se se se nw ne se se se se se se se se se se se se se	977 666 400 333 722 722 966 466 788 1333 1811 156 67 73 1533 99 124 234 1488 63 137 156 67 70 125	20 7 10 5 8 9 13 15 20 20 8 17 17 7 11 15 8 15 25 18 15 20 28 15 20 28 20 28 20 20 20 20 20 20 20 20 20 20 20 20 20	6:30 P 11:30 A 1 P 3 A	9:00 p 12:30 p 5 p 4:15 A	1.21 .16 .06 .48 .09	

^{*} Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

September, 1947	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.64, 18th Minimum 29.62, 22nd Mean semi-daily 30.176	30.45 29.57 30.06	30.65, 1924 28.41, 1938
Range 1.02	.88	1.99, 1938; .57, 1910
Air Temperature, in degrees F.		
Highest 89, 12th, 13th Lowest 29, 27th, 28th Mean 62.9 Range 60 Highest mean daily 79.5, 12th, 13th Lowest mean daily 43.5, 27th Mean maximum 74.9 Mean minimum 50.9 Greatest daily range 37, 29th Least daily range 15, 14th	87.7 33.2 61.7 54.5	97, 1895, 1929 24.5, 1914
Precipitation, in inches		
Precipitation	4.24	14.55, 1938; .52, 1914
Snow	10	16, 1933; 3, 1903
Wind, in miles		
Total movement 3306 Greatest daily movement 282, 30th Least daily movement 33, 5th Mean hourly velocity 4.6 Maximum velocity 40, 29th	3271 4.5 25.9	4,686, 1896; 1,414, 1894 80, 1938
Wind, direction		
Prevailing direction S North, days 4 Northeast, days 1 East, days 0 Southeast, days 9 South, days 9 Southwest, days 1 West, days 2 Northwest, days 4	WSW	
Weather		
Mean relative humidity, percent. 75.0 Mean cloudiness, percent 23.0 Number of clear days. 16 Number of fair days. 8 Number of cloudy days 6 Number of hours bright sunshine 240 Percent of possible hours of	73.6 49.9 10 10 10 201	70, 1934; 27, 1908, 1914 19, 1932; 2, 1907, 1928 19, 1908; 3, 1889 20, 1934; 3, 1941 255, 1916; 106, 1934
bright sunshine 72.6 Thunder and lightning 13th, 15th First frost 27th	54.9 Sept. 21	Oct. 13, 1909, Aug. 22, 1894-95

September weather was not violently extreme but it avoided the normal in almost every category. While the mean temperature for the month held 1.2 degrees above normal, 184 degree-days of heating load were accumulated against a normal of 149 for the largest September heating load since 1938. Recording a maximum temperature of 89 degrees on the 12th and 13th at 1.3 degrees above normal, the 29-degree minimum on the 27th and 28th gave the month's greatest range and lowest minimum since 1942. With the mean date of first frost September 21, September 27 recorded the first frost of this season for the earliest date since 1940.

Precipitation continued the trend of the pest three months with a below normal 2.84 inches for the month and a below normal 28.32 inches accumulated for the year. Normal for the month shows a deficiency of 1.40 inches, and the year short of normal by 5.30 inches. The storm of the 2d yielded 1.21 inches and there was measurable precipitation on six other days of the month. High winds occurred on the 29th with the maximum velocity recorded at 40 miles per hour.

W. H. Thies, Extension Professor of Horticulture, contributes the following report: "The McIntosh apple harvest which normally begins around September 15 is delayed this year for at least two reasons. In the first place, bloom was about 10 days late last spring, and since approximately the same number of days clapses between bloom and maturity, later maturity might be expected this year. Then, too, weather conditions in early September were unfavorable for the coloring of apples. In fact, less color was apparent on September 25 than we commonly find on September 15. Not until the last week in September, when clear days followed cold nights, did the coloring process proceed normally. In spite of this delayed coloring, apples have hung on the tree unusually well this fall. As a result, more McIntosh will be harvested during the first week in October than in the last twenty-five years."

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 706

October 1947

Meteorological Observations

FOR

OCTOBER

1947

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Ter	nper	ature		ity	£	Wind Precipitation					
	-	dmum		imum		Mean Relative Humidity	Number of Hours Bright Sunshine	ng in	ent	illes tr*			
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative	Number Bright 5	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water
1 2 3 4 5 6 7 8 9 10 11 12 13	57 61 69 72 75 80 81 74 64 67 71 73 69	3 P 2 P 4 P 3 P 3 P 3 P 3 P 3 P 3 P 3 P 3 P 3	30 40 32 37 52 51 52 54 42 34 38 42	1 6 A 1 12 P 7 A 4 A 2 A 3 A 6 A 12 P 112 P 7 A 5 A 7 A 3 A 3 A	43.5 50.5 50.5 54.5 63.5 65.5 64.0 53.0 50.5 53.5 55.5 55.5 55.5 55.5 55.5	754 70 78 80 79 76 75 82 57 66 77 69 78	10.2 11.7 8.5 6.6 6.3 8.2 8.3 1.0 9.0 11.3 7.0 9.2 10.8	nw w sw sw s s n ne s se se	195 154 24 62 119 99 109 97 161 74 47 57 109	20 32 3 9 10 10 10 12 20 6 7 7 7			<u></u>
14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30	77 81 86 83 78 78 74 74 83 59 67 75 80 74	3 P 2 P 3 P 2 P 2 P 2 P 2 P 2 P 2 P 3 P 3 P 3 P 3 P 3 P 3 P 3 P	48 8 6 1 7 6 3 4 2 3 6 5 2 3 9 2 9 4 9	12 P 7 A 7 A 5 A 4 A 12 P 6 A 12 P 7 A 12 P 7 A 12 P 7 A 12 P 7 A 7 A	62.5 59.5 66.0 67.0 67.5 70.5 66.0 63.0 55.0 64.0 45.5 45.0 61.0 61.5	68 69 84 88 88 58 41 57 40 73 78	10.3 11.1 6.5 5.4 2.2 3.2 10.9 10.9 10.8 10.8 10.7 8.0 6.6 8.8 2.7	nw se se se s n n n s n ne sw w se	76 52 40 55 76 110 92 156 40 196 214 71 39 43 152	8 7 6 7 7 12 13 25 5 28 25 10 6 8 11	8:30 P	10 р	.08
30 31	68 48 46	3 P 0 A 4 P	48 40 36	12 P 12 P 10 P	58.0 44.0 41.0	88 84 77	0.0 0.0 0.0	se ne ne	119 161 241	15 12 22	6 P 1:30 A 0 A	8:30 P 12 P 11 A	.17 .79 1.00

^{*} Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

October 1947	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.52, 25th Minimum 29.71, 19th Mean semi-daily 30.142 Range 0.81	30.53 29.42 30.06 1.11	30.68, 1929 29.00, 1926 1.47, 1926, .76, 1899
Air Temperature, in degrees F.		
Highest 86 16th Lowest 23, 25th Mean 57,4 Range 63 Highest mean daily 70,5 19th Lowest mean daily 41,0,31st Mean max'mum 71,6 Mean minimum 43,2 Greate t daily range 41,25th Least daily range 8,30th	79.4 23.4 50.5 56.0	90.5, 1908 17, 1936 56.4, 1920, 43.2, 1890
Precipitation, in inches		
Precipitation	3.29	8.81, 1911; .01, 1924
Snow. Maximum presip, in 24 hrs 1.00. 31st Number of days with .01 or more 4	0	15, 1913; 1, 1897, 1924
Wind, in miles		
Total movement3240Greatest daily movement241, 31stLeast daily movement24, 3rdMean hourly velocity4.1Maximum velocity32, 2nd	5.4 29.5	5,467, 1910; 2,540, 1894 42, 1937
Wind, direction		
Prevailing irection SSF North, days 4 Northeast, days 4 East, days 0 Southeast, days 9 South, days 7 Southwest, days 3 West, days 2 Northwest, days 2	W.	
Weather		
Mean relative humidity, percent	69.0 48 10 10 11 175.0	67, 1932, 18, 1924 21, 1938; 1, 1911 17, 1924; 3, 1938 19, 189 <i>t</i> ; 3, 1924 232, 23, 38; 91, 1913
bright sunshine	51.3	
First Frost	Sept. 21	Aug. 22,'94,'95; Oct. 15,'39

The weather during October was bright, warm and dry, with 18 clear and 6 fair days, and all but the last 3 days of the month recording some amount of bright sunshine. Until the storm beginning the 29th, the month had recorded .08 inches of rainfall and the 4 previous months had recorded 10.48, bidding for a record drouth in the 5-month period and developing the woodland into a fire hazard of major proportions. The month finally recorded a deficiency of 1.25 inches below normal and the 5 months period a cumulative 12.52 inches against a normal of 19.46 inches of rainfall.

Examination of the records maintained since 1836 discloses that this month established a record mean daily temperature with 57.4 degrees. October 1920 with 56.4 degrees is nearest this record and 1879 with 56.0 is next. On only 2 days were minimum temperatures of less than 32 degrees recorded. The 23-degree minimum on the 25th and the high of 86 on the 16th gave both the highest maximum and the greatest range for the month since 1941. Seven days of the month recorded mean temperatures in excess of 65 degrees. The heating load for the month is 249 degree-days and cumulative for the season 433, slightly higher than last year's cumulative, 416, but still only 72% of normal heating requirement to October 31.

There were no high winds during the month although the fresh to strong winds which occurred between the 19th and the 24th gave trouble in nearby localities affected by forest fires.

W. H. Thies, Extension Professor of Horticulture, contributes the following report: "The month of October has been very favorable for fruit trees in some cases, and less so in others. On fertile, retentive soils, leaves have functioned later than usual thus encouraging the development of fruit buds and the storage of carbohydrates. This will tend to promote tree vigor and productiveness in 1948. There is little evidence, in properly managed orchards, of a lack of maturity in the trees, a condition which commonly precedes one type of winter injury. On poorer soils, trees have probably suffered from a fluctuating water supply and from certain mineral deficiencies which accompany a water shortage. Such trees lost their leaves even before severe frosts had occurred. October weather has been favorable for the continued development of strawberry 'runners.' In fact, some matter row plantings undoubtedly have more plants than are needed for a full crop."

Correction: An error in barometric readings on page 3 in the June 1947 bulletin gives 30.70 as the maximum on the 1st, 29.49 as the minimum on the 16th and 1.21 as the range. The correct readings are as follows: The maximum was 30.34 on the 21st; the minimum, 29.65 on the 15th; and the range, .69.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 707

November 1947

Meteorological Observations

FOR

NOVEMBER

1947

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Ter	ature		ity	rs.	Wind Precipitation							
	Maximum		Minimum			Humid	of Hou	20	-	es *				
Day	Degrees	Time	Ведгеея	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 3 4 4 5 6 6 7 8 8 9 10 11 11 12 13 11 15 11 15 11 15 12 20 21 22 3 24 25 26 27 28 29 30	55 60 59 53 59 48 48 47 47 42 42 42 42 43 41 43 41 43 41 42 42 42 42 43 43 43 44 43 43 43 43 43 44 45 46 47 48 48 48 48 48 48 48 48 48 48	3 P 3 P 1 P 1 P 1 P 1 P 2 P 2 P 1 2 P 3 A 1 P 3 P 1 2 P 3 P 1 2 P 3 P 1 2 P 3 P 1 2 P 3 P 1 2 P 3 P 1 2 P 3 P 1 2 P 3 P 1 2 P 3 P 1 2 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P 1	31 27 28 41 41 45 40 34 29 26 32 29 26 32 27 28 28 29 26 27 28 28 29 20 20 20 20 20 20 20 20 20 20 20 20 20	7 A 7 A 7 A 7 A 6 A 12 P 7 A 12 P 12 P 12 P 12 P 12 P 12 P 12 P 12 P	43.0 43.5 43.5 47.0 50.0 48.5 40.0 34.0 40.0 34.0 35.5 37.5 33.5 33.5 33.5 42.0 35.5 37.5 37.5 37.5 37.5 37.5 37.5 37.5	57 70 65 77 61 88 80 80 60 66 67 79 64 64 62 55 65 68 66 67 77 76 77	10.3 10.3 2.7 0.0 0.0 0.0 0.0 0.0 4.2 7.5 0.0 0.6 3.2 9.8 9.7 5.0 7.0 0.0 0.0 0.0 4.0 0.0 0.0 0.0 0.0 0.0 0	n w nw nw c c se nw n n nw nw nw nw nw nw nw nw nw nw nw	128 31 99 88 8 56 56 278 193 307 1700 266 51 156 41 388 332 127 99 72 40 52 238 67 75	9 5 10	12:30 P 2 P 11:30 A 5:30 P 0 A 2:30 P	8 P + P 7 P 12 P 2 P 10:30 A	.06 .01 1.74 .27 2.46	Т

^{*} Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

November, 1947	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.55, 21st, 22nd Minimum 29.17, 24th, 25th Mean semi-daily 30.056	30.59 29.30 30.05 1.29	30.87, 1932 28.73, 1904
Range	1.29	1.84, 1904; .89, 1943
Air Temperature, in degrees F.		75 1024 1020
Highest .60. 2nd Lowest 15. 30th Mean 38.0 Range 45 Highest mean daily 50.0 5th 7th. lowest mean daily 24.5, 30th Mean maximum 46.1 Mean minimum 29.8 Greatest daily range 33, 2nd Least daily range 3, 6th	66.0 12.9 38.9 53.0	75, 1924, 1938 _4, 1938 44.1, 1931; 33.6, 1901
Precipitation, in inches		
Precipitation. 5,63 Snow. T, 29th Maximum precip. in 24 hrs. 2,46, 12th	3.41 2.34	8,64, 1927; ,63, 1917 13,50, 1938
Number of days with .01 or more8	, ,	22, 1921; 2, 1904
Wind, in miles		
Total movement3839Createst daily movement388 17thLeast daily movement18, 2ndMean hourly velocity5.3Maximum velocity32, 12th	4531 6.3 30.3	5.978, 1906; 2 589, 1889 44, 1938
Wind, direction		
Prevailing Direction NW North, days 7 Northeast, days 0 East, days 1 Southeast, days 1 South days 4 Southwest, days 0 West, days 4 Northwest, days 13	W.N.W.	
Weather		
Mean relative humidity, percent. 68.7 Mean cloudiness, percent 78 Number of clear days 7 Number of fair days 6 Number of cloudy days 17 Number of hours bright sunshine 101 Percent of possible hours of	70.6 55.1 6 9 15 121 41.3	72 1900 1927 34, 1917 45, 1903; 1 1900, 1911, 1927 16, 112; 4, 189, 130, 132, 135 24, 1927; 9, 1905, 1917 182, 1903; 66, 1927
bright sunshine		Oct. 10 '25: Der. 13 1941

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1946.

REMARKS

The weather for the month was both colder and wetter than normal. The 5.63 inches of precipitation recorded the most rainfall for the month since 1942 and brought the year's cumulative to 35.99 inches which is still 4.33 inches under the normal. The mean temperature for the month was registered at an even 38 degrees, 0.9 degrees below normal for the coldest November since 1939.

No serious storms occurred during the month although 2.46 inches of rain fell on the 12th.

The heating load for the month was 826 degree-days, the greatest for the month since 1939. Cumulative heating load for the season now stands at 1259, which is 130 degree-days below the season normal and indicates only 18.4 instead of the normal 20.3% of mean heating load to have been accumulated at the end of the month.

W. H. Thies, Extension Professor of Horticulture, contributes the following report: "From the standpoint of fruit crops, the weather during November has been quite favorable. Temperatures have been low enough to check tree growth and to encourage hardening of tissues. Only once has the temperature dropped as low as 15° in contrast to near zero weather in late November in years past. Conditions are now ideal for mulching the strawberry planting to provide protection against deeper freezing and consequent heaving of the soil.

Rainfall amounting to 5.63 inches has helped to replenish soil moisture although we are about 5 inches short of normal for the year. In this connection, it should be pointed out that total rainfall is of less significance than the amount which actually penetrates the soil where it falls. An apple tree on a sloping area, for example, may be greatly benefited by contouring or by a heavy mulch which encourages penetration of water otherwise lost in flowering to a lower level. Good management of a fruit planting tends to conserve both moisture and fertility."

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series Bulletin No. 708

December 1947

Meteorological Observations

FOR

DECEMBER

1947

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

> Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

	Temperature				ity	rs		Wind			Precipit	ation		
Day	Max	Time mumi	Min Segrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended .	Water	Snow
11	40 [13	17.1	26.5	65	5.0	50	75	10	-			1 0.
3 4	35 52 49	2 P 2 P 4 P 0 A	20 32 23	5 A 6 A	27.5 42.0 36.0	80 63 47	0.0 0.0 7.4	n w	19 70 193	6 10 25	1.5	2 A	.03	
15	23 30	0 A 2 P	18 19	12 P 7 P 8 P	20.5	79 74	0.0	n n w	73 157	10	8 A	() P	.43	2.5
7 8	39 46 40	1 P 3 P 0 A	22 34 26	0 A 0 A 12 P	30.5 40.0 33.0	75 83 50	2.1 0.0 0.0	W 50	79 103 480	10 18 40	6 P	12 P 10 A	.13 .18	T
10 11 12	35 37 35	3 P 2 P 3 P	19 20	8 A 12 P 7 A	27.0	64 76 74	5.2 7.0 2.0	n w n w	83 88 92	18 13 17	7 P 1 A	12 P 3 A	.15	2.5
13 14 15	32 35 40	2 P 3 P 4 P	12 15 8 12	12 P 7 A 8 A	23.5 23.5 21.5 26.0	62 65 78	2.6 4.3 6.8	n w	108 49 61	18	10 p	12 p	.10	
16 17	51	12 м 3 Р	34 28	12 P 9 P	42.5 31.5	68 54 49	5.0 6.0 2.0	n w n w	218 187 370	32 32 32	3.3	12 P 7 A	.23	
18 19 20	38 28 30	8 A 3 P 3 P	14 12 11	12 P 5 A 7 A	26,0 20,0 20,5	48 60	7.0 5.0	D III	137 98	15				
20 21 23 24 25 26 27 28 29	35 33 23	12 m 3 P 12 P 2 P	16 22 21	0 A 8 A 4 A	25.5 27.5 22.0	64 55 88	2.8 6.0 0.0	n w n w	75 156 130	2.5 2() 14	8 A	12 P	.48	4.0
24 25	31	4 P	10	12 P 8 A	19.5 15.5	64 80	9.0 6.0	W.	314	25	0.7	2:30.\	.03	.25
26 27 28	29 32 31	12 P 1 P 3 P	15 13	2 A 11 P 7 A	18.0 23.5 22.0 15.0	90 83 72 55	0.0 0.7 2.0	n w n n w	196 398 202	25 25 32 32 15	10 A	12 P	.55	8.0
29 30 31	19 26 25	0 A 2 P 8 P	11 4 11	12 P 8 A 7 A	15.0 15.0 18.0	55 45 55	7.0 9.1 0.0	n n	384 157 55	32 15 7				

^{*} Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

		1
December, 1947	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		-
Maximum 30.47, 1st Minimum 29.25, 16th Mean semi-daily 30.014	30.66 29.27 30.06	30.96 1889 28.85, 1915
Range 1.22	1.38	1.78, 1895; 1.01, 1892
Air Temperature, in degrees F.		
Highest 52, 3rd Lowest 0, 25th Mean 25, 6 Range 52 Highest mean daily 42,5, 16th Lowest mean daily 15, 29th, 30th Mean maximum 34,3 Mean minimum 16,8 Greatest daily range 28, 15th Least daily range 2, 23rd	54.3 1.0 27.5 56.2	65.5, 1908 22.5, 1917 36.9, 1891; 17-1, 1917
Precipitation, in inches		
Precipitation	3.39 8.50	7.77, 1901; .58, 1943
Snow 17.50 Maximum precip, in 24 hrs. 0.53, 26th Number of days with .01 or more. 11	10	26.50, '02, T, 1891, 1943 17, 1902; 4, 1892, 1943
Wind, in miles		
Total movement 4849 Greatest daily movement 480, 9th Least daily movement 19, 2nd Mean hourly velocity 65	4710 6.3	6,694, 1925; 3,239, 1918
Maximum velocity	31.4	48, 1938, 1943, 1944
Wind, direction		
Prevailing direction NW North, days 8 Northeast, days 0 East, days 1 Southeast, days 2 South, days 2 Southwest, days 0 West, days 4 Northwest, days 14	WNW	
Weather		
Mean relative humidity, percent 66.6 Mean cloudiness, percent 77 Number of clear days 7 Number of fair days 10 Number of cloudy days 11 Number of hours bright sunshine 115 Percent of possible hours of bright sunshine 40.6 Thunder and lightnine 16 First Snow Dec. 3	69.4 51.9 50 11 128 45.2	71, 1929; 39, 1919 15, 1890; 2, 1922, 1933 16, 709; 4-789, 30, 731, 736, 738 23, 733; 7, 709, 23 172, 1896; 63, 1933

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1946.

ANNUAL SUMMARY

Annual, 1947	Normal	Extremes
(Readings reduced to sea level) Barometer, in inches		
Maximum 30.89, Jan. 1 Minimum 28.63, Mar. 3 Mean semi-daily 30.002 Range 2.26	30.81 28.95 30.01 1.85	31.05, 1920, 1943 28.41, 1938 2 47 1938; 1.38, 1933
Air Temperature, in degrees F.		
Highest 95, Aug. 14, 15, 25 Lowest -1, Jan. 10 Mean 48.4 Range 96 Highest mean daily 84.5, Aug. 14 Lowest mean daily 12.5, Jan. 10, Feb. 21	95.7 -12.2 47.4 107.8	104, 1911 -26, 1904 49.8, 1921; 44, 1904
Mean maximum 58.8 Mean minimum 38.1 Greatest daily range 44, Oct. 25 Least daily range 2, Dec. 23	57.8 36.5	
Precipitation, in inches		
Precipitation .38.32 Snow .43.75 Maximum precip. in 24 hrs. .2.46, Nov. 12	43.70 47.78	59.00, 1938; 30.68, 1908 89.00, 1893; 24.50, 1919
Maximum precip. in 24 hrs2,46, Nov. 12 Number of days with .01 or more	124	146, 1945; 96, 1924
Wind, in miles		
Total movement51530Greatest daily movement480 Dec. 9Least daily movement7. Jan. 15Mean hourly velocity5.9Maximum velocity45. Mar. 3	52,223 5.8 39.5	63,571, 1908; 36,257, 1894 80, 1938
Wind, direction		
Prevailing direction WNW North, days 62 Northeast, days 17 East, days 7 Southeast, days 50 South, days 61 Southwest days 30 West, days 47 Northwest, days 91	W	
Weather		
Mean relative humidity, percent 68,3 Mean cloudiness, percent 51,1 Number of clear days 136 Number of fair days 109 Number of cloudy days 120 Number of hours bright sunshine 2385 Percent of possible hours of bright sunshine 53,5	67.6 51.7 116 123 126 2.353	60, '98, '01, '02; 41, '08, '24 217 1941; 59, 1927 182, 1912; 64, 1936 179, '01, '02; 71, 1910, 1941 3038, 1941; 1864, 1902
Last snow Apr. 21 First snow Dec. 3 Last frost May 15 First frost Sept. 27	April 15 Nov. 6 May 14 Sept. 21	Mar.14, '10; May 11 '07, '45 Oct. 10, '25; Dec. 13, '41 Apr. 14, '43; June 8, '32 Aug. 22, '94, '95; Oct. 13, '09

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 709

January 1948

Meteorological Observations

FOR

JANUARY 1948

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Tei	nper	ature		ity	rs		Wind			Precipit	ation	
	Max	ximum	Mir	imum		Humid	of Hou	b0	it it	es *				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5	28 29 35 34	9 A 12 P 2 P 3 P	22 21 25 26	11 P 9 A 8 A 7 A	25.0 25.0 30.0 30.0	53 86 68 75	2.0 0.0 8.8 4.0	n ne nw e	137 275 352 56	14 20 25 5	2 A	6 Р	.61	3.5
5 6 7 8	33 33 32 35	11 A 1 P 2 P 3 P	29 26 22 17	1 A 12 P 9 A 11 P	31.0 29.5 27.0 26.0	71 79 78 66	3.0 2.0 3.6 4.6	n ne nw nw	137 38 92 67	12 5 9 12	5:30 P 0 A	12 P 3 A	.11	1.25 .25
10 11	39 28 29	3 P 0 A 3 P	18 6 -1	0 A 12 P 4 A	28.5 17.0 14.0	81 47 59	3.6 4.6 7.3	w n nw	137 390 100	32 32 20	3 A	8 A	.14	1.0
12 13 14 15	32 34 31 16	10 P 1 P 0 A 3 P	31 11 -2	5 A 12 P 12 P 8 A	14.0 32.5 21.0 7.0	86 95 64 50	2.0 0.0 9.4 9.4	s s n nw	108 66 91 135	14 10 13 18	1 P 8 A	6 P 6 P	.03	.25 1.5
16 17 18 19	20 30 26 19 31	4 P 3 P 0 A 3 P 3 P	-8 18 4 -9 5	5 A 3 A 12 P 7 A 7 A	6.0 24.0 15.0 5.0 18.0	79 86 64 59	4.9 2.0 6.5 8.6 7.4	s nw nw s	18 23 308 67 130	8 4 25 13 10	6 P 0 A	12 P 7 A	.12	1.0 2.5
20 21 22	31 28 18	9 P 3 P 1 P	17 17 17	4 A 12 P 12 P	24.0 22.5 9.5	92 85 54	0.0 2.0 4.0	W S W	80 31 153	12 20 25	3 A	4 P	.48	3.25
21 22 23 24 25 26 27 28 29	8 26 25 27 21 28	8 P 2 P 1 P 1 P 3 P 2 P	-16 5 -8 5 7	8 A 12 P 6 A 1 A 8 A 7 A	-4.0 15.5 8.5 16.0 12.0 17.5	71 78 67 61 40 47	0.0 6.7 5.8 3.6 9.9 4.5	n n nw nw s nw	108 185 38 81 243 145	20 13 25 20 20 15	2 P 0 A	12 P 9 A	.30	3.0 3.0
30 31	14 20	0 A 4 P	=5 -10	8 A 7 A	4.5 5.0	47 41	9.9 9.9	n n w	315 156	28				

^{*} Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

January 1948

January, 1948	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum. 30.53, 24th Minimum. .29.34, 21st, 22nd Mean semi-daily. .30.087	30.70 29.20 30.07 1.50	31.00, 1927 28.55, 1913 2.18, 1913; 0.97, 1896
Range	1.50	2.10, 1713, 0.77, 1070
	51.1	((0 1022
Highest 39, 9th Lowest -16, 24th Mean 18,0 Range 55 Highest mean daily 32,5, 13th Lowest mean daily -4,0 24th Mean maximum 27,1 Mean minimum 8,8 Greatest daily range 35, 29th 12,12 12,12	51.1 7.1 24.2 58.2	66.0, 1932 —26.0, 1904 34.2, 1913; 13.9, 1918
Least daily range	1265 3817	
Precipitation, in inches		
Precipitation 2,63 Snow 20.5 Maximum precip. in 24 hrs. 0.61, 2nd	$\frac{3.61}{13.32}$	7.15, 1898; 1.07, 1896 33, 1898, 1923; 1.50, '08
Number of days with .01 or more	1.1	15, '93, '20; 4, 1901
Wind, in miles		
Total movement4262Greatest daily movement390, 10thLeast daily movement18, 16thMean hourly velocity5.7Maximum velocity32, 9th, 10th	5055 6.8 32.7	7770, 1908; 2896, 1895 47, 1938
Wind, direction		
Prevailing direction NW North, days 7 Northeast, days 2 East, days 1 Southeast, days 6 South, days 6 Southwest, days 0 West, days 3 Northwest, days 11	WXW	
Weather		
Mean relative humidity, percent. 67.3 Mean cloudiness, percent 82 Number of clear days 10 Number of fair days 12 Number of cloudy days 9 Number of hours bright sunshine 150 Percent of possible hours of bright sunshine 51 Snow 20.5	70.1 55 9 13 137 46.7	78, 1932; 37, 1908 20 1944; 2, 1914 18, 1890; 1926;3,1916, 1944 22,1923,1931; 4,1920,1939 214, 1920; 74, 1932

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1947.

REMARKS

The weather for the month of January was consistently cold and was accompanied by a considerable fall of light density snow. The snowfall of 20.5 inches was considerably over the normal 13.32 inches, but the water yield was only 2.63 inches against a normal of 3.61 inches of water. Of the 38.0 inches of snowfall cumulative for the season, 17 inches remained on the ground on the 31st.

The lowest minimum temperature of the month, -16, occurred on the 24th. This was 10 degrees short of the record - 26 of January 1904 and was 5 degrees warmer than the -21 recorded in January 1935. The maximum of 39 degrees recorded on the 9th is the lowest maximum temperature recorded in the month since the beginning of our records. The normal mean maximum for January is 51.1 degrees while the mean maximum recorded for the current month was 27.1 degrees.

The mean temperature for the month at 18.0 degrees was 0.2 degree above January of 1945 but was considerably below the normal mean of 24.2 degrees. Prior to 1945 the most recent severe January was recorded in 1920. The degree-days of heating load resulting from this very low mean built to 1459 for the month and 3932 cumulative for the season, against a normal of 1265 degree-days for the month and a cumulative of 3817. On January 22, the midpoint of the normal heating load, the cumulative degree-days reached 3431 against a normal of 3425.

W. H. Thies, Extension Professor of Horticulture, contributes the following report: "An unusually heavy blanket of snow during January has offered good protection to plant roots and has been highly favorable for fall seedings. Some smothering of new seedings may occur, however, if a layer of ice forms on top of the snow as a result of freezing temperatures during subsequent rains. The heavy snow cover will have another favorable effect in that a major portion of the winter precipitation will be able to penetrate the soil where it falls, since there has been relatively little freezing of the soil. This assumes of course, a moderate rate of melting of the snow.

The temperature in various parts of the State has dropped below the critical point for peach buds (about 15). This will undoubtedly mean some reduction in the 1948 peach crop. Little or no damage to apple buds is apparent thus far. Fruit growers may expect considerable mouse damage in orchards with heavy grass cover, where control measures were not carried out last fall."

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series Bulletin No. 710 February 1948

Meteorological Observations

FOR

FEBRUARY 1948

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

	Temperature				ity	rs		Wind			Precipit	ation		
	Max	ximum	Min	imum		Humid	of Hou	bo	1	es *				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18	18 23 25 38 30 45 27 40 50 45	4 P 5 P 4 A 0 A 3 P 4 P 3 P 10 A 4 P 4 P 3 P 12 P 8 A 3 P 12 P 12 P	5 1 9 3 2 2 7 6 -7 10 10 17 7 25 16 7 36 27	8 A 1 A 12 P 0 A 7 A 12 P 7 A 12 P 5 A 12 P 6 A 12 P 6 A 12 P 12 P 12 P 12 P	15.0 15.5 17.0 6.0 16.0 14.5 11.5 5.5 6.5 7.5 27.5 18.5 35.0 21.5 23.5 43.0 36.0	56 63 45 72 75 56 66 55 53 78 70 83 77 60 67 52 72	10.0 10.0 10.1 0.0 7.2 4.2 5.0 6.2 10.3 2.0 7.2 0.0 4.0 10.5 2.0 6.6 9.6	nw s w nw nw w s nw n n s n w w w	92 21 232 94 155 19 78 258 113 99 42 109 36 318 403 107 241 205	14 5 22 8 13 9 8 40 15 10 10 18 10 48 40 13 28 28	5 P 2 P 0 A	12 P	.20 .17 .76 .14	2.0
19 20 21 22 23 24 25 26 27 28 29	44 44 30 27 28 31 44 37 41 29 31	12 P 0 A 4 P 4 P 5 P 5 P 2 A 3 P 1 P 2 P	19 22	6 A 12 P 7 A 12 P 7 A 1 A 1 A 12 P 6 A 12 P 12 P	32.0 32.0 21.0 21.0 17.0 15.5 34.5 31.5 30.0 25.5 23.5	72 47 51 79 57 74 84 72 88 72	2.0 1.3 10.8 0.6 10.9 6.0 2.3 0.0 6.3 0.0 4.6	nw w nw ne n s sw n n n nw	29 235 88 127 122 74 243 82 84 85 164	3 25 14 13 11 9 25 17 10 12 14	1 P 4 A 0 A	10:30A 4 P 12 P 3 P	.05 .05 .96	1.0 1.0 6.0 2.0

^{*} Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

February, 1948	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.98, 24th Minimum 29.12, 14th Mean semi-daily 30.221 Range 1.86	30.63 29.24 30.04 1.38	31.05, 1920 28.56, 1895 1.89, '00, '08; .88, '13, '31
Air Temperature, in degrees F.		
Highest 50, 17th Lowest -10, 10th, 11th Mean 21.4 Range 60 Highest mean daily 43.0, 17th	50.5 - 7.4 23.7 58	65.0, 1930 —23, 1943 32.6, 1890; 11.6, 1934
Lowest mean daily5.5, 9thMean maximum32.2Mean minimum10.6Greatest daily range37, 7thLeast daily range6, 4thDegree days of heating load for month1264Degree days cumulative for season5196	32.8 14.5	
Precipitation, in inches		
Precipitation 2.45 Snow 15.0 Maximum precip. in 24 hrs. .96, 28th	3.19 13.99	8.12, 1900; 0.62, 1901 48.75, 1893; 0.50, 1937
Number of days with .01 or more	10	15, '93, '20; 4, 1901
Wind, in miles		
Total movement. 3955 Greatest daily movement 403, 15th Least daily movement 19, 6th Mean hourly velocity 5.7 Maximum velocity 48, 14th	7.1 31.7	6445, 1896; 3438, 1892 50, 1946
Wind, direction		
Prevailing direction WNW North, days 6 Northeast, days 1 East, days 0 Southeast, days 2 South, days 4 Southwest, days 1 West, days 7 Northwest, days 8	WXW	
Weather		
Mean relative humidity, percent 65.0 Mean cloudiness percent 82 Number of clear days 10 Number of fair days 9 Number of cloudy days 10 Number of hours bright sunshine 160 Percent of possible hours of bright sunshine 54.1	66,6 50,9 10 8 10 158	66, 190, 127; 31, 1905 19, 1941; 2, 1927 16, 1920; 2, 1936 16, 1894; 2, 1920 221, 1924; 110, 127, 138

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1947.

REMARKS

The weather for the month was near normal in most respects. A below normal minimum barometer on the 14th occurred on the same date as the near record 48-mph maximum wind velocity, and the maximum barometer reading for the month is topped only by the record of 1920. The 15-inch snowfall is only slightly above the normal, 13.99, but the precipitation in water is still below normal at 2.45 inches for the month. The 23 storms of the current season have recorded a total of 53.0 inches of snow to the month's end, considerably above the normal 38.15 inches but not crowding the 71.5 inches of the 1892-1893 record.

The mean minimum at 10.6 degrees is 3.2 degrees below the normal, and the mean temperature for the month settled at 2.3 degrees below normal. This is reflected in the degree days of heating load of 1264 for the month and 5196 for the season against the normals of 1165 and 4982 degree days. Thus the normal heating load would be 72.6% completed at the month's end, and if the current season is to follow the same trend, the season's total load estimates at 7150 degree days against the normal total of 6850, only 4.4% above normal, but 11.7% above last season.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 711

March 1948

Meteorological Observations

FOR

MARCH 1948

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Ter	mpera	ature		ity	rs		Wind	l		Precip	itatio	n
	Max	kimum	Min	imum		Humidity	of Hours inshine	pt.	ıt	es				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative I	Number of Hour Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5	30 28 40 38 20	4 P 3 P 3 P 3 P 3 P	21 25 10 0	7 A 7 A 2 A 12 P 7 A	17.5 24.5 32.5 24.0 10.0	57 88 74 70 38	10.8 0.0 5.6 1.5 11.4	nw nw nw nw	115 94 193 148 256	12 9 15 20 25	6 A 4 A	12 P 8 A	.78 .03	7.75
6 7 8 9	40 34 36 45 38	5 P 2 P 3 P 4 P 6 P	-14 21 31 28 25	7 A 3 A 7 A 7 A 7 A	13.0 27.5 33.5 36.5 31.5	49 88 61 61 84	11.5 0.0 1.8 4.4 1.0	w s nw s	38 42 179 111 30	25 5 5 22 14	7 а	8 p	.42	3,00
11 12 13 14 15	31 34 35 36	1 A 5 P 2 P 5 P	17 8 6 2 23	12 P 7 A 7 A 7 A	24.0 21.0 20.5 19.0 35.5	73 48 52 62 78	0.0 11.7 11.8 4.0	n n w se	151 200 84 91 136	17 17 10 12 12	12 м	10 р	.32	4.00
16 17 18	48 50 47 50	7 P 0 A 5 P	42 31 22	7 A 1 A 12 P 7 A	46.0 39.0 36.0	95 67 51	1.0 0.0 4.4 12.0	se se nw w	224 159 55	18 15 6	0 а	11 A	.32	
19 20 21 22 23	50 59 68 67 44	9 P 5 P 2 P 3 P 2 P 3 P	27 34 29 43 38	4 A 12 P 5 A 5 A 7 A	38.5 46.5 48.5 55.0 41.0	75 58 78 70 77	0.2 12.0 2.0 1.5 0.0	se nw s nw e	116 235 78 150 129	20 28 11 18 11	1 P 0 A 2 P 1 A	12 P 1 A 5 P 3 A	.21 .01 .33 .10	
24 25 26 27 28	65 51 56 59 51	3 P 4 P 1 P 6 P 0 A	35 30 27 51 28	5 A 6 A 12 P 12 P	50.0 40.5 41.5 55.0 39.5	65 43 44 88 57	9.3 12.4 7.4 0.9 0.0	n ne sw s nw	118 195 162 245 290	13 25 14 13 20	6 P	12 Р	.40	
29 30 31	50 62 69	4 P 4 P 3 P	22 31 33	6 A 1 A 7 A	36.0 46.5 51.0	54 61 66	12.6 8.2 12.6	se se s	146 240 162	22 15 22				

^{*} Based on least time required to blow one mile.

FERNAND E. BARTLETT, Observer

March, 1948	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.87, 5th Minimum 29.38, 27th Mean semi-daily 30.087 Range 1.49	30.57 29.24 30.00 1.34	31.05, 1943 28.47, 1914 2.10, 1914: 0.85, 1915
Air Temperature, in degrees F.		
Highest 69, 31st Lowest -14, 6th Mean 34,9 Range 83 Highest mean daily 55.0, 22nd, 27th Lowest mean daily 10.0, 5th	63.8 6.1 34.4 57.6	85, 1929 -7.5, 1906
Mean maximum. 46,2 Mean minimum 23,6 Greatest daily range. 54, 6th Least daily range 5, 8th Degree days of heating load for month 932 Degree days cumulative for season 6128	43.2 25.4	
Precipitation, in inches		
Precipitation 2.92 Snow 15.0 Maximum precip, in 24 hrs. 0.78, 2nd	3.70 7.47	7.89, 1942; 0.12, 1915 27, 1899; 0, 1921
Number of days with .01 or more10	11	17, 1890, 1913, 1936; 3, 1915
Wind, in miles		
Total movement	5944 7.7 31.2	8182, 1896.; 3006, 1905 48, 1932, 1939, 1941, 1942
Wind, direction		, , , , , , , , , , , , , , , , , , , ,
Prevailing direction WNW North, days 3 No:theast, days 1 East, days 4 Southeast, days 5 South, days 5 Southwest, days 1 West, days 4 Northwest, days 10		
Weather		
Mean relative humidity, percent 65.5 Mean cloudiness, percent 85 Number of clear days. 11 Number of fair days 6 Number of cloudy days 14 Number of hours bright sunshine 172 Percent of possible hours of bright sunshine 46.3	64.2 51.8 11 10 10 199 53.6	68, 1901; 27, 1915 22, 1924; 3, 1901 17, 1900; 1, 1943 21, 1951; 1, 1915 292, 1924, 93, 1901

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1947.

${f REMARKS}$

The month of March brought welcome relief from a winter which had provided both prolonged cold and snow cover of above normal. Early in the month however the record March minimum of -14 was established on the 6th, seven degrees colder than the previous record of -7 reached in 1943 and 1906. This coupled with a 69 maximum on the 31st established a record range of temperature of 83 degrees for the normal range is but 57.6 degrees. The tag end of winter also brought 15 inches of snow fall for the month against a normal of 7.47 giving a total for the season to the month's end of 68.0 inches.

The mean temperature for the month of 34.9 degrees was an even one-half degree above the normal. While comfortable day-time temperatures were reached on most days after the 5th, the morning of the 16th recorded the first overnight period of the season without freezing temperatures and the third night with above-freezing temperatures since last November 12. The degree-days of heating load for the month were 932 against a normal of 950 and the season's total of 6128 against a normal of 5932 degree-days. With the normal heating load 86.5% completed by the end of the month, the heating load for this season can be estimated at very little over the normal of 6850 degree days.

W. H. Thies, Extension Professor of Horticulture, contributes the following report: "One of the most surprising features of the weather during March was the very rapid disappearance of a heavy accumulation of snow. And, contrary to fears of many people, this was accomplished without serious flood conditions. Two or three warm days coupled with the absence of frost in the soil made it possible for the earth to absorb the melting snow "like a sponge". Very little soil erosion has occurred and wells have been replenished.

From an agricultural standpoint, the heavy blanket of snow has, in general, been beneficial. The winter precipitation has been permitted to penetrate the earth where it fell instead of flowing into the lowlands as occurs when the soil is deeply frozen. Fall seedings have come through the winter in realtively good condition. Fruit buds, with the exception of peaches which suffered from low temperatures, look promising. Strawberries look unusually good, having been subjected to little or no injury during the winter. Some injury to conifers exposed to drying winds has, however, been noted and there has been some breakage of young fruit trees due to the heavy snow. It is gratifying to see how generally farmers are growing winter rye on areas subject to both wind and water erosion. This is a move in the direction of soil conservation."

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 712

April 1948

Meteorological Observations

FOR

APRIL

1 9 4 8

H. X STAPLETOX

OBSERVATORY

Latitude, 12° 22° 31″ N. Longitude, 12° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft Height of wind instruments. 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

	Temperature					lity	urs e	1	Wind			Precipitat	ion	
	Maxir	num	Mini	mum		Humidity	r of Hours Sunshine	pt	ديد	es.				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative	Number Bright St	Prevailing Direction	Total Movement	Max. Miles per Hour ^a	Веgan	Ended	Water	Snow
1	59	4 A	54	12 P	56.5	89	0.0	SE	250	28	7 A		.76	
2	54	0 A	38	12 P	46.0	71	1.0	NW	133	22	6:30 A		.10	
3	41	1 P	33	12 P	37.0	49	6.0	NW	474	32	8 1	9:30 A		T
4	55	5 P	30	7.5	42.5	31	12.8	N	339	32			ĺ	
5	73	3 P	39	1.4	56.0	39	6.4	8	179	15				
6	65	11 7	43	12 P	54.0	80	2.2	NE	160	13	12 vi	3 Р	.24	1
7	68	3 P	33	6.4	50.5	55	13.0	SE	145	18			!	
S	50	11 P	39	3.1	44.5	82	0.0	8	88	15				1
9	49	0.4	34	12 P	41.5	49	1.2	N.W	367	40				
10	48	3 P		7.	37.5	42	13.1	W	387	28				!
11	49	12 P	23	5.4	36.0	59	2.3	SE	167	17	1 1	10 P	.44	ļ
12	63	5 P	48	12 P	55.5	76	1.6	W.	190	14	2 \	8 A	.32	
13	52	2 P	44	5 1	48.0	44	0.0	NW	137	12				
14	46	5 1	35	12 P	40.5	85	0.0	N	118	9	7.4	12 P	.78	
15	53	2 P	35	2 A	44.0	75	1.9	NW	122	28				
16	59	2 P		6 4	43.5	70	10.3	7.	148	20			Ì	1
17	47	3 7		[12 P]	38.5	37	11.8	N	400	28		į.	1	
18	57	4 P		5.1	40.5	45	13.4	SW	295	13				1
19	56	2 P	34	2 A	45.0	60	0.7	S	195	13	4 P	10:30 P	.03	
20	71	5 P		5.	57.0	65	1.6	8	198	11				
21	62	11 A		12 P	49.5	72	1.4	NE	191	25	1 1	3 7		
22	65	6 P		6 A	47.0	55	13.6	NE	46	14	5.3	6 A	.01	
23	72	3 P		6 A	52.0	48	8.2	S	107	10				i
24	73	3 P		7 A	58.0	47	8.0	S	121	10			,	1
25	69	3 P		12 P	56.0	43	13.8	SE	154	13			-	
26	68	4 P		6.4	50.0	50	13.9	SW	91	17				
27	62	4 P		4 A	47.0	38	13.9	ZW	121	15				
28	54	3 P		4 1	43.5	59	3.4	E	82	7			-	
29	55	5 P		4.1	47.0	60	2.0	SW	68	9			10	
30	60	2 P	40	12 P	50.0	65	1.5	W	183	32	4 P	6 P	.16	1

^{*} Based on least time required to blow one mile.

FERNAND E. BARILLIT. Observer

April, 1948	Normal	Extremes
Barometer, in inches (Readings reduced to sea level) Maximum	30.48 29.38	30.71, 1911 28.98 1943
Mean semi-daily	$\frac{29.99}{1.11}$	1.52, 1930; .72, 1919
Air Temperature, in degrees F. Highest .73 5th, 24th Lowest .23 11th Mean .47.2 Range .50 Highest mean daily .58.0 .24th Lowest mean daily .36.0 .11th Mean maximum .58.5 Mean minimum .35.8	56.8	90, 1941 8.5, 1923 52, 1921; 41, 1943
Greatest daily range	1	
Precipitation, in inches Precipitation2.87	3.35	6.89, 1929; .55, 1941
SnowT. 3rd Maximum precip, in 24 hrs78, 14th Number of days with .01 or more10		11. 1891; 0. 1910, 1934 18. 1909; 3, 1892
Wind in miles		
Total movement		8208. 1908;3853, 191 7
Maximum velocity40, 9th		40, 1935 1938, 1940, 1945
Wind, direction		
Prevailing direction SW North days 3 Northeast, days 3 East, days 1 Southeast, days 4 South, days 7 Southwest, days 3 West, days 3 Northwest, days 6		
Weather	0.0	
Mean relative humidity, percent 58.0 Mean cloudiness, percent 84 Number of clear days 100 Number of fair days 4 Number of cloudy days 16 Number of hours bright sunshine 179 Percent of possible hours of	51.8 11 9 10 220	75, 1901; 34, 1925, 1927 23, 1941; 3, 1898, 1902 18, 1915; 2, 1894, 1901 22, 1901; 1, 1941 329 1941; 103, 1901
bright sunshine45 Thunder and lightningNo. 4, 30th		Mar. 14. 1910; May 11, 1907

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1947.

REMARKS

While the month of April provided considerable cloudy weather and only 179 hours of bright sunshine against a normal of 220, precipitation continued to remain below normal with only 2.87 inches recorded against a normal of 3.35 inches. In spite of the heavy snow cover from which little was lost to flooded streams, moderately drouthy conditions are noticeable. The precipitation for 1948 so far totals 10.87 inches against a normal of 13.85 inches and the total received since last June 1 is but 33.40 inches. This 11-months' total is compared to our calendar year average of 43.70 inches for emphasis. April has next to lowest monthly average precipitation in our records, and it normally should not have been expected to relieve the deficit in rainfall, but a greater burden of providing moisture for growing crops is now placed upon the precipitation which falls during the growing season.

The cloudiness may have contributed to less extremes in temperatures during the month as a range of 50.0 degrees against an average range of 51.4 degrees was recorded. The normal high for the month is 59.4 while in this month 73 degrees was attained on the 5th. The normal minimum is 22 while 23 degrees was recorded the morning of the 11th. The mean daily temperature for the month was 47.2 degrees, 1.5 degrees above the average. This higher mean daily temperature provided a heating load of 536 degree-days for the month against an average of 579, to bring the season's total to 6664 against a normal of 6511 degree-days. With the normal heating season 95% completed by April 30, warm weather through May and early June could make the total for the current season fall very little over average in heating load.

The wind velocity of 40 mph attained on the 9th ties the record for the month attained in 1938 and equalled in 1940 and 1945. There was more wind movement than normal and on 11 days velocities of 20 mph or above were recorded with 4 days recording maximum velocities above 3 mph.

W. H. Thies, Extension Professor of Horticulture, contributes the following report: "Weather conditions during April have been favorable to apple growers in at least two ways. (1) Relatively cool weather has retarded fruit bud development enough to minimize the danger of frost injury during the blossoming season. (2) Continued dry weather has greatly decreased the danger of pre-blossom infections of apple scab. With McIntosh buds now in the pink stage, we have had only one rainy period of sufficient duration to result in scab infection. Tremendous numbers of scab spores are mature, however, and will be released during the next rain.

Present prospects indicate a heavy apple bloom and, in orchards of good location, a moderate peach bloom. No frost injury to fruit buds has apparently occurred during the past month. With favorable conditions for bee activity during bloom and an absence of damaging frosts during the next three weeks, a larger than normal apple crop may be expected. On lighter soils, newly set fruit trees have required liberal watering."

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 713

May 1948

Meteorological Observations

FOR

MAY

1948

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W.

Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

> Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Te	mpera	ture		dity	Hours hine	,	Wind			Precipitat	tion
	Maxi	mum	Mini	imum .		Humidity	of Ho ınshin	be		8 ÷			
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative	Number of Hour Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water
1	56	2 P	39	1 A	47.5	52	8.4	w	264	22	 		
2	69	5 P	31	5 A	50.0	52	14.0	sw	46	8			i i
3	68	3 P	41	5 A	54.5	42	14.1	NE	102	13			1 1
4	72	3 P	39	5 A	55.5	52	13.1	s	118	17			1 1
5	63	3 P	44	6 A	53.5	66	1.8	SE	180	18			ĺĺ
6	71	2 P	43	5 A	57.0	47	14.2	NE	174	15			1 1
7	51	1 P	44	12 P	47.5	88	0.0	E	104	8	1:30 P	12 P	.42
8	51	11 A	43	2 A	47.0	64	0.0	sw	173	18	0 A	8 A	.29
9	71	5 P	45	1 A	58.0	40	10.2	w	242	18			
10	81	1 P	55	0 A	68.0	58	2.9	w	183	28	}	•	.01
11	81	5 P	50	5 A	65.5	66	5.7	SE	61	10			
12	81	2 P	55	6 A	68.0	67	4.8	NW	112	9			
13	62	0 A	49	[12 P]	55.5	87	0.0	N	105	12	5:30 A	3:30 P	.39
14	49	0 A	45	5 A	47.0	82	0.2	N	176	15	6 A	9 P	.11
15	64	6 P	44	4 A	54.0	62	2.8	N	170	28			
16	59	10 A	37	5 A	48.0	83	0.0	SE	62	9	11:30 A	12 P	.98
17	62	2 P	50	0 A	56.0	94	0.0	SE	115	12	0 A	6:30 P	.55
18	67	3 P	52	7 A	58.5	83	2.5	N	103	25	5 A	5 P	.35
19	63	5 P	46	6 A	54.5	43	8.0	N	215	22			
20	63	4 P	43	5 A	53.0	52	3.0	N	135	13			
21	53	5 P	43	2 A	48.0	92	0.0	SE	108	13	4:30 A	12 P	.59
22	58	2 P	42	6 A	50.0	72	0.0	NW	176	25	0.1	5:30 A	.31
23	69	3 P	44	2 A	56.5	72	0.9	Е	106	20	8:30 P	12 P	.11
24	72	2 P	52	5 A	62.0	70	2.2	SE	91	12	0 A	8 A	.03
25	69	7 P	54	6 A	61.5	80	0.1	NW	58	5	6 A	5 P	.53
26	71	6 P	58	4 A	64.5	82	3.3	SE	31	9]12:30 A	2 A	.13
27	80	3 P	51	6 A	65.5	86	7.3	sw	75	20			T
28	80	4 P	54	0 A	67.0	71	8.3	SE	109	8	0.00		4.4
29	79	3 P	60	5 A	69.5	83	5.4	SE	78	8	2:30 P	3 P	.44
30	72	0 A	63	8 A	67.5	93	0.0	sw	70	9	2 A	11 A	.59
31	71	6 P	61	12 P	66.0	71	0.8	E	72	7			T

^{*} Based on least time required to blow one mile.

LEWIS F. WELLS, Observer

May, 1948	Normal	Extremes
May, 1340	Norman	Extremes
Barometer, in inches (Readings reduced to sea level)	30.39 29.51 29.96 0.88	30.62, 1936 29.10, 1938
Air Temperature, in degrees F. Highest .81, 10th, 11th, 12th Lowest .31, 2nd Mean .57.3 Range .50 Highest mean daily .47.0, 8th, 14th Mean maximum .67.0 Mean minimum .47.6 Greatest daily range .38, 2nd Least daily range .4, 14th Degree-days of heating load for month .256		94.5, 1896, 1911 24.0, 1900
Precipitation, in inches Precipitation	3.60	7.44, 1931; .48, 1903 20, 1901, 1945; 5, 1903
Number of days with .01 or more16	14	20, 1001, 1010, 0, 1000
Wind in miles Total movement		5946, 1907; 2180, 1894
Least daily movement31, 26th Mean hourly velocity5.1 Maximum velocity28, 10th, 15th	6.1	45, 1935
Wind, direction Prevailing direction ESE North, days 6 Northeast, days 2 East, days 3 Southeast, days 9 South, days 1 Southwest, days 4 West, days 3 Northwest, days 3		
Weather Mean relative humidity, percent	55.6	70, 1901, 1902; 30, 1923 21, 1944; 0, 1927 17, 1907, 1926; 5, 1923 20, 1927; 2, 1941 350, 1944; 137, 1927

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1947.

REMARKS

The weather for May deviated very little from normal except in cloudiness and rainfall. The precipitation for the month at 5.82 inches exceeded the monthly mean by 2.23 inches bringing the total precipitation for the calendar year to 16.70 inches, 1.05 inches below normal. Even with the 12 months' precipitation since last June 1 at only 37.23 inches, and this season's total below normal, there are now no indications that water is in inadequate supply. With some amount of precipitation on 18 days of the month the extreme cloudiness held the total hours of sunshine to only a little above half the normal for the month.

Degree-days of heating load for the month totaled 256 bringing the season's cumulative to 6920 degree-days.

- W. H. Thies, Extension Professor of Horticulture, contributes the following report: "Weather conditions during May have exerted a direct influence on fruit crops in several ways: (1) With some rain on 18 of the 31 days and with long periods of wet foliage, at least five apple scab infections have occurred where trees were unprotected. McIntosh growers have found it necessary to apply frequent fungidical sprays or dusts to provide protection against these primary infections. (2) Fruit bloom was delayed somewhat by cold weather and during much of the blooming period bees were relatively inactive. In spite of this interference with bee activity, there appears to be a reasonably good set of fruit in many orchards. The effect of nearby pollenizing varieties on the set of McIntosh is especially noticeable. Trees adjacent to a good pollenizer have, in general, prospects for a good crop.
- "(3) Newly planted trees and small fruit plantings are off to a good start since the roots have become well established before hot, dry weather has occurred. This has been a distinct advantage on lighter soils. (4) The cover crop in orchards has benefited in the same way as pastures and hay fields. Where the soil was well fertilized and adequately limed, more mulch material has grown during the month of May than would have been the case with less rainfall. On heavy and poorly drained soils, however, growers have been at a disadvantage from the standpoint of getting mired with the spraying equipment."

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 714

June, 1948

Meteorological Observations

FOR

JUNE

1948

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Te	mpera	ture		lity	Hours hine	Wind			Precipitation			
	Maximum		Minimum			Humidity	of Ho unshin	B	ţ.	les r*				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative	Number of Hou Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	
1	80	3 P	51	5 A	65.5	70	6.8	NE	54	8			T	
2	84	3 P	49	5 A	66.5	57	11.3	NE	65	13		Ĭ		
3	80	3 P	51	5 A	65.5	48	8.5	SE	85	15			[[
4	83	4 P	51	6 A	67.0	49	9.3	s	82	13				
5	65	6 P	54	12 P	59.5	83	2.8	NW	129	17	5:30 A	2 P	1.27	
6	72	6 P	41	6 A	56.5	48	9.4	W	105	17]]	
7	59	0 A	48	[6 A]	53.5	94	0.0	NW	59	9	7 A	10 P	1.16	
8	66	5 P	54	4 A	60.0	90	0.1	NE	98	12	0 A	11 A	.03	
9	59	6 P	53	6 A	56.0	84	0.0	NE	95	8	2 A	1 P	.24	
10	65	7 P	50	5 A	57.5	91	2.5	SE	58	4	1 A	2 A	.02	
11	79	6 P	55	7 A	67.0	77	5.9	SE	44	12	5:30 A		.01	
12	76	5 P	55	4 A	65.5	79	5.0	SE	50	14				
13	60	7 P	53	7 A	56.5	93	0.0	N	64	6	L 0	11:30 A	1.19	
14	79	5 P	52	7 A	65.5	65	10.0	sw	68	18			T	
15	75	6 P	58	6 1	66.5	80	1.3	s	117	6	5:45 P	6:15 P	.10	
16	71	4 P	54	6 A	62.5	70	6.3	NE	50	4	5:45 P	6 P	.12	
17	71	3 P	49	6 A	60.0	62	8.7	NW	78	18				
18	76	5 P	44	6 A	60.0	56	10.6	NW	64	8				
19	68	3 P	57	7 A	62.5	84	2.2	NW	104	14	5:30 A	10 A	.44	
20	78	5 P	52	5 1	65.0	55	12.2	W	64	8			T	
21	80	3 P	54	4 1	67.0	51	9.8	NW	116	26				
22	75	2 P	49	5 A	62.0	56	7.3	SW	42	5				
23	62	0.4	57	12 A	59.5	86	0.0	SE	100	9	10 A	8 P	.19	
24	85	3 P	55	0.4	70.0	89	4.5	SE	73	32	4 P	12 P	.58	
25	84	4 P	65	12 P	74.5	59	12.4	11.	103	25	0 A	5 A	.09	
26	81	3 P	59	6 7	70.0	68	9.4	s	45	9				
27	75	2 P	60	5 A	67.5	77	2.7	S	87	7] _]	
28	87	3 P	63	2 A	75.0	85	4.9	s	74	9	1:30 A	6:30 P	.20	
29	93	2 P	60	4 A	76.5	77	7.3	s	58	17	3 A	6 A	.03	
30	91	4 P	70	5 A	80.5	67	9.8	W	111	13			T	

^{*} Based on least time required to blow one mile.

LEWIS F. WELLS, Observer

JUNE, 1948	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)	30.30 29.55 29.95 0.75	30.70, 1947 29.24, 1902
Air Temperature, in degrees F.		
Highest 93, 29th Lowest 41, 6th Mean 64.7 Range 52 Highest mean daily 80.5, 30th Lowest mean daily 53.5, 7th Mean maximum 75.3 Mean minimum 54.1 Greatest daily range 35, 2nd Least daily range 5, 23rd Degree-days of heating load for month.79 Degree-days cumulative for season6999	91.2 40.0 65.7 51.3	101, 1919 34, 1891
Precipitation, in inches		
Precipitation	3.75	9.68, 1922; 0.76, 1908
Number of days with .01 or more15	11	17, 1922, 1945; 4, 1 908
Wind in miles		
Total movement 2342 Greatest daily movement 129, 5th Least daily movement 42, 22nd Mean hourly velocity 3.3 Maximum velocity 32, 24th	3585 5.0 24.6	4571, 1908; 1409, 1906 48, 1939
Wind, direction		
Prevailing direction SW North, days 1 Northeast, days 5 East, days 0 Southeast, days 6 South days 6 South west, days 2 West, days 4 Northwest, days 6	WSW	
Weather		
Mean relative humidity, percent	66.9 51.1 10 12 8 257 54.0	71, 1903; 28, 1908 22, 1908, 1941,1943; 1, 1927 23, 1912; 3, 1941 22, 1903; 1, 1923 362, 1908; 102, 1903

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1947.

REMARKS

It not only seemed to rain more than normal in the month of June, it provided the heaviest precipitation for the month since 1941. In 9 other years since the beginning of our records however the 5.67 inches received this month has been exceeded, and it was considerably below the record 9.68 inches of June, 1922.

Accompanying this greater than normal rainfall were the expected less than normal sunshine, more than normal cloudiness, and less than normal mean temperature. While none of these were of a magnitude to approach a record value, the condition is worthy of note for its effect on farming operations. The harvesting and curing of forage crops was very difficult due to precipitation, low temperature, high humidity and soft ground conditions. Planting of some crops was delayed greatly due to soft fields and the "continuous" rain.

Degree-days of heat load for the month were 79 against a normal of 68 and the season's cumulative load was 6999 against a normal of 6848 degree-days.

W. H. Thies, Extension Professor of Horticulture, contributes the following: "With rainfall amounting to 5.67 inches during June, distributed over 19 days, conditions have been very favorable for the continued spread of apple scab and other fruit diseases. Wherever trees are in good vigor, fruits of various kinds have sized rapidly although some yellowing of foliage in areas of relatively high water table has appeared. In strawberry plantings the yield has been somewhat reduced through smaller sized berries where plants were handicapped by root rot, leaf spot, or other unfavorable conditions.

Much of the soluble plant food has been leached from the soils in the Connecticut Valley as a result of continuous rains during the past two months, according to W. G. Colby of the Agronomy Department. Magnesium deficiencies have appeared on fields of potatoes, and nitrogen deficiencies have been evident on both potatoes and onions. Magnesium deficiency symptoms have been aggravated, not only because of the leaching action of heavy rains, but also because of shalllow root penetration which is characteristic of a wet growing season."

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 715

July 1948

Meteorological Observations

FOR

JULY

1948

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W.

Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

> Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Ter	npe ra	ture		dity	ours	Wind			Precipitation			
	Maxi	mum	Mini	mum		Humidity	of Ho unshir	20	ىد	les r*				
Day	Degrees	Time	Degrees	Time	Меан	Mean Relative	Mean Relative Humidity Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Веgan		Ended	Water
1	85	4 P		6 A	77.0	80	6.5	N	37		1	1.	5 1	.21
2	80	4 P	59	3 A	69.5	56	12.5	N	113	14				
3	54	5 P	52	6 A	68.0	59	12.3	E	27	2				
4	86	5 P	54	5 A	70.0	51	12.5	N	42	14			10	0.2
5	\$9	4 P	55	6 A	72.0	69	8.2	sw	34	32		9 Y	12 P 4 P	.93
$\frac{6}{7}$.	85	2 P	62	12 P	73.5	75	7.0	S	65	$\frac{28}{20}$		11 .1	† P	T
	75 83	3 P 5 P	$\frac{56}{52}$	5 A 6 A	$\begin{array}{c} 65.5 \\ 67.5 \end{array}$	$\frac{60}{58}$	$11.0 \\ 12.5$	NW	158 31	20				1
- S - 9	- 86 - 86	э Р 3 Р	52	6 A	69.0	58 58	11.4	S	29	6				
		_	$\frac{52}{59}$	6 A	73.5	57	$11.4 \\ 11.2$	E E	92	14				
$\frac{10}{11}$	$\frac{88}{91}$	5 P 4 P	62	6.4	$\frac{15.5}{76.5}$	62	9.3	ZW	95	12				
$\frac{11}{12}$	94	3 P	68	6 A	81.0	67	12.0	SE	84	8	1			
13	82	12 N	$\frac{63}{70}$	12 P	76.0	83	0.5	SL	103	9		0.4	11:30 г	$1.02^{^{1}}$
14	81	5 P	63	12 P	72.0	63	11.5	N	138	14				
15	83	5 P	55	6 A	69.0	61	12.4	ΣW	64	6				
16	85	4 P	55	6 A	70.0	65	8.8	s	61	8				
17	82	1 P	61	1 6 A	71.5	66	8.1	SE	122	11				
18	88	2 P	62	5 4	75.0	69	8.8	s	66	8	i			
19	83	4 P	63	6 A	73.0	65	8.0	NW	73	12	1			
20	84	6 P	59	: 5 1	71.5	57	12.3	sw	47	8	!			
21	85	4 P	64	6 A	74.5	73	4.2	SE	59	9				
22	84	3 P	64	5 A	74.0	52	10.6	N	97	11				
23	74	2 P	60	12 P	67.0	77	0.0	Е	98	12		3 P		.21
24	79	2 P	57	12 P	68.0	67	8.8	E	97	15	1	8 4		T
25	81	4 P	52	4 A	66.5	58	11.0	NW	99	12		٠	10 -	14
26	81	1 P		3 A	68.5	66	5.0	s	95	10	11:3		12 P	.14
27	86	2 P		4 1	75.5	83	4.0	SE	77	20		8.4	8:30 A	.01
28	87	2 P		12 P	76.5	68	8.4	W	98	17				1
29	86	2 P		4 A	73.5	56	11.3	NW	69	10				
30	90	2 P		4 A	75.0	61	9.9	s	70	11				1
31	87	_	74	5 A	80.5	59	12.0	M_{\perp}	184	18				

^{*} Based on least time required to blow one mile.

LEWIS F. WELLS, Observer

JULY, 1948	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum30.10, 9th	30.27	30.50, 1892
Minimum29.48, 6th	29.59	29.27, 1932
Mean semi-daily29.815	29.96	
Range	0.68	0.97, 1892; 0.47, 1938
Air Temperature, in degrees F.		
Highest94, 12th	93.9	104, 1911
Lowest52, 3rd, 8th, 9th, 25th	46.4	40, 1890, 1898
Mean72.3	70.8	74.7, 1921; 66.3, 1891
Range42	47.7	
Highest mean daily81.0, 12th		
Lowest mean daily65.5, 7th		
Mean maximum84.3		
Mean minimum60.2		
Greatest daily range34, 5th, 9th		
Least daily range12. 13th		
Precipitation, in inches	(1.)	
Precipitation2.95	4.10	14.51, 1897; 0.70, 1929
Maximum precip. in 24 hrs1.02, 13th		20 1017 4 1004
Number of days with .01 or more7	11	20, 1915; 4, 1924
Wind in miles	2.22	
Total movement2524	3422	5097, 1909; 1109, 1894
Greatest daily movement184. 31st		
Least daily movement27, 3rd	4.3	
Wean hourly velocity	4.6	44, 1936
Maximum velocity32, 5th	28.7	11, 1330
Wind, direction		
Prevailing directionSW	SW	
North, days6		
Northeast, days0		
East, days4		
Southeast, days4		
South, days7		
Southwest, days2		
West, days2		
Northwest, days6		
Weather		
Mean relative humidity, percent62.6	68.4	
Mean cloudiness, percent38	50.9	70, 1943; 31, 1924
Number of clear days15	10	22, 1923; 0, 1915
Number of fair days12	1-1	24, '09, '21; 4, 1946
Number of cloudy days4	ī	18, 1889; 0, 1910
Number of hours bright sunshine282	268	371, 1910; 180, 1931
Percent of possible hours of		
bright sunshine61	58	
Thunder and lightning		
5th, 6th, 13th, 23rd, 27th		

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1947.

REMARKS

The weather for July was a little warmer than normal. The high of 94 degrees recorded on the 12th is but 0.1 degree above the normal maximum, the lowest temperature recorded for the month was 52 degrees the morning of the 25th, while the mean temperature for the month became 12.3 against a normal of 10.8 degrees. It will be noted that this increase in mean temperature was accompanied by an above-normal hours of sunshine and a below-normal mean relative humidity. The result was pleasant weather most of the days of the month.

Rainfall of 2.95 inches compares favorably with 2.73 for July of last year but is 1.15 inches below normal for the month. Total precipitation for the calendar year now stands at 25.32 against a normal of 25.30, with a drouth condition ready to assert itself on most of the lighter soils.

W. H. Thies. Extension Professor of Horticulture, contributes the following: "With less than three inches of rainfall and relatively hot weather in Amherst during July, conditions have been favorable for a build-up of certain orchard pests, including European red mite. Bronzing of foliage is becoming more apparent here and there, and 50 or more mites per leaf have been counted. There has been little if any spread of apple scab on the frint, however, since we have had no lengthy rainy period during the month.

"The hay harvest in Massachusetts got off to an early start and has apparently been completed somewhat earlier than usual. Hay was, in general, heavy and of good quality. Corn and potatoes have made excellent growth during the month although some fields show the effects of earlier leaching of essential elements from the soil. Lawns are in fairly good condition, except on lighter soils where drought symptoms are showing. And, as is usual at this season, some gardeners are worried about the rapid growth of weeds forgetting, perhaps, that garden crops and weeds are subject to exactly the same soil and weather conditions."

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 716

August, 1948

Meteorological Observations

FOR

AUGUST

1948

H N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N.

Longitude, 72° 31′ 48″ W

Height of barometer above ground, 36 ft. Above sea level, 252.5 ft Height of wind instruments, 67 feet. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

	1	Ter	npera	ture		lity urs e		Wind			Precipitation			
	Maximum		Minimum			Humic	of Ho unshin	۶t	4	se*			1	
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	
1	83	$\widetilde{3}$ P	58	5 A	70.5	$7\bar{2}$	$+\bar{7}.\bar{7}$	SE	57	9		1	-	
2	84	5 P	62	12 P	73.0	72	6.9	s	95	9				
3	84	3 P	57	6.4	70.5	61	7.5	SE	39	8				
1	82	4 P	61	6 A	71.5	65	6.1	NW	54	10				
5	80	1 P	63	12 P	71.5	66	5.3	NE	94	10		12:00 P	.47	
6	71	3 P	56	12 P	63.5	86	$^{2.0}$	NW	133	18	0 A	1:30 Р	.13	
7	72	4 P	51	5 A	61.5	55	8.9	N	109	14	1		T	
8	78	3 P	51	6 A	64.5	59	s.s	N	94	11			i	
9	78	5 P	50	5 A	64.0	63	7.8	NE	34	9		i		
0	80	4 P	49	6 A	64.5	61	9.5	NE	33	5	1		T	
1	83	1 P	50	6 A	66.5	62	8.9	s	65	10	1			
2	. 73	2 P	66	2 A	69.5	93	0.0		143	25	12:30 A	4:15 Р	1.84	
3	75	11 A	62	12 P	68.5	95	0.0	\cdot s	68	32	11:00 A	5:30 Р	.94	
i	77	3 P	58	3 1	67.5	73	4.0	w	95	20	6:00 \	6:30 A	.01	
5	84	2 P	55	6 A	69.5	59	11.6	NW	79	14			T	
3	84	3 P	52	6 A	68.0	66	10.0	W	25	4			1	
7	82	1 P	57	6 A	69.5	70	5.0	E	36	6			T	
8	79	4 P	53	6 A	66.0	69	9.4	SE	39	5	3		Т	
9	74	5 P	58	5 A	66.0	84	0.3	E		4				
0	78	5 P	65	6 A	71.5	82	1.8	SE	41	3				
1	85	4 P	65	3 7	75.0	7.7	8.6	S	31	6	1	}		
2	85	2 P	61	5 A	73.0	70	8.3	Е	39	4				
3	87	3 P	62	5 A	74.5	73	9.2	E	16	7				
4	. 88	4 P	63	2Λ	75.5	7.9	6.8	SE	59	10			T	
5	96	2 P	68	2 A	82.0	69	9.5	NE	107	9			1	
6	100	3 P	69	6 1	84.5	66	10.8	N	11	4	1	1		
7	97	4 P	69	6.4	83.0	- 67	10.3	S	18	11			1	
8	97	2 P	68	1 6 A	82.5	72	7.8	E	13	7		1		
9	89	1 P	72	6 A	80.5	58	9.0	11.	75	22	0.00	7.1-		
0	86	3 P		6 A		72	7.1	NW	1	İ	6:00 P	7:15 ±		
1	79	1 P	61	6 A	70.0	53	7.7	N	i	1	1	İ	T	

^{*} Based on least time required to blow one mile.

LEWIS F. WEITS. Observer

Barometer, in inches (Readings reduced to sea level)		
(Readings reduced to sea level)		
Maximum30.16, 23rd 30.32 36	.50.	1934
	3.87.	
Mean semi-daily		
Range0.52 0.71		
Air Temperature, in degrees F.		
Highest	100,	1918
Lowest	34,	1940
	2.4,	1903
Range51.0 48.1		
Highest mean daily84.5, 26th		
Lowest mean daily61.5. 7th		
Mean maximum82.9		
Mean minimum59.9		
Greatest daily range33, 11th		
Least daily range		
Degree-days of heating load for month		
Degree-days camulative for season		
Precipitation, in inches		
Precipitation	31.	1894
Snow		
Maximum precip. in 24 hrs1.84, 12th		
Number of days with .01 or more	; 4,	1899
Wind in miles		
Total movement	J20,	1891
Greatest daily movement143, 12th		
Least daily movement25, 16th		
Mean hourly velocity4.6 4.2		
Maximum velocity32, 13th 22.7	40,	1941
Wind, direction		
Prevailing directionE SW		
North, days4		
Northeast, days4		
East, days5		
Southeast, days5		
South, days6		
Southwest, days		
West. days3		
Northwest, days4		
Weather		
Mean relative humidity, percent		
Mean cloudiness, percent32 49.7 67, 1901, 1946;		
Number of clear days		
Number of fair days20 13 25, 1912;		
AT 1 0 1 1 1 1 0 0 10 10 0 10 0 10 0 10		
Number of cloudy days	915	1929
Number of cloudy days	010.	
Number of hours bright sunshine216.6 237 332, 1941; 152. 19 Percent of possible hours of	010.	
Number of hours bright sunshine216.6 237 332, 1941; 152. 19 Percent of possible hours of bright sunshine50.5 55.2	010,	
Number of hours bright sunshine216.6 237 332, 1941; 152. 19 Percent of possible hours of	010.	

The weather for the month of August ranged from warm to hot, with a heat wave the last ten days of the month. From the 21st to the 30th the daily maximums were all 85 degrees or higher; from the 25th to the 28th they were in excess of 95, and the 100-degree temperature recorded on the 26th equals the previous August record on August 7, 1918. There were 25 clear and fair days coupled with average relative humidity. The six cloudy days were the same six which gave some recorded rainfall. Precipitation was 0.68 inch below the monthly mean of 4.24 inches, with more than half of the rainfall in the 1.84-inch storm on the 12th. The moisture condition was slightly more favorable than in 1947 as this one storm recorded more precipitation than the entire month a year ago.

The extremely high temperatures late in the month caused severe losses in poultry flocks not protected with fan ventilation. Birds on range vacated black-roofed shelters, preferring the heat of direct sunshine to suffocation in the shade, Scald developed in some cranberry bogs which appeared to affect the uprights as well as the fruit.

W. H. Thies, Extension Professor of Horticulture, contributes the following: "Weather conditions during August have affected the apple crop in several ways. Extremely high temperatures in late August have probably influenced the amount of sun scald on the fruit ranging up to 20 percent or more in some orchards. similar condition has also been noted on tomatoes.) Relatively warm nights have not favored color development. An absence of prolonged wet periods has meant freedom from late season scab infections in spite of frequent heavy infections in early summer, but considerable russeting of the fruit is in evidence. This is of common occurrence where the relative humidity is high in contrast to an arid climate. In reality, the heavy russeting of some varieties, including Baldwin, may be due to a combination of spray materials and humid weather, particularly in early summer. The average size of apples is quite satisfactory except on drier soils or where the foliage has suffered severe injury from insects or diseases."

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 717

September, 1948

Meteorological Observations

FOR

SEPTEMBER

1948

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 252.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

	Temperature				dity	urs	1	Wind			Precipita	tion	
	Maxi	mum	Mini	imum		Humidity	r of Hours Sunshine	ρū	نيد	* es			1
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative	Number Bright S	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water
1	75	3 Р	48	7.4	65.5	55	11.2	NW	T	14			
2	75	2 P	50	1 A	62.5	72	4.4	E	16	3			
	79	4 P	49	6 A	64.0	70	5.1	SE	48	6			
4	80	3 P	59	7 A	69.5	79	4.6	S	59	5	4:35 P	4:40 P	.01
5	86	4 P	58	3 A	72.0	71	7.4	S	31	6			T
6	88	2 P	61	3 A	74.5	74	6.5	7.	60	6			Т
7	85	4 P	60	3 A	72.5	73	7.1	> W	54	5			
8	86	4 P	57	7 A	71.5	78	7.3	8W	38	9			1 00
9	87	3 P	46	6 A	66.5	77	7.3	>W	15	9	7:15 P	10:45 P	1.02
10	85	3 P	58	6 A	71.5	77	1.3	ΣW	64	10			
11	91	3 P	45	3 A	68.0		7.9	se	31	9			
12	86	10 A	52	0 A	69.0	71	7.8	SE	57	10		1	
13	89	11 A	45	2 A	67.0	75	8.2	S	62	10		! !	
14	72	10 A	60	6.4	66.0	72	10.2	NW	110	12		1	
15	71		45	6 A	58.0	1 -	10.2	NW	63	8			
16	67	5 P	40	6 A	53.5	60	10.3	NW	85	15			
17	72	2 P	34	6 A	53.0	54	5.3	81.	76	12	0.00	4 00	10
18	88	2 P	58	6 A	73.0	76	4.0	81.	105	1 40	3:00 P	4:30 P	.16
1^{α}	79	3 P	57	$= \overline{4} \cdot \Lambda^{3}$	68.0	79	8.3	>W	46	9		40.00	T
20	70	5 P	58	4 1	64.0	93	0.0	N		17	8:00 A	10:00 A	.44
21	67	2 P	46	6 A	56.5	62	8.5	\	131	17			T
22	68	2 P	35	5 A	51.5	64	8.8	`	72	9			
23	66	1 P	38	7 A	52.0	67	4.6	NE	77	12		Ì	
24	66	4 P	45	7 A	55.5	60	9.3	NE.	153	15			
25	72	3 P	44	7 A	58.0	62	9.8	ZM	99	15			
26	75	3 P	40	6 A	57.5	67	9.3	11.	23	5			70
27	81	3 P	36	6 A	58.5	62	9.8	W	23	5		}	Т
28	83	3 P	4.4	6.4	63.5	$6\overline{0}$	9.3	S	27	9			1
29	82	3 P	47	6 A	64.5	64	8.9	SW	22	7		10.00	9.0
30	72	4 P	55	6 A	63.5	81	0.3	E	54	13	5:45 P	12:00 P	.28

^{*} Based on least time required to blow one mile.

LEWIS F. WELLS, Observer

SEPTEMBER, 1948	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum30.53, 17th	30.45	30.65, 1924
Minimum29.15, 29th	29.57	28.41, 1938
Mean semi-daily30.081	30.00	
Range	.88	1.99, 1938; 57, 1910
Air Temperature, in degrees F.		
Highest91, 11th	87.7	97, 1895, 1929
Lowest34, 17th	33.2	24.5, 1914
Mean63.7	61.7	
Range57	54.5	
Highest mean daily74.5, 6th		
Lowest mean daily51.5, 22nd		
Mean maximum78.1		
Mean minimum49		
Greatest daily range46, 11th		
Least daily range12, 14th, 20th		
Degree-days of heating load for month 104		
Degree—days cumulative for season 104		
Precipitation, in inches	4.04	1 (55 1090 , 59 1014
Precipitation	4.24	14.55, 1938; 52, 1914
Maximum precip. in 24 hrs 1.02, 9th		
Number of days with .01 or more5	10	16, 1933; 3, 1903
	10	10, 1000, 0, 1000
Wind in miles		1 000 1000, 1 111 1001
Total movement	3271	4,686, 1896; 1,414, 1894
Greatest daily movement153, 24th		
Least daily movement22, 29th	4.5	
Mean hourly velocity	4.5	80, 1938
Maximum velocity40, 18th	25.9	00, 1000
Wind, direction Prevailing directionSW		
North, days	WSW	
Northeast, days		
East, days		
Southeast, days		
South, days		
Southwest, days4		
West, days		
Northwest, days		
Weather Mean relative humidity, percent69.8	73.6	
Mean cloudiness, percent27.6	43.9	70, 1934; 23, 1947
Number of clear days	10	19, 1932; 2, 1907, 1928
Number of fair days	10	19, 1908; 3, 1889
Number of cloudy days3	10	20, 1934; 3, 1941
Number of hours bright sunshine213	201	255, 1916; 106, 1934
Percent of possible hours of	- (, T	
bright sunshine57.1	54.9	
Thunder and lightning9th		
First frost	Sent 91	Oct 13 1909 · Aug* 22 1894-95

The weather for September continued fair, warm and dry. The 1.91 inches of precipitation was the least since 1940, and more than half was recorded by the storm of the 9th. While measurable rainfall occurred on 5 days, only 3 days are recorded as cloudy, the remainder being fair or clear. The mean temperature for the month was 2 degrees above normal at 63.7, and only about 2/3 the normal heating load accumulated in the month. The highest temperature, 91 degrees on the 11th, is the maximum for the month recorded 5 times in the past 10 years. The lowest temperature, 34 degrees on the 17th, did not produce frost at the station, although higher elevations to the east received their first frost of the season, killing most gardens and susceptible field crops. There were no high winds during the month.

W. H. Thies, Extension Professor of Horticulture, contributes the following report: "A relatively warm, dry September has shown some interesting effects on farm and garden crops. In Amherst, the official low temperature for the month was 34 degrees, and even in lower areas frost damage was light. This made it possible to harvest undamaged tomatoes, green beans, grapes, etc., as late as October 1. Another feature of the September weather was its dryness, with less than half the normal rainfall during the month. This has given farmers a good opportunity to harvest corn, potatoes, fruit crops, and on some farms, a late cutting of alfalfa.

Apples have been rather slow in coloring although they have hung on the trees fairly well. In fact, a considerable percentage of the McIntosh crop was harvested during the last week in September. This is about a week later than in some seasons past. Apple growers will find it necessary to maintain high humidity in storage for at least two reasons. Boxes were very dry at harvest time which means that the wood will absorb water from the apples. Furthermore, there was more russetting of fruit this season and as a result more shriveling may be expected unless a relative humidity of at least 90% is maintained."

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series Bulletin No. 718

October, 1948

Meteorological Observations

FOR

OCTOBER

1948

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 252.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

	Tempera					lity	Hours hine		Wind		1	Precipit	ation
	Maxi	imum	Min	imum		Humidity	of Ho unshin	žį.	4				;
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative	Number of Hour Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Ведип	Ended	Water
ī	79	3 P	57	L 6	68.0	84	5.8	//. 	27	S	()	v 1:30 v	.37
2	77	4 r	53	7 A	65.0	67	6.8		83	. 14			
3	61	2 P	47	7.1	54.0	53	8.8	N	170	22			
4	62	3 P	32	6.4	47.0	52	9.1	NE	103	13			
õ	59	3 P	35	6.4	47.0	61	0	".	53	8			
6	71	4 P	43	, 6 A	57.0	- 63	2.1	.\	98	14			
4	71	2 P	42	4.4	56.5	59	5.0	.\	107	15			4.0
	64	7 P	49	1 1	56.5	93	0	Z.	112	17			.43
9	66	4 P	50	6 A.	58.0	67	7.3	11.	73	10			Γ
()	65	3 P	45	6.4	55.0	65	2.8	8	99	13			4 - 1
1	63	4 P	51 57	6 A	57.0	60	()	E	47	8			.15
9	70	1 P		6 A	60.5	81	0.5	8 E	50	7			T
3	$\frac{62}{56}$	3 P	13	1 1	$52.5 \\ 51.0$	68	2.8 0.1	11.	56	14			Т
i 5	- 63	2 P 3 P	46 45	5 1				11.	47	20			Т
1	- 66 - 66		29	$\frac{4}{7}\frac{\Lambda}{\Lambda}$	$\frac{54.0}{47.5}$	$\begin{pmatrix} 61 \\ 66 \end{pmatrix}$	$\frac{8.4}{7.2}$	ZM	$\frac{154}{70}$	$\frac{20}{13}$			Т
-	74		52		63.0	1.71	3.6	8	199	$\frac{15}{20}$			1
έ,	48	2 P 3 P	38	10 A	43.0	85	0.8	S NE	185	17			.10
}	51	5 P	23	6 A	37.0	73	1.0	Nr.	15	4	2:15	е 2:30 р	
)	49	- 2 P	32	1.0	40.5	84	1.3	N	12	13	~ . I · ·	2,0001	1.00
1	50	3 p	31	6 A	40.5	61	7.3	N.	99	14			Т
2	54	3 P	21	6.1	37.5	61	7.1	ΣM	22	6			T
3	54	2 p	29	; 6 A	41.5	73	0,3	NE	43	20			T
1	53	4 P	42	7 1	47.5	61	()	N	221	20	6:00	a 9:00 a	.01
5	52	3 P	47	8.1	49.5	81	0	N	302	22	6;00		
Ď.	68	2 p	48	1 6 1	58.0	49	5.5		292	22			1
7	66	3 P	35	7 A	50.5	53	7.8	XW	107	12			1
ŝ	68	3 P	30	7. 1	49.0	62	6.6	SE	16	3			1 1
9	61	4 P	34	171	47.5	65	5.6	XW	97	17			1
0	65	4 P	30	8.4	47.5	60	6.8	N	47	14	6:00	A; 8:00 A	Т
1	72	2 P	29	6 4	50.5	60	6.4	N	19	7		8:00 A	T

^{*} Based on least time required to blow one mile.

LEWIS F. WELLS, Observer

OCTOBER 1948	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum30.65, 22nd	30,53	30.68, 1929
Minimum29.68, 9th	29.42	29.00, 1926
Mean semi-daily30.117	30.06	
Range0.97	1.11	1.47, 1926, .76, 1899
Air Temperature, in degrees F.	50 4	0 = 400
Highest	79.4.	90.5. 1908
Lowest21, 22nd	23.4	57.4, 1947, 17, 1936
Mean51.3	50.5	
Range58	56.0	
Highest mean daily68. 1st		
Lowest mean daily37, 19th		
Mean maximum		
Mean minimum		
Greatest daily range43, 31st		
Least daily range		
Degree-days of heating load for month429		
Degree-days cumulative for season533		
Precipitation, in inches		
Precipitation1.13	3.29	8 81, 1911; .01, 1924
Snow		
Maximum precip in 24 hrs0.43, 8th		
Number of days with .01 or more7	9	15, 1913; 1, 1897, 1924
Wind in miles	40=4	* 40* 4040. 0 * 40.04
Total movement3025	4074	5.467, 1910; 2,540, 1894
Greatest daily movement302, 25th		
Least daily movement12, 20th	- 1	
Mean hourly velocity4.07	5.4	10 1027
Maximum velocity22, 3rd, 25th, 26th	29.5	42, 1937
Wind, direction	11.	
Prevailing directionN	* * *	
North, days14		
Northeast, days3		
East, days1		
Southeast, days2		
South, days		
Southwest, days		
West, days		
Northwest, days		•
Weather	30.0	
Mean relative humidity, percent66.8	69.0	0= 1000 10 1001
Mean cloudiness, percent40	18	67, 1932, 18, 1924
Number of clear days4	10	21, 1938; 1, 1911
Number of fair days	10	17, 1924; 3, 1938
Number of cloudy days14	175.0	19, 1896; 3, 1924
Number of hours bright sunshine126.8	175.0	232, '23, '38; 91, 1913
Percent of possible hours of bright sunshine34	51.9	
Thunder and lightning	51.3	
First frost	Sout 21	Now 20101105 - O-1 15100
THE HUST	Sept. 21/2	rug. 22, 94, 95; Oct. 15, 39

The weather for the month of October was very near the normal excepting the amount of precipitation. Of the 1.13 inches received, 0.43 inches was received by the storm of the 8th and the drouthy condition beginning in July has not been adequately relieved. Since the storm of July 13 only 6.96 inches of water has been received and only 3 storms recorded precipitation in excess of 0.5 inches. Since the normal precipitation July through October is 15.11 inches, there is a reason that streams and wells are observed to be very low.

W. H. Thies, Extension Professor of Horticulture, contributes the following report: "In general, October weather was very favorable for farm work. There was little more than an inch of rainfall in Western Massachusetts and the mean temperature was slightly above normal. On one occasion the temperature dropped to 21° causing some worry among apple growers with fruit still on the trees. A few farmers experienced a water shortage. If the soil becomes thoroughly wet before it freezes, conifers will tend to suffer less drying from winter winds.

Winter cover crops and new seedings look unusually well. Aside from the relatively mild weather this may be due, according to Dr. W. G. Colby of the Agronomy Department, to the fact that there has been so little rain to leach the nitrates from the soil. Consequently, cover crops have had a better supply of nitrogen and are therefore making better growth than would have been the case if heavy rains had fallen during October."

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 719

November 1948

Meteorological Observations

FOR

NOVEMBER

1948

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 252.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Te	mpera	ture		midity Hours hine		Wind				Precipitation			
	Maxi	Maximum		um Minimum		of uns]	80		les r*						
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative	Number Bright S	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow	
1	60	2 P	43	7 A	51.5	59	3.9	SE	66	11	<u> </u>				
2	56	3 P	28	7 A	42.0	65	4.9	SE	39	7	1				
3	54	4 P	25	7 A	38.5	76	0.5	N	31	7	7:45 A		.06		
4	59	8 P	47	0 A	53.0	89	0	s	45	8	11:45 A	11:45 P	1.27		
5	70	2 P	57	7 A	63.5	78	2.1	s	65	13			T		
6	72	2 P	61	2 A	66.5	80	0	8	206	32			Т		
7	64	3 P	54	7 A	59.0	61	2.0	W	263	22	4:00 A	10:00 A	.52		
8	63	3 P	40	5 A	51.5	54	1.2	SE	90	\int 6	}	}	T		
9	61	2 P	33	4 A	47.0	80	1.9	SE	53	13					
10	70	6 A	59	0 A	64.5	89	0	SE	202	25	8:30 A]11:30 P	.89		
11	54	3 P	40	7 A	47.0	55	3.3	NW	191	18	Į .		_		
12	51	2 A	29	6 A	40.0		0	E	45	10	1		T		
13	54	3 P	46	0 A	50.0	75	0	W	154	48	4:30 P	8:00P	.02		
14	51	2 P	35	7 A	43.0	48	4.5	NW	240	13			T		
15	50	3 P	33	7 A	41.5	70	0.3	NW	59	12			T		
16	54	2 P	32	6 A	43.0	59	4.2	S	66	10					
17	56	4 P	40	0 A	48.0	81	0	sw	139	17	6:00 A		.02		
18	59	4 P	35	6 A	47.0	35	4.2	W	76	15	6:00 A	8:00 A	T		
19	52	2 P	27	7 A	39.5	85	2.1	N	32	4		12:00M	.98		
20	58	1 P	46	0 A	52.0	98	0.5	N	70	12	0 A	8:15 A	.83		
21	52	1 P	35	6 A	43.5	66	0.9	\mathbf{s}	92	15			T		
22	45	3 P	37	8 A	41.0	88	0	SE	25	3	4:15 P	12:00M	.19		
23	47	4 P	41	0 A	44.0	87	0	NW	22	4	0 A	3:15 A	.10		
24	47	3 P	41	8 A	44.0	96	0.4	N	56	7	6 A	12 N	.09		
25	48	3 P	28	3 A	38.0	73	0.3	NW	110	18					
26	52	3 P	26	8 A	39.0	73	4.1	SE	44	9	0.45	1.00	1.0		
27	56	1 P	37	2 A	46.5	74	0.2	NW	179	28	2:45 A	1:00P	.18		
28	46	12 N	38	7 A	42.0	57	0.2	N	185	13	0.00	10.00	T	~	
29	36	2 P	30	8 A	33.0	100	0	NE	105	9	6:30 A	12:30 P	.07	.7	
30	40	1 P	20	7 A	30.0	96	2.1	s	26	5	_				

^{*} Based on least time required to blow one mile.

LEWIS F. WELLS, Observer

	#1V11V11	
NOVEMBER, 1948	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum30.45, 2nd, 3rd	30.59	30.87, 1932
Minimum29.45, 15th		28.73, 1904
Mean semi-daily30.059		
Range1.00	1.29	1.84, 1904; 89, 1943
Air Temperature, in degrees F.	00.0	77 1004 1000
Highest		75, 1924, 1938
Lowest		4, 1938 44.1, 1931; 33.6, 1901
Range 52	53.0	44.1, 1931, 35.0, 1901
Highest mean daily64.5, 10th	35.0	
Lowest mean daily30, 30th		
Mean maximum		
Mean minimum38.1		
Greatest daily range29, 3rd		
Least daily range6, 23rd, 24th, 29th		
Degree-days of heating load for month 563		
Degree-days cumulative for season1088		
Precipitation, in inches		
Precipitation5.22	3.41	8.64, 1927;.63, 1917
Snow0.75	2.34	13.50, 1938
Maximum precip. in 24 hrs1.27, 4th		
Number of days with .01 or more13	9	22, 1921; 2, 1904
Wind in miles	4507	5,978, 1906; 2,589, 1889
Total movement	4531	3,378, 1900, 2,339, 1809
Greatest daily movement		
Least daily movement22, 23rd Mean hourly velocity4.1	6.3	
Maximum velocity48, 13th	30.3	44, 1938
	00.0	
Wind, direction	WNW	
Prevailing directionSW North, days	*******	
Northeast, days1		,
East, days		
Southeast, days7		
South, days6		
Southwest, days1		
West, days3		·
Northwest, days6	ĺ	
Weather	İ	
Mean relative humidity, percent71.6	70.6	
Mean cloudiness, percent51	55.1	78, 1947; 34, 1917
Number of clear days0	6	15, 1903; 1, 1900, 1911,1927
Number of fair days7	9	16, '12; 4, '89, '30, '32, '35
Number of cloudy days23	15	24, 1927; 9, 1905, 1917
Number of hours bright sunshine44	121	182, 1903; 66, 1927
Percent of possible hours of	41.3	
bright sunshine	41.0	
First frost	Nov. 6	Oct. 10 '25; Dec. 13, 1941
1 11 00 11 00 1	I TAGA' G	Oct. 10 20, Dec. 10, 1341

The weather for the month was both warmer and wetter than normal. Precipitation of 5.22 inches did not exceed last year's 5.63, (normal is 3.41 inches) but in both years November has ended periods of lower than normal precipitation. In both years the late summer and fall have shown very light rainfall producing fire hazards of considerable proportions in woodlands, causing dry wells and low streams, and making the growth of some fall-planted crops very slight until aided by favorable conditions in November.

The mean temperature for the month was 46.3 degrees compared to a normal mean of 38.9 degrees. This was the highest mean daily temperature this station has recorded for the month. The previous record of 44.1 degrees was recorded in 1931.

The heating load of 563 degree-days is the least for the month since records have been kept at this station. The cumulative heating load for the season now stands at 1088 degree-days, which is 301 below normal for the end of the month.

W. H. Thies, Extension Professor of Horticulture, contributes the following report: "The mildest November in recent years has influenced growing things in several ways. With little or no frost in the soil, rainfall has penetrated very largely where it fell. This will tend to benefit plants on drier soils next spring. In the case of conifers, particularly, it will tend to prevent drying from winter winds. The possibilities as regards trunk injury in fruit trees when the weather turns cold will depend, in part, on the degree of maturity. Except where trees were stimulated to make late fall growth, little damage is anticipated.

Rye cover crops have made unusually good growth thus adding more organic matter to the soil, and fall seedings are off to a good start. Strawberry plants have developed well during the fall season and there has been little need for mulching since the lowest temperature in Amherst was 20°F. on November 30. The mild fall weather, according to E. H. Wheeler, Extension Entomologist, has resulted in the laying of many aphis eggs, the adult insects having survived later than usual."

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 720

December, 1948

Meteorological Observations

FOR

DECEMBER

1948

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 252.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		ture	lity	Hours hine	,	Wind			Precipita	Precipitation			
	Maxi	mum	Min	Minimum		Humidity	of Ho unshin	20		es :			
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative	Number of Hour Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour	Began	Ended	Water Snow
1	46	2 P	26	[0 A	36	67	0	ZW	62	18		-	T
21 83	46	2 P	25	7.4	35.5	65	3.0	s	65	8			
3	44	3 P	31	7 A	37.5	77	0	NE	62	8		1	1
4	52	3 P	30	2 A 7 A	41	75	0.3	N	65	8			
5	55	3 P	28		41.5	67	4.1	7111.	62	7			
ii.	48	4 P	35	0.4	41.5	88	0	11.	70	20	8 A	5:45 P	.20
ī	55	3 P	34	± 6 A	45.5	76	2.0	8E	149	15			
`	50	1 P	31	8 A	40.5	59	0.3	ZM	60	29			
9	42	3 P	26	7 A	34	61	1.2	11.	57	7		0.00	1 0 1 1
.0	37	12 P	25	7 A	31	81	0	NW	65	22	12 N	2:00 P	.01 T
1	37	2 P	23	7.4	30	86	0	SW	70	10	1.1		1 1-
13	$\frac{1}{52}$	12 P	22	S A [32	92	0	Z.M.	$\begin{array}{c} -43 \\ 130 \end{array}$	$^{\pm}$ 10	4:45 P	7:00 P	$^{+}$.17 $^{+}$
Lii [4	52 42	4 P 2 P	36 35	4 A	$\frac{44}{38.5}$	51	3.4	11.	$\frac{150}{214}$	28			1
15	31	2 P	23	$\frac{1}{5}\frac{3}{5}$	$\begin{array}{c} 35.5 \\ 27 \end{array}$	51 74	$\frac{1}{1}$ 4.0	NE NE	92	13	7:45 P	9:45 r	$^{+}$.06 $^{+}$ 0.5
16 16	34	2 P 9 P	26	$+$ $\frac{5}{0}$ Λ^{+}	30		10	NE N	101	1.5	4:00 r		
17	$\frac{1}{39}$	3 P	30		$\frac{50}{34.5}$	89 89	0	2.ll	1.93	1 25	1 0 A		
1 8	31	1 8 A	28	1	$\frac{54.5}{29.5}$	± 69	$\begin{vmatrix} 0 \\ 1.9 \end{vmatrix}$	211	± 210	1 15	10 A	1.30 A	60.
19	$\frac{51}{24}$	- 5 A	15	$\begin{bmatrix} 5 & \Lambda \\ 6 & \Lambda \end{bmatrix}$	$\frac{29.5}{19.5}$	$\frac{1}{100}$	1.9	2.11.	74	17	1:30 1	12 1	.10
20E	32	12 P	16	1 7 A	24	1.86	1 0	S	104	4	0 3		
21	29	12 F	. 8	7 4	18.5	98	1.0	1 8	$\frac{104}{25}$	11	1 9 .		T
22	43	2 P	24	5 A	33.5	76	0	SE	164	17	i		
23	36	12 x	31	- 6 A	33.5	66	$\frac{1}{0.5}$	W	136	17			
24	24	1 P	17	1 8 A	$\frac{33.5}{20.5}$	55	1 2.8	ZW	198	22			
25	18	1 2 P	3	6 A	10.5	78	1 0	N	55	22	1		ТТ
26	20	1 P	7	8 3	13.5	1 50	3.5	N	211	22	1		
27	25	1 3 P	1 2	2 A	13.5	1 60	1 0	8	56	5			1
23	41	3 P	$\frac{24}{24}$	5 1			0		72	9			
29	51	12 P	18	1 4 5		1100	0	s	: 51	1 15	1.8	. 12 1	.61
30	58	7 P	50	11 \	54	97	1.0	1 8	221	48		12 1	
31	56	+ - 1	40	12 P	48	1 88	0	E	367	40			.52

^{*} Based on least time required to blow one mile.

LEWIS F. WELLS, Observer

Correction: An error in air temperature readings on page 3 in the November 1948 buletin gives 4° , 1938 as the extreme low. The correct reading is -4° , 1938.

WONTEL	A1411417	
December, 1948	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum30.47, 26th	30.66	30.96, 1889
Minimum28.92, 31st	29.27	28.85, 1915
Mean semi-daily29.976		
Range		1.78, 1895; 1.01, 1892
Air Temperature, in degrees F.		25.5.4000
Highest58, 30th		65.5, 1908
Lowest		-22.5, 1917
Mean32.4	27.5	36.9, 1891; 17.1, 1917
Range56	56.2	
Highest mean daily54, 30th		
Lowest mean daily10.5, 25th		
Mean maximum40.0		
Mean minimum24.6		
Greatest daily range33, 29th		
Least daily range		
Degree-days of heating load for month 1011		
Degree-days cumulative for season2099		1
Precipitation, in inches		
Precipitation2.87	3,39	7.77, 1901; .58, 1943
Snow	8.50	26.50, '02, T. 1891, 1943
Maximum precip. in 24 hrs1.03. 30th		
Number of days with .01 or more11	10	17, 1902; 4, 1892, 1943
Wind in miles		0.004 4027 0.000 1010
Total movement3404	4710	6,694, 1925; 3,239, 1918
Greatest daily movement367. 31st		
Least daily movement25, 21st		
Mean hourly velocity4.6	6.3	10 4000 4040 4044
Maximum velocity48, 30th	31.4	48, 1938, 1943, 1944
Wind, direction		
Prevailing directionNW	RXR	
North, days5		
Northeast, days2		
East, days1		
Southeast, days2		
South, days7		
Southwest, days1		
West, days4		
Northwest, days91		
Weather		
Mean relative humidity, percent76	69.4	
Mean cloudiness, percent50	54.9	71, 1929; 39, 1919
Number of clear days0	\$	15, 1890: 2, 1922, 1933
Number of fair days6	9	16, '09; 4, '89,'30,'31,'36,'38
Number of cloudy days25	14	23, '33; 7, '09, '23
Number of hours bright sunshine27	128	172, 1896; 63, 1933
Percent of possible hours of	1 = 13	<u> </u>
bright sunshine9.5	45.2	
Thunder and lightning	TO.=	
13.	Nov. 6	
	., 101. 0	

ANNUAL SUMMARY

Annual, 1948	Normal	Extremes
Barometer, in inches (Readings reduced to sea level) Maximum	30.81 28.95 30.01	31.05, 1920, 194 28.41, 193
Range2.06	1.85	2.47, 1938; 1.38, 193
Air Temperature, in degrees F.		
Highest100, Aug. 26	95.7	104, 1911
Lowest16, Jan. 24		-26. 1904
Mean48.4	47.4	49.8, 1921; 44, 1904
Range116.0	107.8	
Highest mean daily84.5, Aug. 26		
Lowest mean daily4, Jan. 24 Mean maximum59.0	57.8	
Mean minimum37.7	36.5	
Greatest daily range54, Mar. 6	00.0	
Least daily range3, Jan. 13 & Dec. 18		
Precipitation, in inches		
Precipitation40.01	43.70	59.00, 1938; 30.68, 1908
Snow52.25	47.78	\$9.00, 1893; 24.50, 1919
Maximum precip. in 24 hrs1.84, Aug. 12	11,10	1 111, 1110, 1110, 1110
Number of days with .01 or more119	124	146, 1945; 96, 1924
Wind, in miles		
Total movement	52,223	63,571, 1908: 36,257. 1894
Mean hourly velocity4.8	5.8	
48, Feb. 14, Nov. 13, Dec. 30	39.5	80, 1938
Wind, direction		
Prevailing directionW	11.	
North, days58		
Northeast, days26		
East, days		
South, days63		
Southwest, days20		
Vest, days42		
Vorthwest, days79		
Weather		
dean relative humidity, percent67.8	67.6	
Jean cloudiness, percent58.8	51.7	60, '98, '01, '02; 41, '08, '24
Sumber of clear days85	116	217, 1941; 59, 1927
Sumber of fair days127	123	182, 1912; 64, 1936
Number of cloudy days154	126	179, '01, '02; 71, 1910, 1941 3038, 1941; 1864, 1902
	0 0 = 0	SD:X (Mai' 1861 1989
Number of hours bright sunshine1886	2,353	5055, 1541, 1504, 1502
Number of hours bright sunshine1886 Percent of possible hours of bright sunshine	52.8	5050. 1541, 1604. 1502
Number of hours bright sunshine1886 Percent of possible hours of bright sunshine	52.8 pril 15	
Number of hours bright sunshine1886 Percent of possible hours of bright sunshine	52.8 pril 15 Nov. 6	Mar.14,'10; May 11,'07,'45 Oct. 10, '25; Dec. 13, '41
Number of hours bright sunshine1886 Percent of possible hours of bright sunshine	52.8 pril 15 Nov. 6 May 14	Mar.14,'10; May 11,'07,'45 Oct. 10, '25; Dec. 13, '41 Apr. 14, '43: June 8, '32

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 721

January, 1949

Meteorological Observations

FOR

JANUARY

1 9 4 9

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 252.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Te	mper	ature		dity	Hours	į,	Wind			Precipita	tion	
	Max	imum	Min	imum		Humidity	of He	5.0	د	es *				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative	Number of Hou Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1	40	0 A	28	8 A	34	85	0	SE	138	20	0.1	12:00 P	.04	1.5
2 3	33	1 P	27	0 A	30	63	0	W	167	22				
3	37	2 P	20	7.4	22.5	69	4.4	NW	110	17				
4 5	41 37	2 P	26	7 A	27	80	4.0	7//	69	4		10.00-	1 00	
9 6	55	9 P 2 P	$\frac{27}{34}$	0 A	$\frac{30}{44.5}$	$\frac{100}{71}$	0 3.3	11.	$\frac{10}{191}$	22 22	-	12:00 P 8:45 A	0.03	
7	46	12 N	33	0 A 6 A	39.5	69	0.3	811	167	17	OA	8,49A	.91	
Š	53	2 P	33	6 A	43	78	1.7	N N	198	14				
9	43	2 P	29	6 A	36	92	0	N.	33	4				
10	45	1 P	38	8.4	41.5	88	0.3	ZM	32	10	1			İ
11	33	2 P	30	8.1	31.5	50	0.3	NE	162	13			İ	!
12	29	3 P	16	3 1	22.5	67	0	2.11.	85	25				
13	38	1 P	26	7 A	32	57	2.4	NW	147	20				İ
14	41	12 N	19	12 P	30	59	0.8	N	142	22				
15	31	3 P	13	7 A	22	68	4.0	SE	109	10				
16	51	3 P	28	= 0 A	39.5	69	-2.0	SE	174	17	7:30A	8:30A	T	T
17	51	2 P	37	3 A	44	88	0.4	Z/M_s	90	20				
18	3.9	3 P	29	7 A	34	86	0	E	82	12				
19	48	5 P	35	0 A	41.5	80	0	NW	119	28	3:30A	7:00A	.10	
20	38	3 P	26	12 P	32	56	5.4	NW	291	20			0.0	
21	30	3 P	17	7 A	43.5	82	0	NE	122	15		12:00 P		1.75
22	41	1 P	25	0.4	33	86	5.5	N	100	22	0.4	7:00A	.36	1.25
23 24	35 39	2 P 12 P	$\frac{19}{27}$	7 A	27 33	65 91	$\frac{2.9}{0}$	N	62	13		9:30A	1.0	0.75
25		3 A	23	2 A 12 P	32	63	$\frac{0}{1.7}$	8	34	14	0.4	9:50A	101.	0.75
26	41 23	0 7	17	12 P	20	77	0 1	N	143	$\frac{14}{12}$	10:00A	12:00 P	2.1	2.00
27	31	2 P	18	0 A	$\frac{20}{24.5}$	88	0	XW	$\begin{bmatrix} 115 \\ 59 \end{bmatrix}$	$\frac{12}{10}$	0A	8:15	.07	.25
28	37	12 P	$\frac{15}{25}$	0 A	31	90	0	S	65	28		12:00 N		2.50
29	35	2 P	16	12 P	25.5	46	3.4	11.	156	28	2.00A	12.001		
30	30	$\begin{vmatrix} 2 & 1 \\ 2 & 1 \end{vmatrix}$	9	7 A	19.5	42	7.6	s	114	13			i	
31	30	12 P	10	4 A	20	87	0	ZW	41	22	10:00A	9:00P	.74	4.50

^{*} Based on least time required to blow one mile.

LEWIS F. WELLS, Observer

January, 1949	Normal	Extremes
	2.0711101	
Barometer, in inches (Readings reduced to sea level)		
	90.50	04.00 400
Maximum	$30.70 \\ 29.20$	31.00, 1927
Mean semi-daily30.165	$\frac{29.20}{30.07}$	28.55, 1913
Range		2.18, 1913; 0.97, 1896
		1120, 2010, 0.01, 1000
Air Temperature, in degrees F.	51.1	66.4 1000
Highest	51.1 -7.1	66.0, 1932
Mean	24.2	-26.0, 1904 34.2, 1913; 13.9, 1918
Range		01.5, 1010, 10.0, 1010
Highest mean daily44.5, 6th	1	
Lowest mean daily19.5, 30th		
Mean maximum38.7		
Mean minimum24.5		1
Greatest daily range23, 16th		
Least daily range3. 11th		
Degree Days of Heating Load		
for month		
Degree Days cumulative for Season3817		
Precipitation, in inches		
Precipitation4.52	3.61	7.15, 1898; 1.07, 1896
Snow14.5	13.32	33, 1898, 1923; 1.50, '08
Maximum precip. in 24 hrs1.03, 5th	4.4	47 100 100 4 4004
Number of days with .01 or more11	11	15, '93, '20; 4, 1901
Wind in miles		
Total movement3525	5055	7770, 1908; 2896, 1895
Greatest daily movement291, 20th		
Least daily movement10, 5th	0.0	Ť
Mean hourly velocity4.7	$\frac{6.8}{32.7}$	47 1000
Maximum velocity28, 18th, 28th, 29th	54.1	47, 1938
Wind, direction		
Prevailing direction ·NW	WNW	
North, days7		
Northeast, days2		
East, days1		
Southeast, days		
South, days		
West, days5		
Northwest, days9		
Weather	5 0 4	
Mean relative humidity, percent74 Mean cloudiness, percent64	$\begin{array}{c} 70.1 \\ 55 \end{array}$	70 1022 27 1000
Number of clear days1	99 9	78. 1932; 37, 1908 20, 1944; 2, 1914
Number of fair days	9	18, 1890, 1926; 3, 1916, '44
Number of cloudy days22	13	22, 1923, '31; 4, 1920, '39
Number of hours bright sunshine50.4	137	214, 1920; 74, 1932
Percent of possible hours of]
bright sunshine17.1	46.7	
Thunder and lightning		
?		

The weather for the month of January was characterized by its mildness in comparison with the same month of last year. With a mean daily temperature of 31.8 degrees for the month this was the warmest January since 1933, but the maximum of 55 degrees was reached in 1941. The householder noticed the effect on his purchased fuel with a heating load of 1029 degree-days for the month and a cumulative load of 3128 for the season. Normal for the month and season are 1265 and 3817 degree-days respectively, while a year ago these values stood at 1459 degree-days for the month and 3932 for the season.

The fall of snow was slightly above normal, with total precipitation nearly an inch above normal for the month. Following the fall on the 31st, the month ended with 10.5 inches of snow on the ground. While the rainfall at the end of the previous month had relieved anxiety over short water supply for wells, the more than average precipitation and the snow blanket assured good ground water supply for the remainder of the winter. These combined with less than normal wind movement relieved vegetation of danger of dessication which could have taken place had the relatively mild temperatures been accompanied by drought conditions.

W. H. Thies, Extension Professor of Horticulture, contributes the following report: "The effect on fruit buds of an unseasonably mild winter is of interest to fruit growers and consumers alike. Recent warm weather has caused some swelling of fruit buds although apples and pears have apparently developed less than reaches."

According to J. S. Bailey, "Peach fruit buds appear large for this time of year. This is not surprising since the winter rest perriod ended several weeks ago. The warm weather has also caused internal changes which make the buds more susceptible to winter injury. Should really cold weather come, however, there may still be little injury if the temperature drops slowly; if it drops rapidly to several degrees below zero, severe injury may be expected.

"On February 4 there were practically no dead buds in the University peach orchard. If 10 to 15% of the buds are alive on Washington's Birthday, the chances for a crop are excellent."

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 722

February, 1949

Meteorological Observations

FOR

FEBRUARY

1 9 4 9

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 252.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Te	mpera	ture		lity	Hours hine	7	Vind		Precipitation				
	Maximum		Minimum			Humidity	r of Hour	80		5 *					
Day	Degrees	Time	Degrees	Time	Мевп	Mean Relative	Number Bright St	Prevaili ng Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow	
1	34	2 P	23	6 A	28.5	54	6.9	W	158	22	i i		T	T	
2 3	28	1 P	15	12 P	21.5	60	4.6	W	254	28	İ		ĺ	İ	
	31	1 P	5	7 A	18.0	54	4.3	SE	77	7				j	
4	29	2 P	21	6 A	25.0	86	0	NW	71	6	7:30A	8P		2.00	
5	30	12 N	11	[12 P]	20.5	58	8.9	N	121	22	-	1	T	T	
6	30	3 P	-1	7 A	14.5	54	9.8	NW	-						
7	41	4 P	11	12 P	26.0	77	2.7	N	-	28		_] .15	T	
8	42	3 P	16	7 A	29.0	68	10.3	s	118	15		1			
9	48	1 P	28	12 P	38.0	79	8.4	NW	148	28					
10	38	2 P	16	7 A	27.0	64	2.9	N W	66	7					
11	29	11 A	10	12 P	19.5	60	10.4	N	211	28					
12	32	4 P	0	7 A	16.0	72	8.8	SE	103	15					
13	51	4 P	35	0 A	43.0	68	3.4	SE	156	11	1			ļ	
14	51	2 P	35	9 A	43.0	76	3.0	N	42	7					
15	47	11 P	37	7 A	42.0	76	0.8	NE	41	28		10:45P	.52		
16	42	1 P	35	12 P	38.5	58	0	W	150	15	[10:15P	12:00P	.12		
17	35] 0 A	24	12 P	29.5	65	9.7	W	272	48	0.4	2:30A	.19	1.25	
18	42	5 P	12	6 A	27.0	64	10.6	SE	142	20					
19	54	2 P	33	7 A	43.5	62	6.9	s	150	8	1				
20	46	3 P	30	12 P	38.0	84	3.9	N	76	20	0A	7:30A	.48		
21	39	4 P	22	7.4	30.5	67	10.9	N	142	13					
22	38	10 P	28	2 1	33.0	84	0	s	111	10					
23	45	2 P	31	12 P	38.0	77	10.1	N	162	25					
24	48	3 P	26	7 A	37.0	65	10.0	N	53	9	0 15		0.0	0=	
25	37	0 A	34	8 A	35.5	94	0	N	90	22	6:45A	81	.30	.25	
26	37	11 A	26	12 P	31.5	63	10.1	NW	280	22				-	
27	41	3 P	18	6 A	29.5	66	11.1	NW	122	14	1.	4.0	10	0.00	
28	34	12 N	24	3 A	29.0	90	0	NW	86	32	3Р	12P	.48	9.00	

^{*} Based on least time required to blow one mile.

LEWIS F. WELLS, Observer

February, 1949	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum30.59, 12th	30.63	31.05, 1926
Minimum29.66, 15th	29.24	28.56, 1895
Mean semi-daily30.116	30.04	
Range0.93	1.38	1.89, '00, '08; .88, '13, '31
Air Temperature, in degrees F.		27.0 4000
Highest54, 19th	50.5	65.0, 1930
Lowest1, 6th	$-7.4 \\ 23.7$	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$
Mean30.4	58	32.0, 1330, 11.0, 1334
Range55	90	
Highest mean daily43.5, 19th		
Lowest mean daily14.5, 6th	32.8	
Mean maximum39.3	14.5	
Mean minimum	14.0	
Greatest daily range		
Degree-days of heating load for month 958		
Degree-days of heating load for month 558 Degree-days cumulative for season 4086		
begree-days cumulative for season 4000		
Precipitation, in inches		
Precipitation2.47	3.19	8.12, 1900; 0.62, 1901
Snow13.25	13.99	48.75, 1893; 0.50, 1937
Maximum precip. in 24 hrs52, 15th		
Number of days with .01 or more8	10	15, '93, '20; 4, 1901
Wind in miles		
Total movement3402	4776	6445, 1896; 3438, 1892
Greatest daily movement280, 26th		
Least daily movement41, 15th		
Mean hourly velocity	7.1	50, 1946
Maximum velocity48, 17th	31.7	50, 1340
Wind, direction		
Prevailing directionNW	WNW	
North, days9		
Northeast, days1		
East, days		
Southeast, days		
Southwest, days0		
West, days4		
Northwest, days		
•		
Weather	000	
Mean relative humidity, percent69	66.6	ee 100 107 - 21 1005
Mean cloudiness, percent41.6	$\frac{50.9}{10}$	66, '90, '27; 31, 1905 19, 1941; 2, 1927
Number of clear days	8	16, 1920; 2, 1936
Number of fair days	10	16, 1920, 2, 1930
Number of cloudy days	158	221, 1924; 110, '27, '38
Percent of possible hours of	190	221, 1021, 110, 21, 00
bright sunshine57.16	53.4	
Thunder and lightning	00.1	

The weather for the month continued the mildness recorded for the first month of the new year. With a mean daily temperature of 30.4 degrees, it was the warmest February since 1890 which recorded 32.6 degrees, and is approached only by February 1937 which showed 30.0 degrees. Against a normal mean daily temperature of 23.7 and last year's 21.4 degrees the month showed a mildness that was appreciated. The house-holder had to supply fuel for only 958 degree-days of heating load against last year's 1264 and the normal 1165 degree-days. At the end of the month the cumulative load for the season stood at 4086 against last year's 5196 degree-days and the normal 4982, or only 82 per cent of normal.

The snow fall was nearly normal but the precipitation of 2.47 inches of water was 0.72 inches less than the mean for the month. Some amount of snow covered the ground until the 19th, and from that date the ground was bare until the storm of the 28th. Following a trace of snow on the 5th, the morning of the 6th recorded the lowest temperature of the winter, -1 degree. Excepting the zero recorded the morning of the 12th, the other cold mornings of the month were merely "frosty" and coupled with warm afternoons started the maple sugar season.

W. H. Thies, Extension Professor of Horticulture, contributes the following report: "One might think that an unseasonably mild winter would cause very early development in all kinds of vegetation. But, fortunately, plant life is not greatly influenced by above normal winter temperatures. The mean temperature during February was more than 6° above normal and yet apple buds, although somewhat swollen, remain in a more or less dormant condition. If cool weather prevails during the next few weeks, Mc-Intosh apple trees may still come into bloom around May 10 to 15, as usual.

Concerning the effect of recent weather on forest trees, A. D. Rhodes of the Forestry Department says, "The mild winter has to date been generally favorable for forest trees. Plentiful rainfall, which fell on soil relatively free of frost and therefore well able to absorb it, has restored any deficit in soil moisture that may have existed as an aftermath of last summer's drought. Winter killing of conifers resulting from excessive transpiration, which might have been anticipated, is not likely to occur. Growth this spring should begin fairly early and proceed rapidly unless the weather at that time is unseasonably cold. Cold spring weather may damage buds of some trees, especially those not well protected, which already have shown some response to the spring-like conditions during February."

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 723

March, 1949

Meteorological Observations

FOR

MARCH

1 9 4 9

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 feet. Time used, E. S. T.

		Ter	npers	ture		lity	Hours	•	Wind			Precipitation				
	Maximum		Minimum			Humidity	of Ho unshin	60	ا ،	**						
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative	Number of Hou Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow		
1	36	4 P	23	12 P	29.5	92	5.3	NW.	408	32	(0.A	12A	0.29	7.0		
2	35	1 P	16	6 A	25.5	79	10.9	NW	218	15						
3	. 41	12 N	18	6 A	29.5	7.0	11.3	NW	118	20						
4	37	4 P	9	7 A	23.0	85	11.2	SE	78	13						
5	54	2 P	29	0.4	41.5	7.9	8.1	S	139	12			1			
6	51	2 P	38	8.4	44.5	81	4.4	N	86	12						
7	35	2 P	21	12 M	28.0	59	9.1	YE.	277	22						
8	47	5 P	15	6.4	31.0	35	3.1	ZE	212	15						
9	57	1 4 P	25	+ 6 A	41.0	45	8.8	N	126	14						
10	46	2 P	28	6 A	37.0	60	0	N	119	10	0.		0.09	1 4 0		
11	38	2 P	32	7 A	35.0	100	0 .	NΕ	102	15	6A	21	0.03	1.0		
12	34	1 P	30	7.4	32.0	73	1.1	W	$\frac{301}{320}$	$\frac{40}{28}$						
13	36	6 P	22	7.1	29.0	53	-	W	206	28 22						
14	47	5 P	26	7.1	36.5	55	6.4	W	183	18			1	1		
15	39	2 P	29	4 A	34.0	48	5.2	ZW	$\frac{185}{298}$	22 22						
16	36	3 P	20	7 A 5 A	$\frac{28.0}{30.5}$	42 45	-8.2 + 9.9	2 W	$\frac{236}{236}$	28	3:30A	4:30a		T		
17	39	5 P	22 22		$\frac{30.5}{25.5}$	94	0.0	ZM.	100	18	10:304	6:00P		2.5		
18	29	12 X	9		$\frac{26.5}{26.6}$	1 46	8.3		260	17	10.50.1	0.001		2.0		
19	31	2 P 5 P	2		$\frac{20.0}{19.5}$	53	9.3	2 II.	50	8						
$\frac{20}{21}$	51 57	1 5 P	22	7 A	39.5	- 53 - 82	6.2	- 11	209	25						
	64	3 P	13	L. = .\	53.5	S3	6.0		258	25						
22 23	68	5 P	41	12 P	54.5	1 94	0.8	- 4	217	17	3:00A	9:30A	0.71	1		
24	58	6 P	32	7.4	45.0	66	8.7	N	143	13	1	0.000	''. 1 1			
25	58 58	1 3 P	30	1. 3.	$\frac{43.0}{44.0}$	71	9.9	~ II.	53	13		ì		1		
26 26	- 5 - 75	4 P	42	+ A	58.5	76	$\frac{2.2}{7.3}$	Z.	136	10	i	1				
27	72	5 P	56	6.3	64.0	90	0.2	777	94	11	1			İ		
28	113	3 A	56	8.4	59.5	52	8.8 :	5	371	32	İ					
29	66	4 P	38	7.3	52.0	46	7.9	NE	209	22				İ		
30	60	4 P	37	6.3	48.5	73	4.5	NE	109	13	ĺ					
31	51	4 P	37	7 A	44.0	58	0.8	NE	122	13				ĺ		

^{*} Based on least time required to blow one mile.

LEWIS F. WELLS, Observer

March, 1949	Normal	Extremes
March, 1949	30.57 29.24 30.04 1.34 63.8 6.1 34.4 57.6 43.2 25.4	Extremes 31.05, 1943 28.47, 1914 2.10, 1914; 0.85, 1915 85, 1929 -7.5, 1906
Precipitation, in inches Precipitation	2.70 7.47 11	7.89, 1942; 0.12, 1915 27, 1899; 0, 1921 17,1890,1913,1936; 3,1915
Wind in miles Total movement	5944 7.7 31.2	\$182, 1896; 3006, 1905 48, 1932, 1909, 1941, 1942
Wind, direction Prevailing direction NW North, days 5 Northeast, days 6 East, days 6 Southeast, days 1 South, days 5 Southwest, days 3 West, days 4 Northwest, days 7	WNW	
Weather Mean relative humidity, percent 67 Mean cloudiness, percent 57 Number of clear days 11 Number of fair days 9 Number of cloudy days 9 Number of hours bright sunshine 174.1 Percent of possible hours of bright sunshine 47 Thunder and lightning -	64.2 51.8 11 10 10 199 53.6	68,1901; 27, 1915 22, 1924; 3, 1901 17, 1900; 1, 1943 21, 1901; 1, 1915 292, 1924; 93, 1901

While the weather for the month continued warmer than normal, the 75 degree maximum recorded on the 26th was 10 degrees less than the record March temperature of 1929. In nine other years of record, the mean daily temperature has exceeded the 38.3 degrees of the past month, but never before have the first three months of the year combined greater-than-average mean daily temperatures to produce the mildness of the past season. The heating load for the month was 831 degree-days against a normal of 950 and last year's 932, while the season's cumulative load at the end of the month totaled 4917 against a normal of 5932 and last year's 6128 degree-days.

Except for light snow on the 18th, snow cover disappeared on the 7th. Precipitation was very light at 1.67 inches total water in the more than normal 10.5 inches of snow fall. Measurable precipitation occurred on only five days of the month even though cloudiness exceeded normal and hours of bright sunshine were less than normal.

W. H. Thies, Extension Professor of Horticulture, contributes the following report: 'In spite of the many balmy days during the winter, vegetation is still no more than two weeks ahead of normal. A cold spell now and then, as for example, a low temperature of 2° on March 20 have provided a necessary check on swelling buds and a reminder that spring is not "just around the corner." How mild the past winter has been is well illustrated by these comparisons: January was the warmest since 1933, February the warmest since 1890, and March among the warmest since 1929.

March was a relatively dry month with slightly more than an inch of rainfall in Amherst. Soaking rains and melting snowfall on unfrozen ground in previous months, however, have left the soil well supplied with moisture. Cover crops have wintered well. Fruit buds have thus far suffered no appreciable damage, although growers are already on the alert in the knowledge that apple scab spores are maturing early. Numerous pear psylla adults have been observed in some orchards and red mite of one or more species promises to be a problem.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 724

April 1949

Meteorological Observations

FOR

APRIL 1949

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

	Temperature						s		Wind		1	Precipitation		
	Max	Maximum		imum		Iumidi	f Hour nshine			g				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max Miles per Hour*	Began	Ended	Water	Snow
1 1 2 3 4 4 5 6 7 8 8 9 100 111 122 133 144 155 166 177 188 199 221 222 223 24 225 226 227 288 29 30 31	55 63 60 48 62 49 60 57 49 58 61 72 73 77 64 55 55 50 51 76 62 62 70 74 63 60 79	5 P 2 P 3 P 4 P 12 N 4 P 1 P 2 P 2 P 3 P 1 1 A 3 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P	35 33 40 40 40 27 40 42 36 37 31 32 31 41 49 46 44 33 30 34 42 49 45 30 34 48 44 44	7 A 7 A 6 A 6 A 3 A 0 A 2 A 3 A 5 A 6 A 6 A 4 A 6 A 4 A 6 A 4 A 6 A 6 A 6 A 7 A 8 A 8 A 8 A 8 A 8 A 8 A 8 A 8 A 8 A 8	45. 48. 50. 44.5 44.5 51. 46.5 51.5 57.0 63.0 55.0 49.5 41.5 40.5 44.5 57.5 60.0 60.5 53.0 57.5 60.0 60.5 53.0 63.0	60 62 56 93 52 90 65 74 59 57 52 65 68 83 91 65 69 85 77 66 67 41 71 62 55 53 58	3.6 5.7 5.8 0 8.2 0.3 4.0 4.7 3.5 10.3 8.0 8.7 7.9 6.7 1.1 2.6 3.3 0.6 3.1 6.3 2.2 3.6 6.8 5.1 9.1	ne sw ne ne ne ne ne ne ne ne ne ne ne ne ne	157 77 151 148 83 213 308 121 265 104 89 132 157 163 213 127 214 127 214 132 201 107 264 78	14 8 18 18 18 15 32 28 20 20 17 12 14 22 25 17 15 20 10 22 12 22 11 15	7.30A 2:00 A 0 A 9:30P 0 A 5:00P 2:45P 0 A	11:00A 5:30 P 7:00 A 12 P 9:30A 6:30P 12 P 6:00A	0.73	,

^{*} Based on least time required to blow one mile.

MONTHE	Sommi	111(1
APRIL 1949	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.38, 29th Minimum 29.16, 6th Mean semi-daily 29.920	30.48 29.38 29.99	30.71, 1911 28.98, 1943
Range1.22	1.11	1.52, 1930; .72, 1919
Air Temperature, in Degrees F.		
Highest .79, 30th Lowest .27, 5th, 29th Mean .50.2 Range .52 Highest mean daily .61, 14th	79.4 22.0 45.7 57.4	90, 1941 8.5, 1923 52, 1921; 41, 1943
I.owest mean daily40, 18thMean maximum62.5Mean minimum37.9Greatest daily range.63, 14thLeast daily range40.5, 18thDegree-days of heating load for month.444Degree-days cumulative for season5361	56.8 34.8	
Precipitation, in inches		
Precipitation	3.35 2.17	6,89, 1929; .55, 1941 11, 1891; 0, 1910, 1934
Number of days with .01 or more11	11	18, 1909; 3, 1892
Wind, in miles		
Total movement	5404	8208, 1908; 3853, 1917
Mean hourly velocity 6.7 Maximum velocity 32 6th, 9th	7.5 31.4	40, 1935, 1938, 1940, 1945
Wind, direction		
Prevailing direction N North, days N 5 Northeast, days 10 East, days 2 Southeast, days 0 South, days 2 Southwest, days 0 West, days 2 Northwest, days 3	WNW	
Weather		
Mean relative humidity, percent	61.6 51.8 11 9 10 220	75, 1901; 34, 1925, 1927 23, 1941; 3, 1898, 1902 18, 1915; 2, 1894, 1901 22, 1901; 1, 1944 329, 1941; 103, 1901
bright sunshine	54.7 Apr. 15	Mar. 14, 1910; May 11, 1907



MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 725

May 1949

Meteorological Observations

FOR

MAY 1949

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

		Te	трега	ture		ı,	s	Wind Precipitati						tion	
	Ma	ximum	Min	imum		Humidí	of Hour nshine		į.	8.					
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max Miles per Hour*	Began	Ended	Water	Snow	
1 2 3	80 72 75 84	1 P 1 P 5 P 3 P	48 62 59 44	6 A 5 A 8 A 6 A	64 67 67 64	73 89 90 60	7.3 0.4 3.2 6.3	sw sw sw	133 214 141 63	15 17 14 8	3 P 6 P 0 A	7 P 12 P 6 P	.01 .13 1.11		
4 5 6 7 8	92 72 78 72	4 P 5 P 9 A 4 P	51 55 60 39	5 A 3 A 5 A 6 A	71 63 69 55	69 92 77 62	6.6 5.2 8.6 11.1	sw sw e	97 63 196 91	40 12 20 11	2 A	3 A	.01		
9 10 11	82 70 65	3 P 3 P 3 P	47 52 41	5 A 8 A 5 A	65 61 53	61 65 50	5.5 2.8 3.4	w e e	113 105 187	8 12 12 10	1 A	3 A	.06		
12 13 14 15	58 69 76 72	4 P 4 P 11 A 2 P	37 34 46 46	5 A 3 A 4 A 5 A	47 52 61 59	82 67 74 64	3.1 3.0 1.1 8.4	se w sw s	133 78 126 105	9 8 12	6 P	7 Р	Т		
16 17 18 19	72 85 89 91	4 P 4 P 5 P 3 P	39 46 53 62	4 A 5 A 5 A 5 A	55 66 71 76	71 57 68 75	0.8 4.8 4.8 4.3	sw s w sw	71 58 122 175	7 4 17 28	0 a	12 р	.12		
20 21 22 23	64 68 60 76	1 A 5 P 11 A 12 N	46 38 46 53	7 A 5 A 2 A 1 A	55 53 53 65	93 56 75 75	0.0 5.0 0.0 6.8	ne w sw ne	135 63 131 120	10 8 12 18	2 A 12 N 0 A	12m 1 A	.95 .39 .05		
24 25 26 27	80 64 65 65	1 P 5 P 1 P 1 P	49 53 39 45	6 A 6 A 4 A 6 A	65 59 52 55	74 49 70 68	5.0 10.1 2.6 5.5	w ne w	76 206 107 150	20 32 18 20	6 P 0 A 0 A 7 P	12 P 12 P 3 A 8 P	1.56 .12 .16 .03		
28 29 30 31	62 61 65 73	3 P 1 P 1 P	46 43 36 44	5 A 3 A 3 A 4 A	54 52 50 58	65 64 59 60	4.9 5.3 5.0 10.0	n n n	109 140 120 35	40 25 32 7	, -				

^{*} Based on least time required to blow one mile.

MAY 1949	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.32, 4th Minimum 29.64, 23rd Mean semi-daily 29.98 Range 0.68	30.39 29.51 29.96 0.88	30.62, 1936 29.10, 1938
Air Temperature, in Degrees F.	:	0.5 1007 1011
Highest 92, 5th Lowest 34, 13th Mean 60.0 Range .58 Highest mean daily 76.0, 19th Lowest mean daily 50.0, 30th Mean maximum 72.7 Mean minimum 47.1 Greatest daily range 41, 5th Least daily range 10, 2nd Degree-days of heating load for month 190 Degree-days cumulative for season 5551	86.1 31.2 57.1 54.9	94.5, 1896, 1911 24.0, 1900
Precipitation, in inches		
Precipitation	3.60	7.44, 1931; .48, 1903
Snow	12	20. 1901, 1945; 5, 1903
Wind, in miles		
Total movement 3663 Greatest daily movement 214, 2nd Least daily movement 35, 31st Mean hourly velocity 4.9 Maximum velocity 40, 5th, 28th	6.1 28.8	5946, 1907; 2180, 1894 45, 1935
Wind, direction		
Prevailing direction W North, days 3 Northeast, days 4 East, days 3 Southeast days 1 South, days 2 Southwest, days 9 West, days 9 Northwest, days 0	W	
Weather		
Mean relative humidity, percent. 69.5 Mean cloudiness, percent. 54.7 Number of clear days 18 Number of fair days 18 Number of cloudy days 12 Number of hours bright sunshine. 150.9 Percent of possible hours of bright sunshine. 33.4 Thunder and lightning. 1st, 5th, 18th, 19th, 24th 25th	60.7 52.1 11 11 252 55.6	72, 1948; 30, 1923 21, 1944; 0, 1927 17, 1907, 1926; 5, 1923 20, 1927; 2, 1941 350, 1944; 137, 1927

•

Meteorological Series

Bulletin No. 726

June 1949

Meteorological Observations

FOR

JUNE

1949

II. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

	Temperature								Wind			Precipita	ation	
	Max	imum	Mir	imum		Humidit	of Hours unsbine	1g	nt	les *				
Day	Degrees	Time	Degrees	Тіте	Mean	Mean Relative Humiditv	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max Miles per Hour*	Began	Ended	Water	Snow
1 2 3 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 30 30 30 30 30 30 30 30 30 30 30 30	77 81 82 85 83 74 68 74 76 89 87 87 87 87 87 87 87 87	4 P 3 P 1 P 2 P 4 P 2 P 1 P 5 P 1 P 5 P 1 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 2	44 49 48 50 60 48 57 42 39 51 52 64 68 67 69 70 66 68 70 66 67 58 66 72 68 61 59 59 59 59 59 59 59 59 59 59 59 59 59	5 A A A A A A A A A A A A A A A A A A A	60.5 65.0 65.0 67.5 71.5 65.5 65.5 56.5 70.0 78.5 78.5 74.0 76.5 78.5 79.0 82.0 75.0 76.0 76.0 76.0 76.0 76.0 76.0 76.0 76	53 59 54 62 71 62 71 62 74 65 67 66 67 68 70 62 74 45 54 75 61 66 68	10.1 7.9 7.1 7.4 8.7 7.0 4.8 6.1 7.3 3.1 4.0 7.5 5.3 5.1 5.2 2.1 5.2 4.3 3.3 3.6 6.5 4.0 9.6 8.5 2.7	S SW W N N N N N N N N N N N N N N N N N	50 86 152 145 121 104 112 110 111 176 127 91 178 185 168 122 117 96 85 120 120 104 143 102 136 235 113 76 73 74	7 8 12 11 13 12 28 17 8 13 12 8 14 12 12 9 10 10 10 10 8 11 12 25 25 20 20 20 20 6 6 6 6 6 6 6 6 6 6 6 6 6 6	8 A 4 A 5 P	12 N 9 A 7 P	.07 .39	

^{*} Based on least time required to blow one mile.

	501.11.1	711(1
JUNE 1949	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.40, 10th Minimum .29.58, 22nd Mean semi-daily .29.77 Range 0.82	30.30 29.55 29.95 0.75	30.70, 1947 29.24, 1902
Air Temperature, in Degrees F.		101,1919
Highest 94, 21st Lowest 39, 9th Mean 71.9 Range .55 Highest mean daily 82, 21st Lowest mean daily 55, 8th Mean maximum 84.8 Mean minimum 59,2 Greatest daily range 36, 11th Least daily range 13,28th Degree-days of heating load for month 24.5 Degree-days cumulative for season 5575.5	91.2 40.0 65.7 51.3	34, 1891
Precipitation, in inches		
Precipitation	3.75	9.68, 1922;0.76, 1908
Snow	11	17, 1922, 1945; 4, 1908
Wind, in miles		
Total movement 3632 Greatest daily movement 235, 26th Least daily movement 73, 29th Mean hourly velocity 5.5 Maximum velocity 32, 26th	3585 5.0 24.6	4571, 1908; 1409, 1906 48, 1939
Wind, direction		
Prevailing direction sw North, days 1 Northeast, days 3 East, days 4 Southeast, days 0 South, days 4 Southwest, days 12 West, days 5 Northwest, days 1	WSW	
Weather		
Mean relative humidity, percent. 62.6 Mean cloudiness, percent. 39.7 Number of clear days 0 Number of fair days 21 Number of cloudy days 9 Number of hours bright sunshine 179.9 Percent of possible hours of bright sunshine 39.3 Thunder and lightning 20th	66.9 51.1 10 12 8 257 54.0	71, 1903; 28, 1908 22, 1908, 1941, 1943, 1, 1927 23, 1912;3, 1941 22, 1903; 1, 1923 362, 1908;102, 1903

		•	

Meteorological Series

Bulletin No. 727

July 1949

Meteorological Observations

FOR

JULY

1949

II. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

		Те	mpera	iture		ty			Wind]	Precipita	tion	
		imum		nimum		Mean Relative Humidity	Number of Hours Bright Sunshine	ng no	ent	liles 1r*				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relativ	Number Bright	Prevailing Direction	Total Movement	Max Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8 9 10 11 11 12 13 14 15 16 17 18 19 20 21 22 22 23 24 25 26 27 28 28 29 20 20 20 20 20 20 20 20 20 20 20 20 20	89 91 95 96 88 88 85 85 87 83 85 87 83 88 89 90 92 95 88 88 89 89 90 92 95 88 89 89 89 89 88 89 88 89 88 89 88 89 88 88	2 4 P P P P P P P P P P P P P P P P P P	55 55 69 66 66 73 55 55 63 65 66 67 70 72 74 69 66 65 71 71 70	5 A A A A A A A A A A A A A A A A A A A	72.0 75.0 82.0 81.0 80.5 73.5 71.0 69.0 75.0 75.5 75.5 73.5 72 73.5 80 82 84.5 71.5 74.5 69.5 71.5 74.5 69.5 71.5 74.5 79.8 83.5	58 58 60 59 59 56 48 63 75 81 81 81 82 73 65 79 70 73 68 64 66 68 68 68	4.6 6.2 5.2 9.8 11.0 8.1 10.7 4.3 7.0 11.3 11.6 6.6 12.1 9.4 12.9 9.2 10.3 6.4 12.7 10.3 11.8 11.8 12.9	sw w sw ne sw sw ne sw sw ne sw sw ne sw sw ne sw sw ne ne sw sw ne ne sw sw ne ne w ne ne sw sw ne sw sw ne sw sw ne sw sw ne ne sw sw sw ne ne sw sw sw ne sw sw ne sw sw sw ne sw sw sw ne sw sw sw ne sw sw sw ne sw sw sw sw ne sw sw sw ne sw sw sw sw ne sw sw sw sw ne sw sw sw ne sw sw sw ne sw sw sw ne sw sw sw sw ne sw sw sw sw ne sw sw sw sw ne sw sw sw ne sw sw sw sw ne sw sw sw sw sw ne sw sw sw sw sw sw sw sw sw ne sw sw sw sw sw sw sw sw sw sw sw ne sw sw sw sw sw sw sw ne sw sw sw sw sw sw sw sw sw sw sw sw sw	67 65 79 57 130 89 89 55 127 161 56 127 88 54 104 59 46 136 107 121 109 34 211 94 75 62 112 93 78	9 8 11 22 14 11 10 6 13 14 9 10 8 9 9 6 5 14 9 28 11 13 15 13 6 7 8 8 12 14 14 14 15 16 16 17 18 18 18 18 18 18 18 18 18 18 18 18 18	3 P 11 A 0 A 9 P 1 P 2 A 4 P 2 F 6 A	3 P 5 P 10 P 2 P 7 A 5 P 4 P 9 A	.04 .42 1.00 .02 .04 .43 .43 .28	

^{*} Based on least time required to blow one mile.

LLOYD S. COHAN, Observer

MONTHLY	SUMINIA	114.1
JULY 1949	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum .30.32, 1st Minimum .29.70, 31st Mean semi-daily .29.982	30.27 29.59 29.96	30.50, 1892 29.27, 1932
Range	0.68	0.97, 1892; 0.47, 1938
Air Temperature, in Degrees F.		
Highest 97, 30th Lowest 53, 8th Mean 76,00 Range 44 Highest mean daily 84,5,20th Lowest mean daily 68.0, 10th Mean maximum 87.3 Mean minimum 64.7 Greatest daily range 34, 1st Least daily range 10, 10th Degree-days of heating load for month Degree-days cumulative for season	93.9 46.4 70.8 47.7	104, 1911 40, 1890, 1898 74.7, 1921; 66.3, 1891
Precipitation, in inches		
Precipitation	4.10	14.51, 1897,;0.70, 1929
Snow Maximum precip. in 24 hrs 1.00, 13th Number of days with .01 or more 11	11	20, 1915; 4, 1924
Wind, in miles		
Total movement 2820 Greatest daily movement 211, 23rd Least daily movement 34, 22nd Mean hourly velocity 3.8 Maximum velocity 28, 20th	3422 4.6 28.7	5097, 1909; 1109, 1894 44, 1936
Wind, direction		
Prevailing direction W North, days .4 Northeast, days .7 East, days .2 Southeast, days .0 South, days .0 Southwest, days .13 West, days .3 Northwest, days .2	SW	
Weather		
Mean relative humidity, percent. 68,8 Mean cloudiness, percent. 44,2 Number of clear days	68.4 50.9 10 14 7 268	70, 1943; 31, 1924 22, 1923; 0, 1915 24, 1909, 1921; 4, 1946 18, 1889; 0, 1910 371, 1910; 180, 1931

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 728

August 1949

Meteorological Observations

FOR

AUGUST

1949

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

		Te	mpera	ture		t,y	60		Wind		I	Precipita	tion	
	Max	imum	Mir	imum		Iumidi	f Hour nshine			35				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8 9 10 11 12	84 86 78 81 86 89 91 92 98 98 93 80	5 P 3 P 4 P 4 P 3 P 4 P 2 P 3 P 4 P 1 P 3 P	59 58 69 71 65 61 61 67 65 66 71	5 A 3 A 5 A 6 A 6 A 6 A 5 A 6 A 7 A	71.5 72.0 73.5 76.0 75.5 75.0 76.0 76.5 82.5 81.5 79.5	67 68 82 87 80 79 66 62 60 56	14.1 13.3 12.5 7.0 11.9 13.5 13.6 13.5 10.4 14.2 9.0 1.0	sw sw w nw ne n nw nw sw nw	41 47 48 29 47 32 53 84 75 70 50 45	12 9 6 12 14 13 14 8 8	3 A 4 A	6 P 5 A	.16 .01	
13	80	2 P	68	7 А	74.0	87	0.5	se	58	11	7 P 3 A 6 P	10 P 7 A 7 P	.29 1.05 .03	
14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31	80 83 78 79 75 74 73 83 86 87 80 94 95 92 75 81 77	5 P 3 P 12 N 3 P 4 P 3 P 5 P 1 P 3 P 4 P 5 P 1 P 2 P 2 P 2 P 4 P 2 P 2 P 4 P 2 P 2 P 4 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2	63 61 60 58 59 58 45 47 47 55 64 52 58 65 70 64 55 59	3 A A 3 A A 6 A A 6 A A 7 A A 6 A A 7 A 7 A 7 A	71.5 72.0 69.0 68.5 67.0 66.0 59.0 66.5 75.5 76.0 80.0 81.0 69.5 68	79 69 63 68 84 62 62 62 77 70 56 65 73 75 80 77 92	2.0 12.0 13.9 7.0 6.0 10.7 12.3 11.3 11.7 10.9 12.7 12.0 11.5 8.3 2.5 12.2 1.0	se sw sw w e n sw sw w ne ne s sw w me ne sw w ne ne	58 41 82 90 62 158 149 51 55 60 98 85 126 89 140 214 73 31	12 8 9 15 17 20 7 7 7 14 8 11 9 17 22 7 12	3 A	5 P	1.14	

^{*} Based on least time required to blow one mile.

LLOYD S. COHAN, Observer

AUGUST 1949	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum. 30.27, 16th Minimum. 29.58, 18th Mean semi-daily 29.993 Range. 0.69	30.32 29.61 30.00 0.71	30.50, 1934 28.87, 1930
Air Temperature, in Degrees F.		
Highest 98; 9th, 10th Lowest 45, 20th Mean 72.7 Range 53.0 Highest mean daily 82.5, 9th Lowest mean daily 59, 20th Mean maximum 84.3 Mean minimum 60.8 Greatest daily range 39, 22nd Least daily range 9, 3rd, 12th Degree-days of heating load for month. Degree-days cumulative for season	91,6 43.4 68.6 48.1	100, 1918, 1948 34, 1940 62.4, 1903
Precipitation, in inches		
Precipitation	4.08	8.40, 1948; .31, 1894
Snow	11	16, 1892, 1933; 4, 1899
Wind, in miles		
Total movement	3127 4.2 22.7	4,271, 1910; 1,920, 1894 40, 1941
Wind, direction		
Prevailing direction SW North, days 2 Northeast, days 4 East, days 1 Southeast, days 2 South, days 4 Southwest, days 10 West, days 4 Northwest, days 4	SW	
Weather		
Mean relative humidity, percent. 72.0 Mean cloudiness, percent. 45.3 Number of clear days .21 Number of fair days .5 Number of cloudy days .5 Number of hours bright sunshine .304.5 Percent of possible hours of bright sunshine .70.7	70.2 49.7 9 13 9 237	67, 1901, 1946; 27, 1923 23, 1941; 0, 1915, 1929 25, 1912; 3, 1939 18, 1901, 1928; 2, 1910, 1923 332, 1941; 152, 1915, 1929
Thunder and lightning	 Sept 21	Aug. 22, '94, '95; Oct. 13, '09

£		
		•
		ż

Meteorological Series

Bulletin No. 729

September 1949

Meteorological Observations

FOR

SEPTEMBER

1949

H. N. STAPLETON

OBSERVATORY


Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

		Te	empera	iture		11	w.		Wind			Precipita	ation	
	Ma	simum [Min	nimum		Humidi	of Heur	u.	nt	2*				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sun gine	Preveiting Direction	Total Movement	Max Miles per Hour*	Becan	Ended	Water	Snow
1 \ 2 \ 3 \ 4 \ 5 \ 6 \ 7 \ 7 \ 9 \ 10 \ 11 \ 12 \ 13 \ 14 \ 15 \ 16 \ 17 \ 18 \ 19 \ 22 \ 23 \ 24 \ 27 \ 28 \ 30 \ 31 \ 31	81 74 79 82 86 63 64 63 67 77 77 78 77 67 65 67 66 67 66 66 66 66 66	3 P 4 P 2 P 3 P 1 P 3 P 2 P 4 P 3 P 2 P 4 P 3 P 2 P 4 P 3 P 2 P 4 P 3 P 2 P 4 P 3 P 2 P 4 P 3 P 4 P 3 P 4 P 3 P 4 P 3 P 4 P 3 P 4 P 3 P 4 P 4 P 5 P 5 P 6 P 6 P 7 P 8 P 8 P 8 P 8 P 8 P 8 P 8 P 8	66 44 43 47 62 63 44 49 49 49 42 54 55 67 61 49 53 68 51 44 56 57 48 49 57 58 58 58 58 58 58 58 58 58 58	5 A 7 A 6 A 6 A 6 A 6 A 6 A 6 A 6 A 7 A 6 A 12 P	73.5 59 61 64.5 74 68 53.5 56.5 56.5 57.5 64 62 63.6 62 63.5 73.5 62 63.5 57 60.5 62 56.5 57 60.5 62 56.5 57 60.5 60	67 69 73 80 80 80 80 74 91 80 72 69 69 84 81 73 78 82 73 74 85 71 74 72 74 76 91 72 72 73 74 75 76 77 77 77 77 77 77 77 77 77 77 77 77	10 12.7 7.9 6, 2 5, 5 4, 8 11.7 11.0 9, 6 0, 6 8, 7 6, 8 10.7 7.1 8, 4 11.0 0 4.1 9, 6 0.4 11.0 11.0 0 11.0 11.0 11.0 11.0 11.	sw w w ne ne nw sw e nw ne	172 65 32 62 187 45 26 114 98 66 115 115 87 76 59 89 140 134 50 71 73 104 83 29 78 51 181	18 4 6 5 13 17 7 6 18 12 10 13 14 9 13 11 17 20 6 4 4 15 22 11 6 9 11 11 11 11 11 11 11 11 11	8 A 6 A 0 A 0 A 0 A 0 A 0 A 0 A	7 P 4 P 4 A 8 A 12 P 9 P 6 P	.44 .16 .03 .46 .72 .50 .12 .28 .41	

^{*} Based on least time required to blow one mile.

LLOYD S. COHAN., Observer

MONTHLI	SUMM	
SEPTEMBER 1949	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum .30,57,12th Minimum .29,42, 23rd Alean semi-daily .30,045	30.45 29.57 30.00	30.65, 1924 28.41, 1938
Range	.88	1.99, 1938; .57, 1910
Air Temperature, in Degrees F.		
Highest 86, 5th Lowest 35, 26th Alean 61.7 Range 51 Highest mean daily 75,5 14th lowest mean daily 51, 25th, 26th Mean maximum 71.7 Alean minimum 51.7 Greatest daily range 36, 3rd Least daily range 6, 28th Degree-days of heating load for month 143 Degree-days cumulative for season 143	87.7 33.2 61.7 54.5	97, 1895, 1929 24.5, 1914
Precipitation, in inches		
Precipitation	4.24	14.55, 1938; .52, 1914
Number of days with .01 or more	10	16, 1933; 3, 1903
Wind, in miles		
Total movement 2643 Greatest daily movement 187, 6th Least daily movement 14, 29th Mean hourly velocity 3.7	3271	4,686, 1896; 1,414, 1894
Maximum velocity	25.9	80, 1938
Wind, direction		
Prevailing direction WNW North, days 5 Northeast, days 7 East, days 7 Eoutheast, days 0 South days 5 Southwest, days 7 West, days 2 Northwest, days 3	WSW	
Wealher		
Nean relative humidity, percent	73.6 49.9 10 10 10 201	70, 1934: 23, 1947 19, 1932: 2, 1907, 1928 19, 1908; 3, 1889 20, 1934; 3, 1941, 1948 255, 1916; 106, 1934
bright sunshine	54.9	
First Frost	Sept. 21	Oct. 13, 1909; Aug. 22, 1894-95



Meteorological Series

Bulletin No. 730

October 1949

Meteorological Observations

FOR

OCTOBER 1949

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

		Te	emper	ature		2.	, m		Wind			Precipita	ation	
	-	ximum		nimum		Mean Relative Humidity	Number of Hours Bright Sunshine	iling	ment	Max. Miles per Hour*	_			
Day	Degrees	Time	Degrees	Time	Mean	Mean Relati	Numb Bright	Preveiling Direction	Total Movement	Max. per H	Began	Ended	Water	Snow
1 2	66 64	3 P 4 P	44 36	7 A 7 A	55.0 50.0	66	11.5	ne n	55 48	9 6	1			İ
4	68 74	2 P 4 P	40 50	7 A 6 A	54.0 62.0	79 81	8.0 7.6	s sw	85 105	10			-	
4 5 6	75 68	3 P 4 P	60 40	1 A	67.5	79	6.6	s	146	18	6 A	7 A	.17	
7	66	4 P	44	7 A 2 A	54.0 55.0	72 86	10.2	sw	44 18	5 5	9 p	12 P	.15	
8	72 83	1 P 4 P	50 64	1 A 8 A	61.0 73.5	88	2.8	s	24	3 7	0 A	1 A	.13	
10	86	4 P	62	8 A 7 A	74.0	84	6.5 8.9	se w	37 48	4				
11 12	88 84	3 P 1 P	59 65	5 A 4 A	73.5 74.5	82 64	8.0	se	87	11 20	11 5	12 P	.04	
13	70	3 P	52	4 A 6 A	61.0	65	7.8	se nw	206 88	9	11 P	12 P	.04	
14 15	72 60	3 P 4 P	39 52	6 A	55.5	74	9.7	se	35	7	2	2 р	.27	
16	63	3 P	38	6 A 7 A	56.0 50.5	92 5 4	0 10.1	ne ne	71 117	9 8	2 A	2 P	.21	
17	69	2 P 3 P	34	5 A	51.5	60	8.2	se	69	11				
18 19	66 66	2 P	42 50	5 A 7 A	54.0 58.0	63	3.1 3.1	e e	135	17 15				
20	72	3 P	42	7 A	57.0	65	8.7	n	61	10				
21 22	75 73	4 P 3 P	37 55	5 A 6 A	56.0 64.0	63	8.9 1.3	s se	21 138	6 17	4 P	5 P	.01	
23	68	2 P	48	7 A	58.0	63	9.7	sw	177	18	• • •	, ,	.0.	
24 25	58 57	12 N 2 P	51 28	7 A 6 A	54.5 42.5	61 72	3.3 7.3	nw sw	117 43	14	9 р	12 P	.07	
26	63	2 P	45	2 A	54.0	75	7.5	W	116	28	0 A	7 A	.98	
27 28	55 63	4 P 2 P	36 31	7 A 7 A	45.5 47.0	63 70	9.2 9.4	nw	127 91	10				
29	68	3 P	49	8 A	58.5	78	5.6	se n	37	1				
30	61 64	8 P 3 P	46 48	0 A 12 P	53.5 56.0	98 85	$\frac{0}{2.7}$	sw n	20 90	7 24	3 P 0 A	4 P 10 A	.02	
J1 (01 1	I	10	14 F	50.0	0.5	2.1	- 11	20	24 1	U A	IU A	./4	_

^{*} Based on least time required to blow one mile.

LLOYD S. COHAN., Observer

30.53 29.42 30.06 1.11	Extremes 30.68, 1929 29.00, 1926 1.47, 1926; .76, 1899
29.42 30.06 1.11 79.4	29.00, 1926
29.42 30.06 1.11 79.4	29.00, 1926
79.4	1.47, 1926; .76, 1899
50.5 56.0	90.5, 1908 17, 1936 57.4, 1947; 43.2, 1890
3.29	8.81, 1911; .01, 1924
9	15, 1913; 1, 1897, 1924
4074	5,467, 1910; 2,540, 1894
5.4 29.5	42, 1937
W	
69.0 48 10 10 11 175.0 51.3	67, 1932; 18, 1924 21, 1938; 1, 1911 17, 1924; 3, 1938 19, 1896; 3, 1924 232, 1923, 1938; 91, 1913
	9 4074 5.4 29.5 W

Meteorological Series

Bulletin No. 731

November 1949

Meteorological Observations

FOR

NOVEMBER

1949

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

		Те	mpera	ature		ty	· s		Wind			Precipita	ation	
	Maximum I		Maximum Minimum			fumidi	f Hour nshine			S				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5	59 59 62	2 P 3 P 2 P	39 39 33	8 A 6 A 5 A 8 P	49 49 47.5 47.5	60 72 83	7.3 6.4 3.7	w s ne	166 131 49 164	11 13 8 22	2 A 5 A	3 A	.01	
5 6 7	52 45 42 47	1 A 12 N 3 P 9 P	43 33 31 32	8 P 12 P 2 A 0 A	36.5 39.5	80 67 74 78	4.5 5.4 5.1 2.0	nw nw nw se	115 79 38	13 10 8	5 A 1 P	6 A 2 P	.01	
8 9 10	68 56 64 67	2 P 2 P 4 P	40 43 31 52	6 A 8 A 1 A 8 A	54 49.5 47.5 59.5	72 69 76 85	8.9 7.5 6.9 3.4	w nw se	145 115 77 119	17 7 13 14	6 A	8 A	.03	
11 12 13 14	58 48 63	2 P 7 P 1 P	40 39 40	8 A 8 A 11 P	49 43.5 51.5	62 85 77	0 0 4.6	se ne s nw	87 72 97	11 11 18	10 P 0 A	12 P 2 A	.31	
15 16 17 18	57 54 46 42	2 P 12 N 1 P 3 P	30 32 28 28	8 A 1 A 6 A 7 A	43.5 38 37 35	81 75 76 84	8.5 8.3 5.2 4.9	w nw n ne	30 31 44 35	8 11 15 9				
19 20 21	46 48 43	1 P 2 P 11 A	26 38 28	0 A 2 A 12 P	36 43 35.5	72 76 73	4.7 2.2 6.7	s s w	96 164 104	12 18 13	1 A	2 A	.04	
22 23 24 25	31 44 38 44	1 P 1 P 1 P	18 18 28 35	12 P 0 A 7 A 2 A	24.5 31 33 39.5	68 56 61 95	7.8 5.1 4.6 0	nw se sw nw	249 210 143 49	25 25 9 20	2 A	9 A	.25	
26 27 28	37 22 32	0 A 12 P 2 P	25 12 22	S A 4 P 0 A	31 17 27	62 85 91	8.4 0 0	ne ne ne	361 101 134	32 9 15	8 A 6 P	12 p 7 p	.23	2.3
29 30 31	39 49	8 A 1 P	32 29	0 A 7 A	35.5 39	100 83	0 6	n w	48 60	8	5 A	5 P	.58	

^{*} Based on least time required to blow one mile.

LLOYD S. COHAN, Observer

NOVEMBER, 1949	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum. 30.299, 12th, 13th Minimum 29,376, 25th Mean semi-daily 29.877	30.59 29.30 30.05	30.87, 1932 28,73, 1904
Range	1.29	1.84, 1904; 89, 1943
Air Temperature, in Degrees F.		
Highest 68, 8th Lowest 12, 27th Mean 40 Range 56 Highest mean daily 59,5, 11th Lowest mean daily 17, 27th Mean maximum 48.7 Mean minimum 32.1 Greatest daily range 33, 10th Least daily range 7, 29th Degree-days of heating load for month .742 Degree-days cumulative for season 1151	66.0 12.9 38.9 53.0	75, 1924, 1938 4, 1938 44.1, 1931; 33.6, 1901
Precipitation, in inches		
Precipitation	3.41 2.34	8.64, 1927;63, 1917 13.50, 1938
Maximum precip. in 24 hrs 0.58., 29th Number of days with .01 or more11	9	22, 1921; 2, 1904
Wind, in miles		
Total movement.3133Greatest daily movement361, 26thLeast daily movement30, 15thMean hourly velocity4.6Maximum velocity32, 26th	4531 6.3 30.3	5,978, 1096, ;2,589, 1889 48, 1948
Wind, direction		
Prevailing direction NW North, days 3 Northeast, days 6 East, days 0 Southeast, days 4 South, days 1 West, days 5 Northwest, days 7	WNW.	
Weather		
Mean relative humidity, percent	70.6 55.1 6 9 15 121	78, 1947; 34, 1917 15, 1903; 0, 1948 16, 1912; 4, 1889, 1930, '32, '35 24, 1927; 9, 1905, 1917 821, 1903; 44, 1948
bright sunshine 47.0 Thunder and lightning First Frost.	41.3 Nov. 6	Oct. 10, 1925; Dec. 13, 1941



Meteorological Series

Bulletin No. 732

December 1949

Meteorological Observations

FOR

DECEMBER

1949

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

		Te	mpera	iture		Ą	m		Wind]	Precipita	tion			
	Maximum		Mir	Minimum		nimum		Iumidi	f Hour: nshine			92				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow		
1 2 3 4 5 6	43 35 33 30 31	12 N 12 N 2 P 2 P 3 P	26 27 18 4 23	6 A 3 A 12 P 6 A 9 P	34.5 31.0 25.5 17.0 27.0	75 93 70 83 91	5.4 0 7.9 6.7 0.2	N NE N NW W	90 45 222 55 51	17 8 17 5.5	10 A 0 A 6 P 0 A	12 P 3 A 12 P 2 A	.22 .08 .14 .03	3.5 1.5 1.7 .3		
6 7 8 9 10	25 21 29 28 21	10 A 4 P 1 P 2 P 3 P	10 9 16 8 - 7	12 P 0 A 6 A 8 A 6 A	17.5 15.0 22.5 18.0 7.0	61 89 60 74 78	7.4 0 7.6 6.4 7.4	NW NW NW W NW	217 44 204 108 21	66 32 18 15 1	6 A	12 N	.17	.15		
11 12 13 14 15	34 49 50 38 30	12 P 7 P 6 A 2 P 2 P 3 P	12 34 33 23 15	0 A 1 A 12 P 12 P 8 A	23.0 41.5 41.5 30.5 22.5	95 100 81 71 72	0 0 0 4.6 6.7	W SW N W	13 89 136 86 63	1 12 15 15 15	12 N 7 A 0 A	9 P 12 P 10 A	.07 .68 .38			
16 17 18 19 20	33 40 48 51 49	3 P 6 P 3 P 3 P	10 15 33 39 31	8 A 8 A 0 A 10 P 8 A	21.5 27.5 40.5 45.0 40.0	86 73 88 88 80	7.3 7 0 1.5 3	NW SE SE NW SE	20 70 168 109 58	11 11 8 11 13	5 P	6 P	.01			
21 22 23 24 25	51 61 57 32 38	2 P 3 P 10 A 2 P 2 P 7 P	38 46 34 24 16	4 A 1 A 12 P 7 A 7 A	44.5 53.5 45.5 28.0 27.0	84 87 70 64 72	0 4 0 4.5 7.5	SE SE NW NW SW	198 212 198 123 53	13 15 13 2 7	12 N	(ı P	.18			
26 27	37 58	8 P	27 48	7 A 12 P	32.0 53.0	95 100 59	0 0 5.5	SW SW	20 131 253	13 25 25	12 N 0 A 11 P 4 A	12 P 1 P 12 P	.12 .29 .01 .06			
28 29 30 31	40 37 29 37	1 P 9 A 1 P 3 P	30 18 14 14	12 P 12 P 7 A 7 A	35.0 27.5 16.5 18.5	59 62 48 58	3.5 1 8 8	N N NE N	253 302 282 110	25 25 15 7	# A	6 A	00,			

^{*} Based on least time required to blow one mile.

LLOYD S. COHAN, Observer

DECEMBER, 1949	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum. 31.104, 25th Minimum 29,707, 1st Mean semi-daily 30.259 Range 1,397	30.66 29.27 30.06 1.38	30.96, 1889 28.85, 1915 1.78, 1895; 1.01, 1892
Air Temperature, in Degrees F.		
Highest 61, 22nd Lowest -7, 10th Mean 29,9 Range 68 Highest mean daily 55, 27th Lowest mean daily 5, 10th Mean maximum 38.5 Mean minimum 22.2 Greatest daily range 28, 10th Least daily range 8, 5th, 24th Degree-days of heating load for month 1085.5 Degree-days cumulative for season	54.3 1.09 27.5 56.2	65.5, 1908 —22.5, 1917 36.9, 1891; 17.1, 1917
Precipitation, in inches		
Precipitation 2.44 Snow 8.5 Maximum precip. in 24 hrs. 0.68, 12th	3.39 8.50	7.77, 1901; .58, 1943 26.50, '02, T, 1891, 1943
Number of days with .01 or more13	10	17, 1902: 4, 1892, 1943
Wind, in miles		
Total movement. 3751 Greatest daily movement 302, 29th Least daily movement 13, 11th Mean hourly velocity 5.0 Maximum velocity 66, 6th	4710 6,3 31.4	6,694, 1925; 3,239,1918 48, 1938, 1943, 1944, 1948
Wind, direction		
Prevailing direction NW North, days 8 Northeast, days 3 East, days 0 Southcast, days 5 South, days 0 Southwest, days 4 West, days 3 Northwest, days 8	M.Z.M.	
Weather		
Mean relative humidity, percent. 77.6 Mean cloudiness, percent. 48.8 Number of clear days 12 Number of fair days 6 Number of cloudy days 13 Number of hours bright sunshine. 117.6 Percent of possible hours of	69.4 54.9 8 9 14 128	71, 1929; 39, 1919 15, 1890; 0, 1948 16, '09; 4, '89, '30, '31, '36, '38 25, 1948; 7, '09, '23 172, 1896; 27, 1948
bright sunshine 41.6 Thunder and lightning First Frost	45.2 Nov. 6	

ANNUAL SUMMARY

ANNUAL	Normal	Extremes
Barometer, in inches (Readings reduced to sea level) Maximum 31.104, Dec. 25th Minimum 29.160, Apr. 6th Mean semi-daily 30.022	30.81 28.95 30.01	31.05, 1920, 1943 28.41, 1938
Range	1.85	2.47, 1938; 1.38, 1933
Air Temperature, in degrees F. Highest .98, Aug. 9th, 10th Lowest 7, Dec. 10th Mean .56,9 Range .105 Highest mean daily .82, July 29th Lowest mean daily .5, Dec. 10th	95.7 —12.2 47.4 107.8	104, 1911 —26, 1904 49.8, 1921; 44, 1904
Lowest mean daily 5, bet. Roth Mean maximum 62.14 Mean minimum 41.39 Greatest daily range 41, May 5th Least daily range 3. Jan. 11th & Feb. 25th	57.8 36.5	
Precipitation, in inches		***************************************
Precipitation 33.61 Snow 49.05	43.70 47.78	59.00, 1938; 30.68, 1908 89.00, 1893; 24.50, 1919
Maximum precip. in 24 hrs 1.56, May 24th Number of days with .01 or more 115	124	146, 1945; 96, 1924
Wind, in miles		
Total movement	52,223 5.8 39.5	63,571, 1908; 36,257, 1894 80, 193 8
Wind, direction		
Prevailing direction WNW North, days .57 Northeast, days .56 East, days .16 Southeast, days .27 South, days .36 Southwest, days .71 West, days .48 Northwest, days .54	W.	
Weather	(7.6	
Mean relative humidity, percent.71Mean cloudiness, percent.50.2Number of clear days.111Number of fair days.138Number of cloudy days.116Number of hours bright sunshine.2154.9Percent of possible hours of	67.6 51.7 116 123 126 2,353	60, '98, '01, '02; 41, '08, '24 217, 1941; 59, 1927 182, 1912; 64, 1936 179, '01, '02; 71, 1910, 1941 3038, 1941; 1864, 1902
bright sunshine 47.8 Last snow Mar. 18th First snow Nov. 26th Last frost Apr. 29th First frost Sept. 26th	52.8 April 15 Nov. 6 May 14 Sept. 21	Mar. 14 '10; May 11, '07, '45 Oct. 10, '25; Dec. 13, '41 Apr. 14, '43; June 8, '32 Aug. 22, '94, '95; Oct. 13, '09

Meteorological Series

Bulletin No. 733

December 1950

Meteorological Observations

FOR

JANUARY

1950

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

		Те	mpera	ature		;y			Wind		1	Precipita	ition	
	Ma	ximum	Min	nimum		Iumidi	f Hour nshine			8				
Day	Degrees	Time	Degrees	Time	Меап	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5	35 35 51 62 58	3 P 8 P 8 P 3 P 0 A	25 25 33 50 42	3 A 0 A 3 A 7 A 12 P	30 30 42 56 50	80 90 99 88 79	0 0 0 1.7 0	NW W S S	32 9 70 189 113	2 2 13 13 18	4 P 6 A	5 P 10 A 4 P	.01 .20	
6 7 8 9	44 38 25 32 33	0 A 0 A 0 A 5 P 8 P	37 24 5 6 23	12 P 12 P 10 A 2 A 7 A	41.5 31 15 19 28	100 78 55 72 94	0 5.6 8.2 0.5 0	N N N S	60 256 339 89 36	10 28 28 9 5	5 A 0 A	12 P 5 A	.97 .27	
11 12 13	41 32 34 59	6 A 4 P 12 P 1 P	25 15 23 34	12 P 8 A 0 A 0 A	33 23.5 28.5 46.5	56 63 99 7 3	2.2 7.0 0 3.7	N NE N W	333 91 28 320 189	40 13 5 40 9	7 A 5 A 7 P	9 P 6 A 8 P	.29	
15 16 17 18 19	39 45 36 48 26	2 P 11 A 1 P 3 P 0 A	27 31 27 29 16	8 A 12 P 8 A 4 A 7 A	33 38 31.5 38.5 21	73 61 56 83 60	4.1 0 4.1 2.6 5.7	W W SE W	223 160 201 147	25 13 25 7	1 A 2 A	8 A 8 A	.08	1.1
20 21 22 23 24	25 18 32 39 37	3 P 1 P 8 P 4 P 9 A	13 13 17 31 29	8 A 8 A 0 A 2 A 12 P	19 15.5 24.5 35 33	68 86 96 99	8.5 0.2 0.3 0	W S N W	120 49 15 22 74	22 6 2 4 10	3 A 1 A 10 A	6 P 7 P	.08 .10	1.5 1.1
25 26 27 28	34 65 45 32	3 P 1 P 0 A 2 P	28 33 22 14	6 A 0 A 12 P 8 A	31 49 33.5 23	100 93 68 66	0 4.3 3.8 6.8	NE SW N	108 123 271 86 99	5 25 25 5 12	0 A	9 A	.10	
29 30 31	52 46 27	3 P 0 A 0 A	29 27 20	1 A 12 P 8 A	40.5 36.5 23.5	98 50 95	2.2 3.1 0	SE NW NW	224 160	22 17	4 A	7 P 11 A	.42	3.4

^{*} Based on least time required to blow one mile.

LLOYD S. COHAN, Observer

MONITILI	3011111	7 11 C 1
JANUARY, 1950	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum. 30.696, 12th Minimum. 29,581, 14th Mean semi-daily 30.252	30.70 29.20 30.07	31.00, 1927 28.55, 1913
Range1.115	1.50	2.18, 1913; 0.97, 1896
Air Temperature, in Degrees F.		
Highest 65, 26th Lowest 5, 8th Mean 32,3 Range 60 Highest mean daily 53, 4th Lowest mean daily 8, 8th Mean maximum 39,5 Mean minimum 24,8 Greatest daily range 32, 26th Least daily range 5, 21st Degree-days of heating load for month 1015 Degree-days cumulative for season 3251	51.1 -7.1 24.2 58.2	66.0, 1932 —26.0, 1904 34.2, 1913; 13.9, 1918
Precipitation, in inches		
Precipitation .4.33 Snow .7.1 Maximum precip. in 24 hrs. 1.09, 10th	3.61 13.32	7.15, 1898; 1.07, 1896 33, 1898, 1923; 1.50, '08
Number of days with 01 or more17	11	15, '93, '20; 4, 1901
Wind. in miles		
Total movement 4036 Greatest daily movement 339, 8th Least daily movement 9, 2nd Mean hourly velocity 5.4 Maximum velocity 40, 11th, 14th	5055 6.8 32.7	7770, 1908; 2896, 1895 47, 1938
Wind, direction		
Prevailing direction NW North, days 9 Northeast, days 4 East, days 0 Southeast, days 2 South, days 4 Southwest, days 1 West, days 7 Northwest, days 4	WNW	
Weather		
Mean relative humidity, percent. 79.9 Mean cloudiness, percent. 75.1 Number of clear days 3 Number of fair days 9 Number of cloudy days 19 Number of hours bright sunshine. 74.6 Percent of possible hours of bright sunshine 25.4 Thunder and lightning First Frost.	70.1 55 9 13 137 46.7	78, 1932; 37, 1908 20, 1944; 1, 1949 18, 1890; 1926; 3, 1916, 1944 22, 1923, '31, '49; 4, 1920, '39 214, 1920; 50,4, 1949

REMARKS

While the month produced no record highs for mildness it did couple a low heating load to the season's already low figure which was appreciated by the householder. At the mean midpoint of the heating season which fell on the 22nd, less than 3000 degree days had accumulated. Late season severe weather which could follow can still bring the seasonal figure up, but will likely run less than average for the season.

Precipitation at an accumulated 4.33 inches of water is somewhat above the mean of 3.61 inches for the month and will aid in replacing the ground water deficit which has been accumulating over the past 18 months of below mean rainfall.

W. H. Thies, Extension Professor of Horticulture, contributes the following report "There has been little visible swelling of peach buds in the University orchard thus far and the number of live buds at present gives promise of a good crop. There is reason to believe, however, that sub-zero temperatures after this date would cause more damage than in early winter. More than 800 hours of temperatures below 45° are required to bring peach trees out of their winter rest period, and thus set the stage for a response to warmer weather. Around 40° they go through their rest period most rapidly. Above 45°, and also at temperatures much below freezing, the process is slowed up. Because of this situation, peach buds in Massachusetts have not responded as much to the mild January weather as one might suppose.

A few comments concerning other fruits may also be of interest. Apple and pear buds are still apparently in a strictly dormant stage. It is doubtful if any bud injury has occurred. An occasional case of bark splitting on the trunk may be traced to late stimulation and retard maturity. All raspberry varieties are probably out of their winter rest period, and there has been some evidence of tip-killing on canes of tender varieties, such as Washington. Strawberry plants, if protected by a layer of suitable mulch material, and in good condition. The absence of a snow cover during most of the winter season, has left unmulched plantings exposed to rather wide extremes of temperature. This condition may be reflected in damaged plants and a smaller crop of berries next June."

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 734

February 1950

Meteorological Observations

FOR

FEBRUARY

1950

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N.

Longitude, 72° 31′ 48″ W.

Height of barometer above ground, 36 ft. Above sea level, 253.5 ft.

Height of wind instruments, 67 ft.

Time used, E. S. T.

		Те	mpera	ture		, A.	m		Wind		. 1	Precipita	ition	
	Maximum		Mir	Minimum		Iumidi	f Hour nshine		t	.9				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7	31 35 34 35 32	2 P 2 P 1 A 1 P 4 P	20 27 22 18 15	0 A 2 A 8 P 8 A 8 A	25.5 31 30 26.5 23.5	90 100 51 59 44	3.6 0 7.3 4.5 7.1	NW NW W N	33 37 306 196 143	5 5 32 28 11	1 A	8 A	.21	1.7
8	34 33 26	2 P 2 P 4 P	15 26 - 4	0 A 3 A 8 A	24.5 29.5 11	77 57 60	0 7.5 7.8	NE NE N	21 160 127	9 25 13	11 A	6 P	.16	1.1
9 10 11	46 38 39	12 N 4 P 4 P	20 29 32	0 A 7 A 5 A	33 33.5 35.5	94 73 93	0 5.6 0.6	SW NE NW	147 142 34	20 13 6	l A 8 A	4 P 2 A 9 A	.02	1.7 0.1 T
12 13 14 15 16 17 18	43 33 29 33 39 42 40	2 P 4 P 12 P 4 P 3 P 2 P 3 P	26 25 24 26 28 28 31	8 A 4 A 8 P 0 A 8 A 5 A 8 A	34.5 29 26.5 29.5 33.5 35 35.5	75 80 92 100 89 87 67	6.4 2.3 0 0 4.3 6.1 4.2	N N N N	48 79 112 175 75 34 82	10 10 15 18 9 11	7 P 0 A 0 A 6 A	12 P 12 P 12 P 11 A	1.3 .90 .83 .05	1.4 9.1 1.0 0.7
19 20 21 22 23 24	41 21 20 16 29 41	2 P 0 A 2 P 2 P 3 P 5 P	26 - 4 - 8 - 8 - 8 11 19	0 A 12 P 6 A 8 A 0 A 7 A 12 P	33.5 12.5 6 12 20 30 17	85 58 49 87 89 71 58	2.5 9.3 8.7 1.8 0.9 4.7 8.6	NW E E E E NW NW	147 420 133 110 231 206 373	22 28 9 14 28 28 32	7 A 10 P 0 A 0 A	8 A 12 P 12 P 9 A	.04 .09 .65 .19	0.3 0.6 5.2 0.3
25 26 27 28	24 25 17 30	1 P 2 P 4 P 1 P	10 - 2 - 4 - 5	7 A 6 A 7 A	17 11.5 6.5 12.5	58 67 60 68	7.7 9.3 6.6	NW N SW	128 263 131	22 25 14	5 P	12 Р	.15	0.6

^{*} Based on least time required to blow one mile.

FEBRUARY, 1950	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum. 30.77, 8th Minimum. 29.45, 24th Mean semi-daily 30.13 Range 1.32	30.63 29.24 30.04 1,38	31.05, 1920 28.56, 1895 1.89, '00, '08; .88, '13, '31
Air Temperature, in Degrees F.		
Highest .46, 9th Lowest 8, 21st Mean .24.5 Range .54 Highest mean daily .34.5, 9th	50.5 7.4 23.7 58	65.0, 1930 -23, 1943 32.6, 1890; 11.6, 1934
Lowest mean daily 0, 20th Mean maximum 32.4 Mean minimum 16.4 Greatest daily range 35, 28th Least daily range 5, 14th Degree-days of heating load for month 1132 Degree-days cumulative for season 4383	32.8 14.5	
Precipitation, in inches	ĺ	
Precipitation 3.99 Snow 23.8 Maximum precip. in 24 hrs 94, 14th	3.19 13.99	8.12, 1900; 0.62, 1901 48.75, 1893; 0.50, 1937
Number of days with .01 or more13	10	15, '93, 1920; 4, 1901
Wind, in miles		
Total movement	7.1 31.7	6445, 1896; 3402, 1949 50, 1946
Wind, direction		70, 1710
Prevailing direction N North, days 12 Northeast, days 3 East, days 3 Southeast, days 1 South, days 0 Southwest, days 2 West, days 1 Northwest, days 6	WNW	
Weather		
Mean relative humidity, percent	66.6 50.9 10 8 10 158	66, '90, '27; 31, 1905 19, 1941; 2, 1927 16, 1920; 2, 1936 16, 1894; 2, 1920 221, 1924; 110, '27, '38
Thunder and lightning First Frost.		

With the exception of the 8th the mild weather of the previous month continued until the 20th when a precipitous drop in temperature occurred which carried through to the end of the month. This greatest daily wind movement during the month combined with this temperature change provided the most uncomfortable winter conditions to that point of the season. The low of —8 degrees on the morning of the 21st however only approximated the —7.1 mean lowest temperature for the month, and the mildness earlier provided a mean of 24.5 degrees against an average mean of 23.7. The heating load continued very moderate at 1132 degree-days for the month to accumulate 4383 degree days for the season. The lengthening days with fair to clear cloud conditions helped offset the cold air masses and heavy nocturnal radiation on clear nights.

The precipitation of 23.8 inches of snow which yielded 3.99 inches of water provided a margin above the mean of 3.19 inches for the month which will eventually be available to the ground water

supply.

William G. Colby, Professor of Agronomy, contributes the following report: "The relatively mild temperatures during the first 19 days of the month, coupled with somewhat more than average rainfall, helped tremendously in replenishing badly depleted soil moisture reserves. The last 10 days of the month were colder than normal but the ground was covered with a heavy blanket of snow, therefore preventing deep penetration of frosts into the soil."

"This means that soil moisture reserves will be still further replenished when the present snow cover melts. Field crop conditions to this point in the season continue favorable. A similar condition prevailed last year. Our summer drought would have been more serious had it not been for the absorption by the soil water produced

from a comparatively heavy, late winter fall of snow."

Meteorological Series

Bulletin No. 735

March 1950

Meteorological Observations

FOR

MARCH 1950

II. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments. 67 ft Time used, E. S. T.

		Te	empera	ature	_ +	<u>,</u>			Wind		I	Precipita	tion	
	Ma	ximum	Min	nimum		Humidi	f Hour nshine		t	s s				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1	37	2 P	19	12 P	28	76	1.6	W	217	25	0 A 11 A	1 A 1 P		Т
2 3 4 5 6 7 8 9	20 15 26 44 42 37 57 45 25	12 N 1 P 4 P 4 P 2 A 4 P 12 P 0 A 4 P	$ \begin{array}{c c} 7 \\ -2 \\ -9 \\ 11 \\ 11 \\ 4 \\ 27 \\ 15 \\ 14 \end{array} $	12 P 6 A 6 A 0 A 12 P 7 A 0 A 12 P 5 A	16 6.5 8.5 27.5 26.5 20.5 42 30 19.5	56 50 55 68 52 53 91 56 53	8.2 9.3 9.7 6.0 10.1 8.0 0 4.7 9.6	NW NW NW S NW SW S NW NW	305 331 42 193 338 81 107 455 387	32 25 8 18 32 8 40 40 32	8 A 9 A	11 p 8 p	.72	T
11 12 13 14 15	31 45 33 34 42 36	6 P 12 N 0 A 3 P 12 N 3 P	14 24 23 21 16 18	6 A 0 A 12 P 8 A 7 A 7 A	22.5 34.5 28 27.5 29 27	79 73 85 59 63 56	3.1 8.2 0 8.8 7.8 9.5	S NW N NW W	118 108 119 240 236 126	12 18 14 20 39 12	7 P 7 A 0 A 11 A	10 P 12 P 8 A 12 N	.02 .15 .07	1.2 1.2 T
16 17 18 19 20	38 35 39 49	4 P 5 A 2 P 4 P	18 18 18 14 19	6 A 12 P 7 A 7 A	28 26.5 26.5 34	47 64 53 62	9.0 3.6 10.5 9.5	N NW NW S	120 120 334 212 75	12 40 18 7	11 p 0 a	12 P 6 A	.03	.5 1.8
21 22 23 24 25	42 47 43 43	11 A 2 P 3 P 3 P 11 A	29 24 37 35 31	5 A 7 A 8 A 5 A 6 A	35.5 35.5 40 39 38.5	78 84 92 70 66	.3 7.2 0 0 5.2	N E N NW N	87 85 185 206 137	17 11 14 22 18	11 A 3 A 0 A	5 P 12 N 1 A	.08 .85 .01	.7
26 27 28 29 30 31	52 41 65 62 41 57	2 P 2 P 4 P 2 P 4 P 4 P	27 37 37 40 28 20	5 A 0 A 4 A 8 P 3 A 6 A	39.5 39 51 51 34.5 38.5	56 96 95 83 56 49	9.2 0 4.9 4.5 9.5 11.4	N SE S N N	75 66 67 127 378 126	11 10 13 25 25 25 14	8 A 6 P 5 A	12 P 7 P 5 P	.07 .01 .58	

^{*} Based on least time required to blow one mile.

LLOYD S. COHAN, Observer

MARCH, 1950	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum. 30.557, 4th Minimum. 29.188, 23rd Mean semi-daily 29.945 Range 1.369	30.57 29.24 30.04 1.34	31.05, 1943 28.47, 1914 2.10, 1914; 0.85, 1915
Air Temperature, in Degrees F.		
Highest. 65, 28th Lowest -9, 4th Mean 30.7 Range. 74 Highest mean daily 47.5, 28th Lowest mean daily 3.5, 3rd Mean maximum 49.3 Mean minimum 20.6 Greatest daily range 37, 31st Least daily range 4, 27th Degree-days of heating load for month 1085 Degree-days cumulative for season 5468	63.8 6.1 34.4 57.6 43.2 25.4	85, 1929 -7.5, 1906
Precipitation, in inches		
Precipitation 2.67 Snow 5.6 Maximum precip. in 24 hrs. .85, 23rd	3.70 7.47	7.89, 1942; 0.12, 1915 27, 1899; 0, 1921
Number of days with .01 or more 13	11	17, 1890, 1913, 1936; 3, 1915
Wind, in miles		
Total movement	7.7 31.2	8182, 1896; 3006, 1905 48, 1932, 1939, 1941, 1942
Wind, direction		
Prevailing direction. NW North, days 10 Northeast, days East, days 1 Southeast, days 1 South, days 5 Southwest, days 1 West, days 2 Northwest, days 11	WNW	
Weather		
Mean relative humidity, percent. 67 Mean cloudiness, percent. 51.1 Number of clear days 14 Number of fair days 7 Number of cloudy days 10 Number of hours bright sunshine 186.4 Percent of possible hours of bright sunshine 50.4 Thunder and lightning 8th, 28th First Frost.	64.2 51.8 11 10 10 199 53.6	68, 1901; 27, 1915 22, 1924; 3, 1901 17, 1900; 1, 1943 21, 1901; 1, 1915 292, 1924; 93, 1901

The month was the coldest March since 1941 and continued the cold weather ushered in by the abrupt change which occurred February 20. The maximum of 65° was only a moderate highest temperature for the month, while the low of -9° was well below the mean of +6.1. The heating load of 1085 degree days was also the highest since the same month of 1941, while the cumulative total of 5468 remained below the normal expected at the month's end.

High winds occurred on three days of the month with a maximum velocity of 40 mph. Total wind movement however, at 5683, was below the mean of 5764. Precipitation at only 2.67" was more than an inch below the mean for the month and may be expected to have its effect on seasonal deficiency shown more fully after crops are planted. Thunder and lightning was recorded on both the 8th and

the 28th, the two milder days near each end of the month.

W. H. Thies, Extension Professor of Horticulture, contributes the following report: "A survey of Massachusetts peach orchards reveals severe damage from cold weather. All of the peach buds in some orchards have apparently been destroyed while a scattering of uninjured buds is found in a few varieties. In a Worcester County orchard examined April 5, about 10% of the buds on one odd variety were alive. Buds of other varieties showed brown centers. Several growers have reported enough live buds here and there to suggest a very light crop this season.

In the University orchard, J. S. Bailey reports no live buds at all on Elberta, Halehaven, Golden Jubilee and many other varieties. A

few live buds were observed on the Greensboro variety.

The present peach situation is not surprising in view of the wide variation in winter temperatures, and the fact that we are near the northern limit of the peach belt. Peaches are among the most tender of the fruits grown in this area. But if peach growers are able to harvest a good crop three years out of four under our climatic conditions the gamble in setting a peach orchard is justifiable provided a better than average soil and site are selected."

Meteorological Series

Bulletin No. 736

April 1950

Meteorological Observations

FOR

A P R I L

II. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

		Te	mpera	iture .		٤	œ		Wind		I	Precipita	tion	
	Max	imum	Min	nımum		lumidi	Hour suine			0				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunsuine	Prevailing Direction	Total Movement	Max Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 11 12 11 12 13 14 15 16 17 18 19 20 21 10 10 10 10 10 10	52 55 75 52 48 52 46 51 46 55 56 77 74 70 74 70 75 77 77 77 77 77 77	2 P 3 P 5 P 2 P 8 A 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 4 P 4 P 4 P 4 P 4 P 4 P 4 P 4 P 4	22 33 33 50 25 23 28 34 25 27 30 30 30 33 31 42 40 44 37	6 A 12 P 0 A 7 A 12 P 12 P 5 A 6 A 6 A 4 A 12 P 0 A 0 A 1 5 A 6 A 6 A 12 P 6 A 6 A 6 A 6 A 6 A 6 A 6 A 6 A 6 A 6	37 44 44.5 62.5 45.5 35.5 35.5 40 40 38 31.5 39 40 45.5 42.5 56 57 49.5 47	56 67 88 79 50 41 42 54 45 62 54 100 75 63 52 50 60 60 62	9.6 6.0 6.1 0 9.1 11 11.4 9.1 10.6 2.4 6.3 0 6.9 7.8 12.1 6.6 10.9 7.8 0 6.1	SE W SE S NW NW N N N N N N N N N N N N N N N N	120 186 97 175 237 260 241 238 236 199 91 288 78 251 183 67 186 120 147 145	13 18 10 20 28 25 22 18 25 20 18 25 11 22 20 20 10 25 17 12 17	1 A 6 A 0 A 5 A 0 A 0 A 0 A	6 A 12 P 1 A 3 P 7 P 12 P 6 A 8 P	.10 .17 .04 .16	6,6 .7
22 23 24 25 26 27 28 29 30	50 53 56 62 47 64 61 43 60	1 P 6 P 6 P 4 P 0 A 4 P 5 P 0 A 4 P	36 32 31 29 42 41 44 33 29	3 A 2 A 5 A 6 A 8 A 0 A 5 A 12 P 6 A	43 42.5 43.5 45.5 44.5 52.5 52.5 38 44.5	61 86 57 61 90 78 83 82 61	6.1 1.2 10.4 10.6 0 5.9 4.0 0 10.5	NW S N SW NE SW S S S S S S S	150 89 127 75 122 102 121 94 79	20 14 12 14 9 18 9 10 15	4 P 4 A 10 P 0 A 8 A	8 P 12 N 12 P 1 A 3 P	.04 .31 .08 .04 .14	.3

^{*} Based on least time required to blow one mile.

LLOYD S. COHAN. Observer

APRIL, 1950	Norma!	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum. 30.31, 10th Minimum. 29.66, 23rd Mean semi-daily 29.95	30.48 29.38 29.99	30.71, 1911 28.98, 1943
Range 0.65	1.11	1.52, 1930; .72, 1919
Air Temperature, in Degrees F.		
Highest 75, 4th Lowest 22, 1st Mean 43.8 Range 53 Highest mean daily 57, 4th	79.4 22.0 45.7 57.4	90 1941 8.5, 1923 52, 1921; 41, 1943
Lowest mean daily	56.8 34.8	
Precipitation, in inches		
Precipitation	3.35 2.17	6.89, 1929; .55, 1941 11, 1891; 0, 1910, 1934
Maximum precip. in 24 hrs	11	18, 1909; 3, 1892
Wind, in miles		
Total movement 4646 Greatest daily movement 288, 12th Least daily movement 67, 17th	5404 7.5	8208, 1908; 3853, 1917
Mean hourly velocity	31.4	40, 1935, 1938, 1940, 1945
Wind, direction		
Prevailing direction NW North, days 9 Northeast, days 1 East, days 0 Southeast, days 5 South, days 5 Southwest, days 2 West, days 1 Northwest, days 7	WNW	
Weather		
Mean relative humidity, percent. 66.5 Mean cloudiness, percent. 56 Number of clear days 10 Number of fair days 11 Number of cloudy days 9 Number of hours bright sunshine. 189.1 Percent of possible hours of bright sunshine 47.2	61.6 51.8 11 9 10 220 51.7	75, 1901; 34, 1925, 1927 23, 1941; 1, 1949 18, 1915; 2, 1894, 1901 22, 1901; 1, 1941 329, 1941; 103, 1901
17.25		Mar. 14, 1910; May 11, 1907, 45

This menth proved a little reluctant to usher in the spring season. After a high for the month of 74° on the 4th it was a full two weeks before any 70 temperature again appeared. The low of 22° on the first reached the mean lowest temperature for the month and the mean daily of 43.8° was nearly two degrees below the average. Householders continued to feel the dregs of winter as 637 degree days of heating load accumulated the seasonal total of 6105 degree days.

The 3.64 inches of precipitation added nearly 0.3 inch over the

mean to the moisture reserve for the coming growing season.

W. H. Thies, Extension Professor of Horticulture, contributes the following report: "Gool weather during April, with its retarding effect on fruit buds, has been favorable to fruit crops in Massachusetts. This applies to fruits of all kinds. The situation may be summed up as follows: The stage of bud development is about 10 days behind that of certain seasons in the recent past. McIntosh apple buds, now in the Pre-Pink stage in some areas, are better able to withstand freezing temperatures that if they were approaching full bloom. At these two stages there is a difference of at least 3° or 4° in susceptibility to frost damage. This may mean a big difference in the number of blossoms setting fruit. McIntosh records in Amherst show full bloom occurring, on the average, around May 15 with a range of nearly 10 days in either direction. May 20 is the estimated date of full bloom this year.

As this is being written, there has been negligible low temperature damage to apple buds, and the stage is set for at least a normal crop. Peach buds suffered severely during the winter in some orchards. This must be expected occasionally since we are near the northern limit of the peach belt, and peaches are among the most tender of the fruits grown in this area. In well located orchards, however, there were still enough live peach buds of certain varieties on May 1 to provide at

least a light crop.

Strawherry plants in mulched plantings came through the winter in good condition. Bloom is likely to be delayed enough to escape appreciable frost damage. Even though a few of the early blossoms do show black centers, it will mean only the loss of the earliest berries and not any extensive reduction in the number of quarts per acre."

Meteorological Series

Bulletin No. 737

May 1950

Meteorological Observations

FOR

MAY

1950

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

		Te	mpera	ature		, A	80		Wind		1	Precipita	tion	
	Ma	ximum	Min	nimum		Iumidi	f Hour nshine			20				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Вецяя	Ended	Water	Snow
1 2 3	46 68 71	3 P 4 P 4 P	43 42 43	6 A 6 A	44.5 55 57	87 65 53	9.1 10.6	SE S	89 127 118	17 14	6 A	12 N	.34	
4 5 6 7	73 59 85 73	5 P 4 P 5 P 12 N	43 51 49 49	6 A 9 A 1 A 9 P	58 55 67 61	74 75 76 41	8.1 0 2.6 12.7	S S NW	105 89 150 328	13 11 17 48			т	
8 9 10	58 74 66	4 P 4 P 6 P	34 34 47 50	5 A 5 A 2 A	46 54 56.5 57	42 46 77 53	13 11.3 0.2	SW S	399 114 157 141	28 17 14 20	11 P 0 A	12 P 1 P	.02 .38	
11 12 13 14	64 72 78 68	12 N 3 P 3 A 4 P	42 40 38	6 A 5 A 6 P 6 A	57 59 53	52 63 50	10.3 10.6 10.5 13.9	NW N NW N	52 118 161	32 20 20	7 P	8 P	.01	
15 16 17 18 19 20 21	62 72 75 54 51 63 76	5 P 4 P 3 P 2 P 2 P 1 P 4 P	40 50 45 48 45 45 45 38	5 A 6 A 5 A 6 A 5 A 0 A 5 A	51 60 51 48 54 57	74 67 64 88 88 69 59	5.2 8.5 10.9 0 0 5.7 12.5	NW N SW SE NE NE NE	80 140 92 83 82 179 165	6 9 18 10 7 20 20	3 A 1 A	12 P 2 P	.26	
22 23 24 25 26 27	70 71 67 64 79 80	2 P 2 P 11 A 1 P 5 P 2 P	35 42 56 57 55 49	5 A 4 A 7 A 3 A 5 A 5 A	52.5 56.5 61.5 60.5 67 64.5	54 72 89 89 68 61	13.5 8.8 0.6 0 11.8 13.4	SW S SE E N	118 144 78 74 66 87	18 13 28 10 8 12	4 A 0 A	12 P 5 P	.41 .45	
28 29 30 31	72 64 75 77	2 P 3 P 1 P 5 P 2 P	46 54 58 59	5 A 6 A 5 A 3 A 5 A	64.5 59 59 66.5 68	68 89 85 84	13.4 10.1 0.3 6.5 5.6	SW NW SW SW W	88 63 74 102	12 9 9 7 12	8 A 1 A	12 P 8 A	.37 .11	

^{*} Based on least time required to blow one mile.

MONTILL	JUNIMIAN	(1
MAY 1950	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.32, 9th Minimum 29,59, 13th Mean semi-daily 30.03 Range .83	30.39 29.51 29.96 0.88	30.62, 1936 29.10, 1938
Air Temperature, in Degrees F.		
Highest .85, 6th Lowest .34, 8th, 9th Mean .57,9 Range .51 Highest mean daily .64,5, 30th, 31st Lowest mean daily .43, 8th Mean maximum .68,6 Mean minimum .46 Greatest daily range .40, 9th Least daily range .3 1st Degree-days of heating load for month .230 Degree-days cumulative for season .6335	86.1 31.2 57.1 54.0	94.5, 1896, 1911 24.0, 1900
Precipitation, in inches		
Precipitation 2.77	3.60	7.44, 1931; .48, 1903
Snow	12	20, 1901, 1945; 5, 1903
Wind, in miles		
Total movement	4504 6.1 28.8	5946, 1907; 2180, 1894 45, 1935
Wind, direction		
Prevailing direction. WSW North, days. 5 Northeast, days 2 East, days. 1 Southeast, days. 3 South, days. 6 Southwest, days 6 West, days 2 Northwest, days 6	"	
Weather		
Mean relative humidity, percent 68.4 Mean cloudiness, percent 58.3 Number of clear days 13 Number of fair days 8 Number of cloudy days 10 Number of hours bright sunshine 226.3 Percent of possible hours of bright sunshine 50 Thun ler and lightning 50 Last Snow Last Frost	60.7 52.1 11 11 25.2 55.6	72, 1948; 30, 1923 21, 1944; 0, 1927 18, 1949; 5, 1923 20, 1927; 2, 1941 350, 1944; 137, 1927

The month made a valiant beginning to catch up the slowness of the season, providing a maximum of 85° on the 6th, only to have this mildness dispersed by a dust storm in the teeth of a 48 mile wind on the 7th. The mean daily then dropped bock into the 50's and high 40's until the last week of the month when seasonable warmth was again obtained. In the month 2.77 inches of precipitation fell; nearly $\frac{3}{4}$ inch below the mean and only two stormy periods recorded rainfall in excess of $\frac{1}{2}$ inch: en the 18th-19th 0.68 inches, and the 24th-25th 0.86 inches. The mean daily temperatures of 57.9° although 0.8° above average contributed to the 230 degree days of heating load which brought the season's total to 6335 at the month's end.

The last snow neted April 29, and the last frost April 30, in last month's bulletin, have not been disputed by the month's records.

W. H. Thies, Extension Professor of Horticulture, contributes the following report: "Weather conditions during May have had some interesting effects on plant growth. Lawns look unusually well and the hay crop is promising, thanks to quite ideal growing conditions. Peas, spinach and such cool weather crops have made excellent progress. But corn, beans, and others which thrive in hot weather have been marking time. Asparagus got off to a slow start but has made up for it since. Seldom have we seen strawberries look any more promising. Here is a fruit plant which seems to best under cool, cloudy conditions. We have prospects of a good strawberry crop. But instead of the first berries appearing around June 1 as they sometimes do in the Connecticut Valley, they are more lightly to appear around June 10.

Apple bloom was at least a week later than normal, and was spread over a longer period. The height of McIntosh bloom in Amherst was around May 22. Bee activity was frequently held up by cool or rainy weather. Honey bees are quite inactive when the temperature drops below 60 although bumblebees continue to fly at temperatures at least 10° lower. In spite of weather interference in bee activity, there were enough warm, sunny hours in most orchards to insure pollination. Contrary to a common notion, none of the fruits grown in this area except nuts, are wind pollinated. Bees of one kind or another are almost entirely responsible for the transfer of pollen from one blossom to another. And that means suitable weather for bee flight."

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 738

June 1950

Meteorological Observations

FOR

JUNE

1950

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W

Height of barometer above ground, 36 ft. Above sea level, 253 5 ft.

Height of wind instruments, 67 ft Time used, E. S. T.

		Те	mpera	ture		ty	6		Wind		P	Precipita	tion	
	Max	imum	Mir	imum		Humidi	of Hour inshine	co _	nt	**		1		
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Венап	Ended	Water	Snow
1	69 78	10 A	64	2 A 5 A	66.5 67.5	96	0 13.9	W NE	132 110	9	3 A	9 P	1.35	
3	79 67	1 P 3 P	59 53	1 A 8 A	69 60	74 73	5 10.3	sw nw	193 200	25 20	9 p 0 a	12 P 8 A	.03	
5	76	5 P	45	1 A	60.5	52	13	w	84	12	0.1		,	
6 7	84 89	1 A 5 P	46 53	5 A 5 A	65	58 62	8.2 10.3	SW	65 42	6				
8	90	3 P	60	6 A 5 A	75 75.5	62	9.5 11.2	SW	101 110	11				
9 10	90 85	1 P	61 63	5 A 4 A	74	81	6.9	S W	159	18	11 A	12 P	.81	ļ
11	71 74	4 P 4 P	57 42	6 A 5 A	64 68	61 59	12.7	NE NW	211 51	17				
13	78	2 P	49	5 A	63.5	59	11.1	W	125	10		_		
14 15	70 78	11 A 4 P	53 53	2 A 2 A	61.5 65.5	81 78	3.3	NW NW	79 48	7 7	4 P	7 Р	.06	
16	79	2 P	54	4 A	66.5 55.5	70	9.2 8.7	w	121 261	13	4.5	9 3	.61	
17 18	68 69	9 A 11 A	43 42	12 P 1 A	55.5	70 50	12.1	N NE	197	20	+ A			
19 20	60 76	1 P 2 P	41 53	1 A 5 A	50.5 64.5	82 72	0.3	W	13 96	12	10.A	[11 A	.01	
21	79	4 P	62	4 A	70.5	74	5.6	N	126	18	3 A	5 A	.07	
22 23	75 89	4 P 1 P	47 53	5 A 5 A	61 71	55 62	11.4	NW	98	$\frac{8}{10}$				
24	90	2 P	62	5 A	76	79	4.7	s	84	1.5	4 P	5 P	.03	
25 26	89 91	2 P 4 P	65	5 A 5 A	77 76	73	8.0	NW N	46 64	17				
27	86	2 P	63	5 A	74.5	69	6.0	s	157 173	18 20	2 P	4 P	.31	
28 29	77 82	5 P 1 P	52	5 A	67	61	9.1	NW	55	12	7 P	S P	Т	
30	85	4 P	63	5 A	74	71	9.5	s	113	8				

^{*} Based on least time required to blow one mile.

JUNE 1950	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30,302,7th Minimum 29,522,17th Mean semi-daily 29,902 Range 780	30.30 29.55 29.95 0.75	30.70, 1947 29.24, 1902
Air Temperature, in Degrees F.		
Highest 91, 26th Lowest 41, 19th Mean 66 Range 50 Highest mean daily 71, 26th Lowest mean daily 50.5, 4th, 19th Mean maximum 78.1 Mean minimum 54.1 Greatest daily range 41, 2nd Least daily range 5 1st Degree-days of heating load for month 68.5 Degree-days cumulative for season 6404	91.2 40.0 65.7 51.3	101, 1919 34, 1891
Precipitation, in inches		
Precipitation	3.75	9,68, 1922; 0,76, 1908
Snow. Maximum precip, in 24 hrs	11	17, 1922, 1945; 4, 1908
Wind, in miles		
Total movement 3535 Greatest daily movement 261, 17th Least daily movement 13, 19th Mean hourly velocity 4.9 Alaximum velocity 32, 17th	5.0 24.6	4571, 1908: 1409, 1906 48, 1939
Wind, direction		
Prevailing direction W North, days 3 Northeast, days 3 East, days 0 Southeast, days 5 South, days 5 Southwest, days 4 West, days 9 Northwest, days 6	WSW	
Weather		
Mean relative humidity, percent. 67.7 Mean cloudiness, percent 39.8 Number of clear days 11 Number of fair days 16 Number of cloudy days 3 Number of cloudy days 268.2 Percent of possible hours of bright sunshine 58.7 Thunder and lightning 10th, 17th, 27th Last Snow. Last Frost	66.9 51.1 10 12 8 257 54.0	71, 1903; 28, 1908 22, 1908, 1941, 1943; 0, 1949 23, 1912; 3, 1941 22, 1903; 1, 1923 362, 1908; 102, 1903

Meteorological Series

Bulletin No. 739

July 1950

Meteorological Observations

FOR

JULY 1950

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N.

Longitude, 72° 31′ 48″ W

Height of barometer above ground, 36 ft. Above sea level, 253.5 ft

Height of wind instruments, 67 ft

Time used, E. S. T.

-		Т	emper	ature		<u>.</u>	, m		Wind]	Precipita	ation	
	Ma	ximum I	Mi	nimum	i	Humidi	of Hour inshine	G0 _	ıt	les *				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Ведан	Ended	Water	Snow
1 1 2 3 4 4 5 6 6 7 8 8 9 10 11 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29	84 85 83 76 85 82 89 87 88 82 86 90 75 77 87 91 88 83 83 83 79 84 88 87 88 88 88 88 88 88 88 88 88 88 88	3 P 2 P 5 P 11 A 4 P 2 P 2 P 12 N 4 P 2 P 12 N 4 P 2 P 12 N 4 P 2 P 2 P 2 P 2 P 3 P 2 P 2 P 3 P 2 P 3 P 2 P 3 P 2 P 3 P 2 P 3 P 2 P 3 P 2 P 3 P 2 P 3 P 2 P 3 P 2 P 2	60 48 65 65 62 68 51 55 61 61 46 62 70 64 46 55 65 65 65 65 65 65 65 65 65 65 65 65	6 A A A A A A A A A A A A A A A A A A A	72 66.5 74 70.5 73.5 73 66.5 72 74 70.5 72 78 77 68 62.5 64 79.5 72 62 66 68 69 74 71 67.5 71 72	64 60 71 89 79 68 63 60 64 69 92 80 75 59 83 62 64 64 77 68 63 63 64 64 64 65 71 68 68 63 64 64 64 65 65 66 66 66 66 66 66 66 66 66 66 66	10.8 9.4 10.2 0.2 7.2 9.6 13.4 12.9 11.9 4.8 .45 6.5 8.2 12.1 11.3 3.2 8.4 9.5 12.6 0.0 13.3 14.9 14.0 12.9 12.1 12.9 14.9 11.9 11.9 11.9 11.9 11.9 11.9 11	N SW S S S NW N N SE S S NW S S S W W N N N N N N N N N N N	106 69 122 36 59 155 120 25 51 28 39 87 79 186 31 107 68 48 35 31 77 123 137 75 187	13 10 13 7 7 13 17 5 6 4 10 11 12 25 6 9 9 9 9 9 9 9 9 9 9 14 17 17 17 17 18 18 18 18 18 18 18 18 18 18 18 18 18	2 A 3 A 8 P 0 A 0 A 7 P 6 A 5 P	4 A 4 P 12 P 1 A A 11 P 4 P 3 P 4 P 3 A A 6 P	.14 .38 .13 .01 .05 .05 .15	
$\frac{30}{31}$	88 90	2 P 2 P	62 60	5 A 6 A	75 75	69 75	13.1 8.8	w sw	89 51	12 28	5 P	8 P	.18	

^{*} Based on least time required to blow one mile.

MARIUS P. VAVOUDES, Observer

	JOIVINIAI	
JULY 1950	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.334, 15th Minimum 29.704, 29th Mean semi-daily 30.001 Range 0.630	30.27 29.59 29.96 0.68	30.50, 1892 29.27, 1932 0.97, 1892 0.47, 1938
Air Temperature, in Degrees F.		
Highest 91, 18th Lowest 46, 15th, 21st Mean 70.6 Range 45 Highest mean daily 79.5, 18th Lowest mean daily 59.5, 20th Mean maximum 82.7 Mean minimum 58.8 Greatest daily range 37, 2nd Least daily range 10, 6th Degree-days of heating load for month. Degree-days cumulative for season.	93.9 46.4 70.8 47.7	104, 1911 40, 1890, 1898 66.3, 1891, 76.0. 1949
Precipitation, in inches		
Precipitation	4.10	14.51, 1897; 0.70, 1929
Snow. Maximum precip. in 24 hrs 1.23, 18th Number of days with .01 or more11	11	20, 1915; 4, 1924
Wind, in miles		
Total movement2348Greatest daily movement186, 14thLeast daily movement25, 8thMean hourly velocity3.2Maximum velocity28, 18th, 31st	34.22 4.6 28.7	5097, 1909; 1109, 1894 44, 1936
Wind, direction		
Prevailing direction WSW North, days 4 Northeast, days 0 East, days 0 Southeast, days 1 South, days 10 Southwest, days 4 West, days 6 Northwest, days 6	SW.	
Weather		
Mean relative humidity, percent 70 Mean cloudiness, percent 47.7 Number of clear days 14 Number of fair days 13 Number of cloudy days 4 Number of hours bright sunshine 293.2 Percent of possible hours of bright sunshine 63.2 Thunder and lightning 4th, 18th, 31st Last Snow. Last Frost 5	68.4 50.9 10 14 7 268	70, 1943; 31, 1924 22, 1923; 0, 1915 24, 1909, 1921; 4, 1946 18, 1889; 0, 1910 371, 1910; 180, 1931

The month was quite normal with regard to temperature and other conditions except precipitation. The mean figures over the period of observation indicate a 4.10 inches precipitation expected for the month with but 2.83 inches recorded; of which 1.23 inches fell on the 18th. While this storm was appreciated, it did not offset the development of drouthy conditions which slowed the growth of tobacco and made it unnecessary to mow lawns which were not sprinkled.

Even though a mean daily temperature of 59.5 degrees was recorded on the 20th, no heating load is considered in this locality to accumulate during July and August under conditions which are near the mean values.

Meteorological Series

Bulletin No. 740

August 1950

Meteorological Observations

FOR

AUGUST 1950

H. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N.

Longitude, 72° 31′ 48″ W.

Height of barometer above ground, 36 ft. Above sea level, 253.5 ft.

Height of wind instruments, 67 ft. Time used, E. S. T.

		Te	mpera	iture		ty	s		Wind		I	Precipitation			
	Maximum		Maximum Minimum			Humidi	of Hour nshine	b0	Į.	- S *					
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Вецап	Ended	Water	Snow	
1 2 3 4 5	77 70 72 80 79	12 N 2 P 4 P 2 P 3 P	66 58 61 60 53	4 A 5 A 5 A 5 A 6 A	71.5 64 66.5 70 66	88 76 76 67 70	0.6 4.2 7.4 11.4 10.6	N N S NW NW	62 70 37 86 70	11 7 6 13 18	4 A	12 N	.28		
6 7 8 9	75 76 84 88 85	2 P 4 P 5 P 3 P 4 P	59 52 46 52 67	6 A 5 A 5 A 6 A 5 A	67 64 65 70 76	67 66 67 67 71	9.6 12.1 12.8 11.0 8.1	NE NE N W	124 68 34 56 77	13 9 8 13 8	5 A 9 P	6 A 12 P	.01		
10 11 12 13 14	81 78 81 79	12 N 4 P 3 P 5 P	63 50 50 53	5 A 6 A 6 A 4 A	72 64 65.5 66	68 58 56 63	7.1 12.7 12.6 9.8	NW N N N	79 84 74 19	15 8 13 3) r	12 F	.02		
15 16 17 18	83 84 84 85	4 P 3 P 2 P 3 P	50 59 57 53	5 A 5 A 6 A 4 A	66.5 71.5 70.5 69	60 75 83 86	9.7 9.5 6.6	NW S S S	29 119 76 104	7 13 9 13	5 P	11 Р	.16		
19 20 21 22 23	80 73 76 77 80 84	12 N 0 A 5 P 2 P 4 P 4 P	67 62 52 46 46 58	5 A 11 P 6 A 6 A 7 A 5 A	73.5 67.5 64 61.5 63 71	88 100 69 72 78 82	3.3 0 11.9 10.6 10.5 9.3	S SW NW SW S	85 82 117 28 70 54	8 10 17 5 8 5 7	2 P 9 P	12 P 12 P	1.15 .01		
24 25 26 27 28	83 86 87 84	3 P 3 P 3 P 3 P	58 61 62 61	5 A 5 A 1 A 3 A	70.5 73.5 74.5 72.5	80 83 79 80	10.3 6.1 8.8 9.4	SW SW S	69 58 86 153	10 11 13	7 P	9 P	.13		
29 30 31	85 78 71	2 P 3 P 6 P	70 63 62	4 A 6 A 5 A	77.5 70.5 66.5	91 79 95	2.9 8.7 0	S NE SE	109 58 31	12 10 5	4 P 12 P	5 P 6 A	.01		

^{*} Based on least time required to blow one mile.

MARIUS P. VAVOUDES, Observer

MONTAL	501-11-1	
AUGUST 1950	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.240, 24th Minimum 29.747, 20th Mean semi-daily 30.022 Range 0.493	30.32 29.61 30.00 0.71	30.50, 1934 28.87, 1930
Air Temperature, in Degrees F.		
Highest 88, 9th Lowest 46, 8th, 22nd, 23rd Mean 68,8 Range 42 Highest mean daily 74, 28th Lowest mean daily 62, 21st Mean maximum 80,2 Mean minimum 57,3 Greatest daily range 38, 8th Least daily range 11, 1st, 3rd, 20th Degree-days of heating load for month Degree-days cumulative for season	91.6 43.4 68.6 48.1	100, 1918, 1948 34, 1940 62.4, 1903
Precipitation, in inches		
Precipitation	4.08	8.40, 1948; .31, 1894
Snow. Maximum precip. in 24 hrs 1.15, 19th Number of days with .01 or more 9	11	16, 1892, 1933; 4, 1899
Wind, in miles		
Total movement.2268Greatest daily movement153, 28thLeast daily movement19, 14thMean hourly velocity3.1Maximum velocity18, 5th	3127 4.2 22.7	4,271, 1910; 1,920, 1894 40, 1941
Wind, direction		
Prevailing direction WSW North, days 5 Northeast, days 3 East, days 0 Southeast, days 1 South, days 9 Southwest, days 5 West, days 2 Northwest, days 6	SW	
Weather		
Mean relative humidity, percent. 75.5 Mean cloudiness, percent. 55.9 Number of clear days 12 Number of fair days 13 Number of cloudy days 6 Number of hours bright sunshine 2593 Percent of possible hours of bright sunshine 58.2 Thunder and lightning 5th 10th 16th 18th 28th 29th	70.2 49.7 13 237 55.2	67, 1901, 1946; 27, 1923 23, 1941; 0, 1915, 1929 25, 1912; 3, 1939 18, 1901, 1928; 2, 1910, 1923 332, 1941; 152, 1915, 1929

The weather for the month was quite comfortable with the highest temperature, 88 degrees, reached on the 9th. The mean daily of 66.8 degrees, however, was accompanied by the mean relative humidity of 75.5% so that rather stuffy weather would have been experienced with much increase in temperature.

Only 2.94 inches of rain of the mean expected 4.08 inches was recorded and 1.15 inches of the total fell on the 19th. This broke the drouthy period to which farmers were giving serious consideration. The total wind movement of only 2268 miles against a mean of 3127 miles indicates that even normal wind could have made the effect of the drouth much more serious. Cloudiness above the mean value also undoubtedly contributed to making the drouthy conditions more bearable.

Meteorological Series

Bulletin No. 741

September 1950

Meteorological Observations

FOR

SEPTEMBER 1950

II. N. STAPLETON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

	Temperature						æ		Wind		I	Precipita	tion	
	Maximum		Mir	imum		Iumidi	f Hours nshine			8				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hour Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 22 22 23 24 25 26 27 28 29 29 20 20 20 20 20 20 20 20 20 20 20 20 20	84 81 82 75 68 75 80 84 84 84 72 61 63 69 65 74 70 68 58 76 65 53 60 53 60 65 77 70 68 77 70 68 70 70 70 70 70 70 70 70 70 70 70 70 70	12 P 1 P 2 P 3 P 4 P 5 P 4 P 2 P 3 P 12 N 2 P 3 P 12 N 2 P 3 P 12 N 2 P 4 P 2 P 4 P 2 P 4 P 2 P 3 P 12 N 2 P 3 P 4 P 2 P 3 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 4 P 4 P 4 P 4 P 5 P 4 P 4 P 5 P 4 P 4 P 5 P 4 P 5 P 5 P 4 P 5 P 5 P 6 P 6 P 7 P 8 P 8 P 8 P 8 P 8 P 8 P 8 P 8 P 8 P 8	70 64 56 52 48 46 49 53 51 50 48 49 32 44 29 32 46 50	2 A 8 A 1 5 A 6 A 6 A 6 A 6 A 5 A 6 A 5 A 6 A 6 A 6 A 6 A 6 A 6 A 6 A 6 A 6 A 6	77 72.5 64 63.5 860.5 63.5 68.5 68.5 57 56 58.5 57 52 59.5 54 62 52.5 52 52 52 52 54.5 62 62 62 62 62 63 64.5 65 65 66 67 62 62 63 64 65 66 66 66 67 68 67 68 68 67 68 68 68 68 68 68 68 68 68 68 68 68 68	94 85 98 82 73 67 73 77 77 77 92 90 74 76 85 84 72 72 72 74 88 86 87 86 86 86 87 88 88 88 88 88 88 88 88 88 88 88 88	2.5 9.1 0.0 8.3 11.1 11.9 10.8 7.9 0.3 0.0 7.9 9.9 6.7 4.7 11.5 11.7 8.1 0.0 10.4 8.3 10.7 5.9 6.4 6.8 5.9 6.6	SW N N N N N N N N N N N N N N N N N N N	78 51 53 97 149 45 224 115 48 163 189 39 42 110 218 85 54 49 108 87 50 249 232 92 20 35 35 38 24 24 24 25 26 27 28 28 28 28 28 28 28 28 28 28 28 28 28	12 12 7 20 18 6 4 3 7 8 17 14 8 4 13 20 13 7 8 5 9 9 4 25 18 18 18 18 18 18 18 18 18 18 18 18 18	12 A 4 A 0 A 8 A 0 A 1 A 6 A 5 A 0 A	5 P 12 P 1 A 8 P 2 P 7 A 5 A 12 P 4 A	.93 .37 .05 .05 .24 .31 .08 .12 .05 .07	

^{*} Based on least time required to blow one mile.

MARIUS P. VAVOUDES, Observer

. 101(11121		
SEPTEMBER 1950	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum	30.45 29.57 30.00	30.65, 1924 28.41, 1938
Mean semi-daily 30.044 Range 1.015	.88	1.99, 1938; .57, 1910
Air Temperature, in Degrees F.		
Ilighest .84, 1st Lowest .29, 25th Mean .58, 9 Range .55 Highest mean daily .72, 5, 1st Lowest mean daily .42, 25th Mean maximum .69, 4 Mean minimum .48, 8 Greatest daily range .35, 8th, 18th Least daily range .5, 11th Degree-days of heating load for month .209 Degree-days cumulative for season .209	87.7 33.2 61.7 54.5	97, 1895, 1929 24.5, 1914
Precipitation, in inches		
Precipitation	4.24	14.55, 1938; .52, 1914
Snow	10	16, 1933; 3, 1903
Wind, in miles		
Total movement2530Greatest daily movement249, 23rdLeast daily movement15, 8thMean hourly velocity3.5Maximum velocity25, 23rd	3271 4.5 25.9	4,686, 1896; 1,414, 1894 80, 1938
Wind, direction		
Prevailing direction. WNW North, days. 8 Northeast, days. 0 East, days. 0 Southeast, days. 1 South, days. 3 Southwest, days. 6 West, days. 4 Northwest, days. 8	WSW	
Weather		
Mean relative humidity, percent 80.5 Mean cloudiness, percent 55 Number of clear days 110 Number of fair days 77 Number of cloudy days 77 Number of hours bright sunshine 2087 Percent of possible hours of bright sunshine 56 Thunder and lightning 0	73.6 49.9 10 10 10 201 54.9	70, 1934; 23, 1947 19, 1932; 2, 1907, 1928 19, 1908; 3, 1889 20, 1934; 3, 1941, 1948 255, 1916; 106, 1934
First Frost	Sept. 21	Oct. 13, 1909; Aug. 22, 1894-95

With the last half of the month registering very cool temperatures, the mean for the month ran nearly three degrees below the average of 61.7 degrees. The low temperatures accompanied by less than average wind movement made the drouth more bearable. Of the 2.34 inches of precipitation recorded, 0.93 came in one storm on the first day of the month which spread the remainder very thin in a month, with an expected average 4.24 inches total rainfall.

Heating load for the month was 200 degree days, slightly over the average figure but hardly a measure of the seasonal load to be anticipated.

The first frost of the season occurred the night of the 24-25th

Meteorological Series

Bulletin No. 742

October 1950

Meteorological Observations

FOR

OCTOBER

EARLE COX

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

		Te	mper	ature		ty	o o		Wind			Precipitation			
	Ma	Maximum		mum Minimum		fumidi	f Hour nshine		,	20					
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max Miles per Hour*	Вецап	Ended	Water		
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 12 22 23 24 25 27 28 29 20 30 30 30 30 30 30 30 30 30 30 30 30 30	84 83 82 70 62 68 68 71 77 61 69 61 57 62 63 72 77 67 67 64 44 45 53 49 67 77 74 77	4 P 2 P 2 P 2 P 2 P 3 P 3 P 3 P 1 A 4 A 4 A 1 P 1 P 1 O A 3 P 4 P 3 P 2 P 2 P 2 P 2 P 3 P 4 P 3 P 4 P 3 P 4 P 2 P 2 P 4 P 4 P 5 P 5 P 6 P 6 P 7 P 8 P 8 P 8 P 8 P 8 P 8 P 8 P 8 P 8 P 8	48 53 53 54 50 32 33 33 36 44 47 59 50 39 42 43 37 58 59 42 43 37 44 47 47 47 47 47 47 47 47 47 47 47 47	6 A 6 A 6 A 6 A 6 A 6 A 6 A 6 A 6 A 7 A 8 A 7 A 7 A 7 A 6 A 7 A 7 A 6 A 7 A 7 A 6 A 7 A 7	668 688 608 647 49 52 57.5 64 556.5 58 60 53.5 48.5 61 67.5 63 48.5 50.5 43.5 50.5 43.5 50.5 43.5 55.5	777 855 799 700 755 744 833 799 777 774 889 726 880 744 886 886 887 738 886 897 887 997 838 887 997 838 887 997 888 888 888 888 888 888 888 88	8.6 9.9 6.8 10.5 11.1 9.4 8.1 6.3 6.3 0.0 8.1 1.5 4.1 8.5 9.9 10.2 9.3 6.3 9.3 10.3 5.1 9.5 0.0 0.0 5.1 9.5 0.0	NW SW SE W NW SW S NW S NW S NW S NW S N	37 53 108 178 73 41 66 84 79 96 72 246 269 37 44 108 55 120 221 57 65 118 52 96 221 43 39 71 99 90 55	7 6 13 12 13 13 11 17 7 12 9 8 8 28 20 4 7 13 6 10 25 7 18 20 5 7 18 20 18 20 18 20 18 20 20 20 20 20 20 20 20 20 20 20 20 20	5 A 2 A 7 A 7 A 5 A	12 P 4 P 1 P 8 A 8 A	1.01 .71 .09 .01 .03		

^{*} Based on least time required to blow one mile.

MARIUS P. VAVOUDES, Observer

OCTOBER 1950	Normal	Extremes
Barometer, in inches	TTOTMAT	
(Readings reduced to sea level)		
Maximum	30.53 29.42	30.68, 1929 29.00, 1926
Mean semi-daily	30.06	
Range	1.11	1.47, 1926; .76, 1899
Air Temperature, in Degrees F.		
Highest 84, 1st Lowest 22, 27th Mean 54.2 Range .62 Highest mean daily 68, 2nd, 3rd Lowest mean daily 36.5, 28th Mean maximum 65.4 Mean minimum 43.4 Greatest daily range 38, 6th Least daily range 2, 12th Degree-days of heating load for month 344	79.4 23.4 50.5 56.0	90.5, 1908 17, 1936 57.4, 1947; 43.2, 1890
Degree-days cumulative for season 553		
Precipitation, in inches		
Precipitation 1.87 Snow —	3.29	8.81, 1911; .01, 1924
Maximum precip. in 24 hrs 1.01, 10th Number of days with .01 or more 6	9	15, 1913; 1, 1897, 1924
Wind, in miles		
Total movement 2968 Greatest daily movement 269, 14th Least daily movement 37, 1st, 15th Mean hourly velocity 3,98	4074 5.4 29.5	5,467, 1910; 2,540, 1894 42, 1937
Maximum velocity	27,3	12, 1707
Wind, direction		
Prevailing direction WNW North, days. 4 Northeast, days 0 East, days. 1 Southeast, days 1 South, days 5 Southwest, days. 3 West, days 7 Northwest, days 10	W	
Weather		
Mean relative humidity, percent. 80.22 Mean cloudiness, percent 45 Number of clear days 14 Number of fair days 10 Number of cloudy days 7 Number of hours bright sunshine. 200.8 Percent of possible hours of bright sunshine. 58.5 Thunder and lightning.	69.0 48 10 10 11 175.0 51.3	67, 1932; 18, 1924 21, 1938; 1, 1911 17, 1924; 3, 1938 19, 1896; 3, 1924 232, 1923, 1938; 91, 1913
First Frost		



Meteorological Series

Bulletin No. 743

November 1950

Meteorological Observations

FOR

NOVEMBER

1950

EARLE COX

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T

		Te	mpera	ature		r.	ø		Wind		Precipitation			
	Ma	ximum	Mir	nimum		Jumidi	f Hour nshine		t	sa,		!		
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Ведап	Ended	Water	Snow
1 1 2 2 3 3 4 5 5 6 6 7 1 8 9 9 10 11 12 13 14 15 16 17 18 18 19 20 21 22 22 23 24 5 26 27 28 29 1 2 3 3 3 1	72 82 48 47 64 65 65 65 65 65 65 44 47 63 38 37 46 40 40 40 37	3 P 2 P 1 P 8 A 8 A 8 A 9 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P 1	42 444 43 442 37 31 32 50 54 36 29 21 22 21 42 38 37 22 43 37 42 37 43 42 43 37 43 42 43 44 42 43 44 44 45 46 47 47 47 47 47 47 47 47 47 47 47 47 47	1 A 7 A 5 A 1 A 1 A 1 A 1 A 1 A 1 A 1 A 1 A 1	57 63 45.5 44.5 50.5 44.5 57.5 59.5 44.3 37.5 32.5 33.5 36.5 33.5 40 40 53 33.5 46 40 53 33.5 47 40 53 33.5 33.5 33.5 33.5 33.5 33.5 33.5	79 76 89 91 70 69 78 82 87 77 28 86 85 77 81 81 81 88 89 75 81 88 89 91 88 88 88 88 88 88 88 88 88 88 88 88 88	9.1 8.0 0.0 0.0 5.0 8.6 9.0 4.2 1.2 9.0 9.0 9.0 9.0 8.0 5.0 6.5 0.0 8.3 3.4 0.0 0.0 2.3 6.5 0.4 4.0 0.2	WWNNNNWSWNWNWSNNWNWSNNWNWSNNWNWNWNWNWNW	13 90 135 119 235 102 94 140 151 225 40 22 22 50 68 145 174 208 330 343 30 343 30 343 141 550 252 78 91 140 177	3 20 15 8 25 9 9 13 13 28 5 2 2 2 11 12 13 13 14 10 32 32 28 6 6 20 48 11 17 17 18 18 18 18 18 18 18 18 18 18 18 18 18	10 P O A 4 P O A 2 P O A 2 A	12 P 1 A 12 P 1 A 12 P 4 A	.28 .79 .80 .05 .07 1.35 .44 2.28 .39 .03 .12 T	0.5 T

^{*} Based on least time required to blow one mile.

MARIUS P. VAVOUDES, Observer

NOVEMBER 1950	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.447, 14th Minimum 29.352, 25th Mean semi-daily 29.978 Range 1.095	30.59 29.30 30.05 1.29	30.87, 1932 28.73, 1904 1.84, 1904; 89, 1943
Air Temperature, in Degrees F.		
Highest 82, 2nd Lowest 19, 23rd Mean 42.8 Range .63 Highest mean daily 63, 2nd Lowest mean daily 26, 23rd Mean maximum 51.5 Mean minimum 34.2 Greatest daily range 38, 2nd Least daily range 5, 3rd-4th	66.0 12.9 38.9 53.0	82 ,1950 4, 1938 44.1, 1931; 33.6, 1901
Degree-days of heating load for month 667 Degree-days cumulative for season 1219.5	783	
Precipitation, in inches		
Precipitation	3.41 2.34	8.64, 1927; 0.63, 1917 13.50, 1938
Snow. 0.5 Maximum precip. in 24 hrs. 2.28, 25th Number of days with .01 or more 11	9	22, 1921; 2, 1904
Wind, in miles		
Total movement 4623 Greatest daily movement 550, 25th Least daily movement 13, 1st Mean hourly velocity 0.3 Maximum velocity. 48, 25th	4531 6.3 30.3	5,978, 1906; 2,589, 1889 48, 1948, 1950
Wind, direction		
Prevailing direction W North, days 3 Northeast, days 1 East, days 1 Southeast, days 0 South, days 10 Southwest, days 3 West, days 2 Northwest, days 10	WNW	
Weather		
Mean relative humidity, percent. 82.5 Mean cloudiness, percent 56 Number of clear days 11 Number of fair days 8 Number of cloudy days 11 Number of hours bright sunshine. 138.4 hrs Percent of possible hours of bright sunshine. 47.2 Thunder and lightning 0	70.6 55.1 6 9 15 121 41.3	78, 1947; 34, 1917 15, 1903; 0, 1948 16, 1912; 4, 1889, 1930, '32 '35 24, 1927; 9, 1905, 1917, 1949 182, 1903; 44, 1948
First Frost	Nov. 6	Oct. 10, 1925; Dec. 13, 1941

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1949.

SUMMARY

The outstanding feature of the weather during the month was the high wind and rainfall which came on the 25th. The daily movement of wind was 550 miles, which is the highest that has occurred since February 15, 1940. The recorded velocity of 48 miles per hour was a record for the month of November. A total of 2.28 inches of rainfall came with the wind storm.

A new record for air temperature in November occurred on November 2 when the air temperature was at 82° F. In general, warm weather prevailed throughout the month giving a mean daily of 42.8° F. as compared to a normal of 38.9° F. The degree days of 667 was below the normal of 783.

The rain which came with the storm on the 25th resulted in a total precipitation for the month of 6.60 inches, which was considerably above the normal of 3.41. The total precipitation to the end of November was 37.23 inches which is somewhat below the normal of 40.31 inches.

The mean relative humidity of 82.5 per cent was far above the normal of 70.6.

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 744

December 1950

Meteorological Observations

FOR

DECEMBER

1950

EARLE COX

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

		Te	emper	ature		2	s		Wind		1	Precipita	ation	
		ximum		nımum		Mean Relative Humidity	Number of Hours Bright Sunsume	ling	nent	Miles our*				
Day	Degrees	Time	Degrees	7 ime	Mean	Mean Relati	Numb Bright	Prevailing Direction	Total Movement	Max Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22	38 36 56 64 43 43 42 50 45 41 39 40 37 33 33 35 37 32 26 27 34	10 A 2 P 3 P 12 P 2 P 2 P 2 P 1 A P 12 P 12 P 12 P	26 21 26 51 30 24 42 31 25 34 30 24 28 42 31 25 31 25 31 25 31 30 31 30 31 30 31 30 31 30 31 31 31 31 31 31 31 31	12 P 6 A A 6 A O A 12 P 7 A A 3 A A 12 P 7 A O A 12 P 7 A 12 P 6 A 12 P 6 A 8 A 3 A	32 28.5 41 57.5 36.5 33.5 35 46 38 33 32.5 34.5 29.5 26.5 33 19.5 16	83 82 90 90 74 74 80 97 97 90 89 74 75 72 86 79 77 77 77 77 83 83 83 86	1.7 4.4 5.7 0.0 5.0 0.0 6.6 1.5 1.2 7.2 4.9 0.0 0.0 0.0 7.3	NW S S S N N NE SE SW N N NW S N N N N N N N N N N N N N N	98 35 60 230 410 196 153 149 52 67 129 129 129 129 129 129 148 2 31 224 50 9	14 8 9 18 28 17 28 9 7 14 14 13 5 4 3 3 12 2 2 0 15 11 11 11 12 12 13 14 14 15 15 16 16 17 17 18 18 18 18 18 18 18 18 18 18 18 18 18	12 N 2 A 2 A 11 A 0 A	12 P 12 P 6 A 12 P 8 P	1.81 .65 .39	T 2.0 1.0
23 24 25	40 39 22	3 P 2 P 1 P	24 30 14	8 A 7 A 12 P	32 34.5 18	80 87 68	0.0 0.0 6.0	s s NW	43 90 297	12 32 28	2 Р	4 P	.06	1.0
26 27 28	15 16 30	3 P 3 P 3 P	12 - 3 - 3	6 A 7 A 6 A	13.5 6.5 13.5	77 73 73	1.0 9.1 9.1	W W S	7112 72 47	11 13 8	12 N	3 P	.06	2.0
29 30 31	33 34 37	5 P 2 P 2 P	9 22 26	0 A 8 A 0 A	21 28 31.5	95 99 97	0.0 3.5 4.1	N N N	30 0 25	11 0 20	8 A 2 A	12 P 7 A		0.5 2.0

^{*} Based on least time required to blow one mile.

MARIUS P. VAVOUDES, Observer

DECEMBER 1950	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.569, 7th Minimum 29.384, 4th Mean semi-daily 30.027 Range 1.185	30.66 29.27 30.06 1.38	31.104, 1949 28.85, 1915 1.78, 1895, 1.01, 1892
Air Temperature, in Degrees F.		
Highest 64, 4th Lowest -3, 27th, 28th Mean 29,1 Range 67 Highest mean daily 57,5, 4th Lowest mean daily 6,5, 27th Mean maximum 37.0 Mean minimum 21.8 Greatest daily range 33, 28th Least daily range 3, 26th Degree-days of heating load for month1114.0 Degree-days cumulative for season 2333.5	54.3 1.09 27.5 56.2	65.5, 1908 —22.5, 1917 36.9, 1891; 17.1, 1917
Precipitation, in inches		
Precipitation4.64	3.39	7.77, 1901; .58, 1943 26.50, 1902, T, 1891, 1943
Snow 8.5 Maximum precip. in 24 hrs 1.81, 4th Number of days with .01 or more 9	8.50 10	17, 1902; 4, 1892, 1943
Wind, in miles		
Total movement3289Greatest daily movement410, 5thLeast daily movement0, 30thMean hourly velocity4.4Maximum velocity32, 24th	4710 6.3 31.4	6,694, 1925; 3,239, 1918 66, 1949
Wind, direction		
Prevailing direction WNW North, days 11 Northeast, days 1 East, days 0 Southeast, days 1 South, days 8 Southwest, days 2 West, days 4 Northwest, days 4 Northwest, days 4	WNW	
Weather		
Mean relative humidity, percent. 83.8 Mean cloudiness, percent	69.4 54.9 8 9 14 128 45.2	71, 1929; 39, 1919 15, 1890, 0.1948 16, '09; 4, '89, '30, '31, '36, '38 25, 1948; 7, '09, '23 172, 1896; 27, 1948
First Frost	Nov. 6	

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1949.

ANNUAL SUMMARY

ANNUAL	Normal	Extremes
Barometer, in inches (Readings reduced to sea level) Maximum 30.77, Feb. 8th Minimum 29.18, Mar. 23rd Mean semi-daily 20.00 Range 11.59	30.81 28.95 30.01 1.85	31.10, 194 28.41, 193 2.47, 1938; 1.38, 193
Air Temperature, in degrees F.		
lighest	95.7 12.2 47.4 107.8	104, 191 —26, 190 56.9, 1949; 44, 190
owest mean daily 0, Feb. 20th dean maximum 59.5 dean minimum 38.3 Greatest daily range 41, June 2nd least daily range 2 Oct. 12th Degree-days cumulative for season 6404	57.8 36.5	
Precipitation, in inches		
Precipitation 42.17 Snow 53.1 Maximum precip, in 24 hrs. 2.28, Nov. 25th	43.70 47.78	59.00, 1938; 30.68, 1908 89.00, 1893; 24.50, 1919
Number of days with .01 or more 132	124	146, 1945; 96, 1924
Wind, in miles		
Total movement	52,223	63,571, 1908; 36,257, 1894 705, 1909
east daily movement 0, Dec. 30th dean hourly velocity 5.0 daximum velocity 48, Nov. 25th, May 27th	5.8 39.5	80, 1938
Wind, direction		
Prevailing direction WNW North, days 83 Northeast, days 18 East, days 7 Southeast, days 17 South, days 70 Southwest, days 39 West, days 47 Northwest, days 84	W.	
Weather		
Mean relative humidity, percent 74.7 Mean cloudiness, percent 55.2 Number of clear days 128 Number of fair days 125 Number of cloudy days 112 Number of hours bright sunshine 2273.0 Percent of possible hours of	67.6 51.7 116 123 126 2,353	60, '98, '01, '02; 41, '08, '24 217, 1941; 59, 1927 182, 1912; 64, 1936 179, '01, '02; 71, 1910, 1941 3038, 1941; 1864, 1902
bright sunshine 49.5 Last snow April 29th First snow Nov. 21st Last frost Apr. 30th First frost Sept. 25th	52.8 April 15 Nov. 6 May 14 Sept. 21	Mar. 14 '10; May 11, '07, '45 Oct. 10, '25; Dec. 13, '41 Apr. 14, '43; June 8, '32 Aug. 22. '94, '95; Oct. 13, '09

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 745

January 1951

Meteorological Observations

FOR

JANUARY

1951

EARLE COX

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

		Te	mpera	ture		ty.	9		Wind		I	Precipita	tion	
	Max	ximum	Mir	imum		Tumidi	of Hour nshine		t	es.				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5	36 35 44 55 38	2 P 3 P 8 P 12 N 2 P 2 P	13 15 35 34 24	7 A 5 A 0 A 12 P 8 A	24.5 25 39.5 44.5 31	88 90 92 82 56	7.9 3.5 0.0 0.0 0.0	SE SW NW	55 35 163 211 119	7 6 13 28 13	1 P	2 Р	.07	
6 7 8 9	44 38 19 30 38	2 P 0 A 6 X 2 P 3 P	26 19 10 4 10	0 A 12 P 12 P 7 A	35 28.5 14.5 13 24	78 88 83 81 81	0.0 0.0 6.0 8.0 7.1	S N NW S	91 168 231 85 72	15 11 22 10 6	10 A 0 A	12 P 5 A	.50 .03	7.0 0.5
11 12 13	31 38 39	5 P 3 P 3 P	11 30 23	6 A 0 A 12 P	21 34 31	91 70 69	0.0 3.0 7.9	N NW N	16 194 188	1 25 14	8 P	10 P	.02	0.3
14 15 16 17	32 43 33 40	8 P 2 P 1 P 4 P	7 28 26 17	7 A 0 A 12 P 4 A	19.5 35.5 29.5 28.5	85 86 68 71	1.0 3.0 3.0 0.5	NW NW S	14 119 338 120	28 25 12	5 P 0 A	12 P 1 P	.38	4.0
18 19 20 21	43 53 50 46	2 P 4 P 3 P 10 A	30 26 28 17	7 A 6 A 7 A 12 P	36.5 39.5 39 31.5	77 87 94 79	3.0 6.9 0.0 0.0	S SW SW NW	29 20 18 254	4 8 8 32	7 A	8 A	Т	
22 23 24 25 26	23 36 45 39 31	3 P 12 P 11 A 3 P 4 P	11 9 33 27 22	8 A 7 A 12 P 8 A 7 A	17 22.5 39 33 26.5	56 76 91 89 71	8.5 0.0 0.0 2.0 5.5	N SW N	231 34 147 51 165	14 13 15 9 18	6 A	3 P	.91	
26 27 28 29 30	28 26 29 22	3 P 3 P 2 P 10 A	20 18 19	8 A 6 A 12 P 12 P	24 22 24 13	74 76 91 72	0.0 2.5 1.5 5 1	SW SW NW NW	165 138 36 89 161	8 6 11 25	4 P 0 A	12 P 11 A	.23 .16	1.0
31	10	12 P	-1	7 A	4.5	74	0.0	N	163	12	9 A	12 P	.46	5.5

^{*} Based on least time required to blow one mile.

MARIUS P. VAVOUDES, Observer

MONTHLI	Somm	IAIXI
JANUARY, 1951	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.728, 31st Minimum 28,996, 15th Mean semi-daily 30.064 Range 1.732	30,70 29,20 30,07 1,50	31.00. 1927 28.55, 1913 2.18, 1913; 0.97, 1896
Air Temperature, in Degrees F.		
Highest 55, 4th Lowest -4, 9th Mean 27,4 Range 59 Highest mean daily 44,5, 4th Lowest mean daily 4,5, 31st Mean maximum 35,9 Mean minimum 18,9	51.1 -7.1 24.2 58.2	66.0, 1932 —26.0, 1904 34.2, 1913; 13.9, 1918
Greatest daily range	1265	
Precipitation, in inches		
Precipitation	3.61 13.32	7.15, 1898; 1.07, 1896 33,1898, 1923; 1.50, '08
Maximum precip. in 24 hrs 0.91, 24th Number of days with .01 or more 10	11	17, '50; '20; 4, 1901
Wind, in miles		
Total movement	5055 6.8 32.7	7770, 1908; 2896, 1895 47, 1938
Wind, direction		
Prevailing direction WNW North, days 8 Northeast, days 0 East, days 0 Southeast, days 1 South, days 5 Southwest, days 5 West, days 2 Northwest, days 10	WNW	
Weather		
Mean relative humidity, percent. 79.5 Mean cloudiness, percent 51.0 Number of clear days 6 Number of fair days 8 Number of cloudy days 17 Number of hours bright sunshine 85.9 Percent of possible hours of bright sunshine 29.2 Thunder and lightning	70.1 55 9 13 137 46.7	78, 1932; 37, 1908 20, 1944; 1, 1949 18, 1890; 1926; 3, 1916, 1944 22, 1923, '31, '49; 4, 1920, '39 214, 1920; 50.4, 1949
First Frost.		

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1950.

SUMMARY

The mean daily temperatures the forepart of January were above normal. This, together with above normal mean daily temperatures during November and December, resulted in an accumulated degree days of 3,013 to January 22, which is the mean mid-point of the heating season. This is about 10 per cent below the normal of 3,352. The mean daily for the month of 27.4° F. was more than 3° F. above the normal of 24.2° F.

The precipitation during the month of 3.28 inches was below the normal of 3.61 inches.

A MEAN RELATIVE HUMIDITY of 79.5 per cent was considerably above the normal of 70.1 per cent and added to the ventilation problems in poultry houses and dairy barns.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 746

February 1951

Meteorological Observations

FOR

FEBRUARY

1951

EARLE COX

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

		Te	mpera	iture		,	10		Wind		I	Precipita	tion	
Day	Degrees	Lim e mumin	Degrees Min	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 22 23 24 25 26 27 28 28 29 20 30 30 30 30 30 30 30 30 30 30 30 30 30	48 46 20 42 41 53 8 14 22 35 42 41 38 40 39 45 36 36 36 36 36 36 36	9 P 0 A 8 P 4 P 1 P 2 P 3 P 4 P 4 P 4 P 4 P 2 P 3 P 4 P 2 P 3 P 4 P 4 P 2 P 3 P 4 P 1 2 P 3 P 4 P 1 P 2 P 3 P 4 P 1 P 2 P 3 P 4 P 2 P 2 P 2 P 2 P 2 P 3 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 3 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2	10 17 7 16 14 15 26 6 2 —6 12 26 33 33 20 15 11 20 33 34 34 34 32 26 29 34 32	O A 12 P 7 A 2 A 7 A 12 P 8 P 7 A O A 3 A 12 P 8 A 7 A 7 A 12 P 8 A 7 A 7 A 12 P 7 A 12 P 7 A 12 P 7 A 12 P 7 A 12 P 7 A 12 P 7 A 12 P 7 A 12 P	29 31.5 13.5 27.5 28 39.5 7 8 8 23.5 24.5 28.5 24.5 35.5 37.3 36.5 37.3 36.5 37.3 31.5 41.5 41.5 41.5		0.0 5.5 0.3 7.5 7.1 7.0 0.0 6.9 8.9 9.0 5.0 0.1 0.0 8.9 9.0 0.0 0.0 0.0 0.0 0.0 0.0 0	S NW N NW NW S S NW NW NW S S NW NW NW NW NW NW NW NW NW NW N NW NW N	76 330 60 122 2 31 241 416 143 24 62 462 462 179 148 25 179 148 25 31 34 79 12 298 432 471 298 187	20 32 11 18 8 20 32 40 11 7 8 6 10 32 332 332 28 18	11 P O A 2 P 4 P O A 7 P 7 A 0 A 11 A 2 P O A	12 P 12 P 6 A 12 P 3 P 12 P 9 A 8 P 12 P 5 A 2 P 3 P	.02 .03 .03 .04 .05 .05 .03 .03 .03	.5 1.5 4.0

^{*} Based on least time required to blow one mile.

MONTHEL	SUMM	AKI
FEBRUARY, 1951	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30,850, 10th Minimum 29,200, 7th Mean semi-daily 30,161	30.63 29.24 30.04	31.05, 1920 28.56, 1895
Range 1.650	1.38	1.89, '00, '08; .88, '13, '31
Air Temperature, in Degrees F.		
Highest 56, 27th Lowest -6, 10th Mean 29,6 Range .62 Highest mean daily 45, 27th	50.5 7.4 23.7 58	65.0, 1930 —23, 1943 32.6, 1890; 11.6, 1934
Lowest mean daily 7, 8th Mean maximum 38.3 Mean minimum 21.0 Greatest daily range 38, 1st Least daily range 2, 8th	32.8 14.5	
Least daily range 2, 8th Degree-days of heating load for month 990 Degree-days cumulative for season	1166.7	
Precipitation, in inches		
Precipitation 4.61 Snow 6.0 Maximum precip. in 24 hrs 1.33, 7th	3.19 13.99	8.12, 1900; 0.62, 1901 48.75, 1893; 0.50, 1937
Number of days with .01 or more	10	15, '93, 1920; 4, 1901
Wind, in miles		
Total movement .4088 Greatest daily movement .471, 24th Least daily movement .2, 5th Mean hourly velocity .6.1 Maximum velocity .32, 22nd, 23rd, 24th	7.1 31.7	6445, 1896; 3402, 1949 50, 1946
Wind, direction	31.7	30, 1710
Prevailing direction. NNW North, days. 13 Northeast, days. 1 East, days. 0 Southeast, days. 0 South, days. 5 Southwest, days. 2 West, days. 1 Northwest, days. 6	WNW	
Weather		
Mean relative humidity, percent. 79.0 Mean cloudiness, percent 55.4 Number of clear days 8 Number of fair days 5 Number of cloudy days 15 Number of hours bright sunshine. 99.0 Percent of possible hours of bright sunshine 33.5 Thunder and lightning 33.5	66.6 50.9 10 8 10 158	66, '90, '27; 31, 1905 19, 1941; 2, 1927 16, 1920; 2, 1936 16, 1894; 2, 1920 221, 1924; 110, '27, '38
First Frost.		

REMARKS

The weather for the month was generally warm except for a cold period on the 8th, 9th, and 10th. The last 11 days of the month were above 30 degrees mean daily and 8 of the 11 were above 35 degrees mean daily. The mean daily for the month was 29.6 degrees. The warm weather during the month gave a total of 990 degree days for the month, as compared to a normal of 1166.7. The cumulative degree days to the end of the month was 4428 which is only slightly above that of 1950 which was 4383. Only 6 inches of snow fell during the month, as compared to a normal of 13.99, but precipitation for the month was 4.61 inches, as compared to a normal of 3.19. Considerable cloudiness occurred throughout the month and the hours of bright sunshine were 99 which was far below the normal of 158. The mean relative humidity of 79 per cent was also far above the normal of 66.6.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 747

March 1951

Meteorological Observations

FOR

MARCH 1951

EARLE F. COX

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

		Te	mpera	ture		ty	8		Wind		I	Precipita	tion	
	Max	imum	Mir	imum		fumidí	f Hour nshine			8				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2	26 39	3 P 2 P	22 22	12 P 0 A	24 30.5	82 79	0.0 5.0	E NW	136 119	20 20	8 A	12 Р	.18	2.0
3 4 5	34 34	1 P 8 P	23	8 A 6 A	28.5 30.5	70 94	5.1 0.0	E N	138 49	17 8	7 P 0 A	12 P 5 P	.04 .50	.5 3.0
5 6 7 8 9	41 50 51 42 34	5 P 4 P 1 P 4 P 12 N	28 22 34 31 23	5 A 7 A 7 A 6 A 12 P	34.5 36 42.5 36.5 28.5	84 72 88 65 69	9.7 7.8 0.6 0.0 4.5	N SW W NW	88 56 95 158 149	13 9 10 17 22	2 P	4 P	.03	
10 11 12 13	36 45 54 42	2 P 3 P 4 P 2 P	20 25 35 35	7 A 3 A 6 A	28 35 44.5 38.5	58 73 69 79	10.2 5.0 9.5 0.0	N N N	371 382 217 131	32 32 17 11				
14 15 16	42 39 48	4 P 2 P 1 P	37 36 32 30	0 a 0 a 6 a	39.5 37.5 40 37	89 89 82 82	0.0 0.0 0.0 0.0	E S W	179 99 95 107	17 10 13 14	8 a 0 a	12 P 4 A	.55 .02	
17 18 19 20	44 47 45 43	4 P 4 P 4 P 3 P	29 32 29	4 A 6 A 5 A 12 P 7 A	38 38.5 36 38	67 79 79	0.0 0.0 5.0	N SE N	143 135 243 129	18 15 25 18	11 P 0 A	12 P 9 A	.02 .98	
21 22 23 24 25	50 34 48 56	3 P 5 P 4 P 2 P	26 21 19 36	12 P 3 A 6 A	27.5 33.5 46	67 87 74 83	6.2 7.0 2.5 2.0	S N S W	260 115 91	25 15 22	2 A 5 A 0 A	11 A 12 P 12 N	.17 .17 .25	1.5 1.0
26 27 28	41 33 50 65	2 P 4 P 5 P 4 P	29 23 17 34	6 A 6 A 6 A 7 A	35 28 33.5 49.5	59 57 63 69	8.5 10 9 7	NW NW S	311 274 83 97	40 28 9 9				
29 30 31	46 51 55	8 P 2 P 2 P	41 41 40	7 A 3 A 12 P	43.5 46 47.5	96 89 81	0.0 0.0 4.5	NW SE NW	31 118 219	9 20 20	6 A 0 A 0 A	12 P 12 P 6 A	.45 1.62 .15	

^{*} Based on least time required to blow one mile.

MARIUS P. VAVOUDES, Observer

MARCH, 1951	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum .30.491, 3rd Minimum .29.518, 24th Mean semi-daily .30.076	30.57 29.24 30.04	31.05, 1943 28.47, 1914
Range	1.34	2.10, 1914; 0.85, 1915
Air Temperature, in Degrees F.		
Highest 65, 28th Lowest 17, 27th Mean 36,5 Range 48 Higher and dilly 16, 20th 31cm	63.8 6.1 34.4 57.6	85, 1929 —9, 1950
Highest mean daily .46, 30th, 31st Lowest mean daily .24, 1st Mean maximum .41.1 Mean minimum .29.0 Greatest daily range .33, 27th	43.2 25.4	
Least daily range	949 5876	
Precipitation, in inches		
Precipitation	3.70 7.47	7.89, 1942; 0.12, 1915 27, 1899; 0, 1921
Maximum precip. in 24 hrs 1.62, 30th Number of days with .01 or more	11	17, 1890, 1913, 1936; 3, 1915
Wind, in miles		
Total movement	5944	8182, 1896; 3006, 1905
Mean hourly velocity	7.7	48, 1932, 1939, 1941, 1942
Wind, direction		
Prevailing direction NNW North, days 10 Northeast, days 0 East, days 4 Southeast, days 2 South, days 4 Southwest, days 2 West, days 3 Northwest, days 6	WNW	
Weather		
Mean relative humidity, percent. 76.6 Mean cloudiness, percent 58.9 Number of clear days 6 Number of fair days 10 Number of cloudy days 15 Number of hours bright sunshine. 119.1 Percent of possible hours of bright sunshine. 32.2 Thunder 16th	64.2 51.8 11 10 10 199 53.6	68, 1901; 27, 1915 22, 1924; 3, 1901 17, 1900; 1, 1943 21, 1901; 1, 1915 292, 1924; 93, 1901
First Frost		

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1950.

REMARKS

The month of March was generally mild with a mean daily of 36.5 degrees, as compared to a normal of 34.4 degrees. One abrupt temperature change occurred during the month when the temperature rose from a low of 17 degrees on one day to a high of 65 degrees on the following day. These two figures were the minimum and maximum for the month. The accumulated degree days for the season were 5321, as compared to 4917 in 1949 and 5468 in 1950. Rainfall of 1.62 inches on March 30 gave a precipitation for the month of 5.13 inches, as compared to a normal of 3.70 inches. As with previous months, the mean relative humidity contained far above normal with a reading of 76.6 per cent, as compared to a normal of 64.2 per cent.

MASSACHUSETTS

Meteorological Series

Bulletin No. 748

April 1951

Meteorological Observations

AGRICULTURAL EXPERIMENT STATION

FOR

APRIL

1951

EARLE F. COX

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

		Т	emper	ature		ķ			Wind	l	:	Precipita	ation
	Ма	ximum 1	-	nimum		Humidit	of Hours unshine	8.6	ıt	les *			
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 30 30 30 30 30 30 30 30 30 30 30 30	63 47 44 49 61 65 55 68 52 67 63 57 63 57 63 57 63 57 63 57 68 52 67 68 69 69 69 69 69 69 69 69 69 69 69 69 69	4 P 3 P 4 P 3 P 5 P 4 P 3 P 4 P 12 N 1 P 1 P 2 P 3 P 4 P 1 P 2 P 3 P 4 P 1 P 2 P 3 P 4 P 2 P 3 P 4 P 2 P 3 P 4 P 2 P 3 P 4 P 2 P 3 P 4 P 2 P 3 P 4 P 2 P 4 P 5 P 6 P 6 P 7 P 8 P 8 P 8 P 8 P 8 P 8 P 8 P 8	31 42 36 33 33 36 34 46 38 39 43 35 42 40 36 43 42 40 31 32 33 41 30 45 40 40 40 40 40 40 40 40 40 40 40 40 40	6 A 8 A 8 A 6 A 6 A 6 A 6 A 6 A 6 A 6 A	47 44.5 40 42 39.5 48.5 50.5 46.5 53.5 47.5 51 49.5 51.5 46.5 41 42 50.5 46 48 47 48.5 55.5 55.5 55.5 55.5 55.5	68 83 87 70 73 63 69 80 80 73 81 87 75 72 69 72 74 70 60 61 73 70 62 83 81 70 69 70 60 61 70 60 60 60 60 60 60 60 60 60 60 60 60 60	6.5 0.0 0.5 3.0 6.0 11.0 9.5 0.2 7.1 5.0 0.2 8.5 3.0 2.2 1.0 2.5 11.5 12.0 2.1 9.1 7.7 0.0 4.3 11.1 7.0 2.2 12.1	W W NW NW NW N N N NE SE S S SW SW SW SW SW NW S N N N S N N N S N N N S N N N S N N N S N N N S N N N N S N N N N S N	48 63 180 117 74 183 147 101 93 65 107 102 151 189 250 151 216 136 168 225 95 143 293 79 280 60 226 239	111 132 28 114 122 200 133 111 188 200 40 15 22 28 20 18 20 13 113 113 113 12 28 20 13 13 113 114 115 116 117 118 118 119 119 119 119 119 119 119 119	5 P 10 A 10 P 0 A 6 A 2 A	12 P 3 P 1 P 1 P 1 2 P 4 A 8 A 10 A 12 P 5 P 12 P 8 A	.12 1.64 .23 .21 .05 .37 .01 .01 .06 .07

^{*} Based on least time required to blow one mile.

MARIUS P. VAVOUDES, Observer

APRIL, 1951	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.395, 22nd Minimum 29.230, 15th Mean semi-daily 29.899	30.48 29.38 29.99	30.71, 1911 28.98, 1943
Range	1.11	1.52, 1930; 0.65, 1950
Air Temperature, in Degrees F.		
Highest 71, 29th Lowest 30, 5th, 24th Mean 48.4 Range .41 Highest mean daily 59, 29th Lowest mean daily 39.5, 5th Mean maximum 59.1 Mean minimum 37.9	79.4 22.0 45.7 57.4 56.8 34.8	90, 1941 8.5, 1923 52, 1921; 41, 1943
Greatest daily range	579 6455	
Precipitation, in inches		
Precipitation	3.35 2.17	6.89, 1929; .55, 1941 11, 1891; 0, 1910, 1934
Snow	11	18, 1909; 3,1892
Wind, in miles		
Total movement	7.5 31.4	8208, 1908; 3853, 1917 40, 1935, 1938, 1940, 1945
Wind, direction		
Prevailing direction W North, days 5 Northeast, days 2 East, days 0 Southeast, days 3 South, days 4 Southwest, days 5 West, days 4 Northwest, days 7	WNW	
Weather		
Mean relative humidity, percent. 72.6 Mean cloudiness, percent 50.0 Number of clear days 7 Number of fair days 10 Number of cloudy days 13 Number of hours bright sunshine. 169.7 Percent of possible hours of bright sunshine. 42.3 Thunder and Lightning 10th	61.6 51.8 11 9 10 220 54.7	75, 1901; 34, 1925, 1927 23, 1941; 1, 1949 18, 1915; 2, 1894, 1901 22, 1901; 1, 1941 329, 1941; 103, 1901
Last Snow First Frost	Apr. 15	Mar. 14, 1910; May 11, 1907.'45

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1950.

REMARKS

The mean daily temperature for the month of April was 48.4 degrees as compared to a normal of 45.7 degrees. There were 500 degree days for the month and the normal is 579. The cumulative degree days for the season of 5821 is far below the normal of 6455.

The precipitation for the month was about normal with a total of 3.63 inches. The normal for the month is 3.35 inches. No snow fell during the month, while normally a snowfall of 2.71

inches would occur.

As with previous months, the mean relative humidity continued far above normal. The reading was 72.6 per cent as compared to a normal of 61.6 per cent. Considerable cloudy and foggy weather occurred during the month. The hours of bright sunshine were only 169.7 as compared to a normal of 220. Foggy weather occurred nine days during the month when the reading was taken at 8 a.m.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 749

May 1951

Meteorological Observations

FOR

MAY

1951

EARLE F. COX

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

-		Τe	empera	ature		2	8		Wind		Precipitation			
Day	Ма	Time	Degrees K	Time m	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 22 23 24 25 26 27 28 28 29 20 20 20 20 20 20 20 20 20 20 20 20 20	76 64 61 68 68 68 68 67 61 61 74 86 87 77 77 81 77 78 77 77 77 77 77 77 77 77 77 77 77	4 P 2 P 2 P 5 P 4 P 12 N 5 P 4 P 12 N 5 P 4 P 12 N 6 P 12 N 6 P 12 N 6 P 12 P 6 P 12 P 12 P 12 P 14 P 12 P 14 P 14 P 14	34 38 32 32 35 47 48 43 43 43 44 50 44 45 66 45 56 46 45 56 47 56 47 56 47 56 66 47 56 56 56 56 56 56 56 56 56 56 56 56 56	6 A 6 A 6 A 7 A 7 A 6 A 6 A 6 A 6 A 6 A	55 57 48 46.5 50 51.5 58.5 66.5 52.5 68.5 64.5 64.5 64.5 65.5 63.5 63.5 63.5 63.5 63.5 64.5 65.5 66	49 49 45 57 51 63 55 58 57 91 93 65 66 70 70 77 88 82 89 96 74 75 75 83 91 76 76	11.4 8.8 9.1 8.9 12.0 7.5 4.2 5.0 10.5 0.0 2.0 12.5 10.9 10.0 12.5 11.0 0.6 0.3 9.5 11.0 0.3 9.5 0.0 2.5 11.0 0.0 7.9 9.5 11.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.	N N NW NW NW NW NW NW NW NW NW SW SW SW SW SW SW SW SW SW S S S S	41 213 117 79 65 162 175 185 94 43 108 124 45 59 81 24 45 136 141 42 93 84 134 228 190 67 96 66 79 91 05 105	13 40 20 8 11 11 12 22 22 20 10 8 14 18 8 8 13 17 25 12 6 8 11 22 12 25 25 20 10 8 11 11 11 25 25 25 11 25 15 15 15 15 15 15 15 15 15 15 15 15 15	6 P 6 A 0 A 5 P 2 A 4 A 0 A	8 P 7 P 3 A 6 P 12 P 12 N	.02 .87 .32 0.1 .11 .12 .79	

^{*} Based on least time required to blow one mile.

MONTILL	SUMM	
MAY, 1951	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.487 Minimum 29.436 Mean semi-daily 29.890 Range 1.051	30.39 29.51 29.96 0.88	30.62, 1936 29.10, 1938
Air Temperature, in Degrees F.		
Highest 87, 16th Lowest 31, 14th Mean 58,9 Range 56 Highest mean daily 73, 22nd Lowest mean daily 42.5, 11th Mean maximum 71.7 Mean minimum 46.0 Greatest daily range 43, 14th, 19th Least daily range 6, 24th Degree-days of heating load for month 211 Degree-days cumulative for season 6032	86.1 31.2 57.1 54.9	94.5, 1896, 1911 24.0, 1900
Precipitation, in inches		
Precipitation	3.60	7.44, 1931; .48, 1903
Snow	12	20, 1901, 1945; 5, 1903
Wind, in miles		
Total movement3256Greatest daily movement228, 24thLeast daily movement41, 1stMean hourly velocity4.4Maximum velocity40, 2nd	4504 6.1 28.8	5946, 1907; 2180, 1894
Wind, direction		
Prevailing direction NW North, days 10 Northeast, days 2 East, days 2 Southeast, days 1 South, days 4 Southwest, days 5 West, days 2 Northwest, days 5	W.	
Weather		
Mean relative humidity, percent. 72.2 Mean cloudiness, percent 37.8 Number of clear days 9 Number of fair days 12 Number of cloudy days 10 Number of hours bright sunshine 211.5 Percent of possible hours of bright sunshine 46.8 Thunder and Lightning 9th Last Frost 14th	60.7 52.1 11 11 9 252 55.6	72, 1948; 30, 1923 21, 1944; 0, 1927 18, 1949; 5, 1923 20, 1927; 2, 1941 350, 1944; 137, 1927

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1950.

REMARKS

The degree days for the month of May were slightly below normal, the reading being 211 as compared to a normal of 245. This brought the cumulative degree days for the season to a total of 6,032, as compared to a normal of 6,700. This was somewhat higher than the mild winter of 1948-1949 and is slightly below the cumulative degree days for the winter of 1949-1950. The cumulative degree days for the season to the end of May 1949 were 5,551 and the cumulative degree days for the heating season to the end of May 1950 were 6,335.

In general the weather during the month was mild and dry. The last frost for the month occurred on May 14. The precipitation was below normal with a reading of 2.96 inches as compared to a normal of 3.60 inches. Following the rainfall of .14 inches on April 25 no appreciable amount of rain fell until May 11. A total of 1.19 inches fell on May 11 and 12. There was very little rainfall following this until May 24 and 27 when a total of 1.50 inches was

recorded.

Although the rainfall was below normal, the relative humidity continued to be far above normal as it had in preceding months. The mean relative humidity for the month was 72.2 per cent as compared to a normal of 60.7 per cent. The number of hours of bright sunshine of 211.5 was low as compared to the normal of 252 hours.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 750

June 1951

Meteorological Observations

FOR

JUNE

1951

EARLE F. COX

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

		Te	empera	ature	-	,			Wind		I	Precipita	tion	
	Maximum		Min	nimum		Humidi	f Hour nsbine	nsbine	+	9				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunsbine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15	88 78 67 80 80 77 70 74 77 67 68 76 77 55	3 P 1 P 5 P 3 P 4 P 3 P 1 P 1 P 2 P 5 P 1 P	53 65 60 58 61 48 49 39 47 53 48 47 49 51 48	5 A 4 A 5 A 6 A 5 A 4 A 5 A 4 A 5 A 4 A 5 A 4 A 5 A	70.5 71.5 63.5 69 70.5 62.5 59.5 62 60 58 61.5 63 53	73 89 95 84 73 62 64 59 61 60 61 78 94	4.1 0.0 0.0 1.7 6.6 9.3 12.4 11.2 9.2 5.5 2.2 10.4 5.9 0.0 0.1	SW S S N SW N SW N SW N SE N N SE NW W	64 85 70 76 127 38 105 63 78 135 118 73 98 87 156	20 7 22 17 5 28 13 20 28 25 11 13 14	6 P 1 A 0 A 3 P	9 P 8 P 6 A 8 P	0.06 1.42 0.33 0.27	
16 17 18 19 20 21 22 23	74 80 83 88 83 82 82 76	7 P 4 P 1 P 2 P 4 P 4 P 4 P 6 P	49 55 58 58 60 66 55	4 A 5 A 5 A 6 A 5 A 4 A 5 A 6 A	61.5 67.5 70.5 73 71.5 74 69 70.5	88 91 77 72 74 82 73 94	2.4 2.0 4.2 6.2 8.0 6.4 10.1 1.4	N SE SE S S SE SE	68 69 69 73 112 160 88 134	9 11 10 8 10 12 13	5 P 10 A 7 P	10 A 6 P 1 P	0.01 0.01 0.07	
24 25 26 27 28 29 30	87 84 78 82 81 72 73	4 P 2 P 5 P 5 P 2 P 11 A 12 N	58 67 50 62 59 61 60	5 A 5 A 5 A 3 A 6 A 5 A 4 A	82.5 75.5 64 72 70 66.5 66.5	74 69 59 88 77 89 91	11.2 12.0 13.8 6.1 12.3 1.1 2.7	SW NW NW S SE S	59 210 62 85 75 92 74	14 25 10 14 9 18 9	12 N 3 P	8 P 4 A	0.08 0.17 T 0.61 0.01	

^{*} Based on least time required to blow one mile.

WALTER S. FOSTER, Observer

MONTILI	SUMIM	
JUNE 1951	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.103, 12th Minimum 29.738, 27th Mean semi-daily 29.931 Range 365	30.30 29.55 29.95 0.75	30.70, 1947 29.24, 1902
Air Temperature, in Degrees F.	1 1	
Highest. 88, 1st, 19th Lowest 39, 8th Mean. 77,2 Range 49 Highest mean daily 82,5,24th Lowest mean daily 52,15th Mean maximum 76,5 Mean minimum 55,3 Greatest daily range 35,1st,8th Least daily range 4,14th Degree-days of heating load for month 68 Degree-days cumulative for season 6100	91.2 40.0 65.7 51.3	101, 1919 34, 1891
Precipitation, in inches	ľ	
Precipitation	3.75	9.68, 1922; 0.76, 1908
Naximum precip. in 24 hrs 1.42, 14th Number of days with .01 or more9	11	17, 1922, 1945; 4, 1908
Wind, in miles	ļ	
Total movement2803Greatest daily movement210 25thLeast daily movement38.6thMean hourly velocity3.9Maximum velocity28,7th, 10th	3585 5.0 24.6	4571, 1908; 1409, 1906 ~ 48, 1939
Wind. direction		
Prevailing direction SSW North, days 6 Northeast, days 1 East, days 0 Southeast, days 6 South, days 8 Southwest, days 4 West, days 2 Northwest, days 3	wsw	
Weather		
Mean relative humidity, percent. 77.2 Mean cloudiness, percent 59.3 Number of clear days 6 Number of fair days 10 Number of cloudy days 11 Number of hours bright sunshine 178.5 Percent of possible hours of bright sunshine 39.0 Thunder and Lightning 12st Frost	66.9 51.1 10 12 8 257 54.0	71, 1903; 28, 1908 22, 1908, 1941, 1943; 0, 1949 23, 1912; 3, 1941 22, 1903; 1, 1923 365, 1947; 102, 1903

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1950.

REMARKS

The dry period, which began May 28, extended until June 13 before any rainfall occurred. Following this period of drought rainfall occurred on nine of the eighteen remaining days in June. Total precipitation for the month was 3.04 inches as compared to a normal of 3.75 inches.

Cloudy weather prevailed throughout the month. Bright sunshine occurred only 39 per cent of the possible hours. The normal for the month is 54 per cent. As with preceding months, the relative humidity continued high. The mean for the month was 77.2 per cent as compared to a normal of 66. 9per cent.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 751

July 1951

Meteorological Observations

FOR

JULY 1951

EARLE F. COX

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

	Temperature					r,	ø	Wind Precipitation					tion	
	Ma	ximum	Min	nimum		Iumidi	f Hour isbine			20				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4	74 80 84 83	3 P 5 P 6 P 2 P	64 53 51 58	6 A 5 A 5 A 4 A	69 66.5 67.5 70.5	89 70 65 78	4.8 4.9 10.1 9.2	W W W S	80 101 66 95	7 12 15 14	8 A	10 A	0.10	
5	83	12 N	69	3 A	76	81	7.8	s	142	20	$ \begin{bmatrix} 1 \text{ A} \\ 3 \text{ P} \\ 7 \text{ P} \end{bmatrix} $	2 A 4 P 12 P	.09	
6 7 8 9 10 11 12	79 89 87 85 87 88 83	3 P 3 P 3 P 3 P 2 P 2 P	56 48 57 62 63 63 67	3 A 5 A 5 A 4 A 3 A 4 A 3 A	67.5 68.5 72 73.5 75 75.5	69 63 69 65 82 82 95	7.3 12.0 10.7 11.4 7.7 8.9 0.	NW W S S S SE SE	157 41 76 155 130 70 38	22 5 8 13 8 10 5	5 P 5 P 4 A 9 A	6 P 6 P 5 A 8 P	.08	
13 14 15 16 17 18 19	82 86 88 91 84 77 75	4 P 5 P 4 P 4 P 1 P	65 60 57 61 65 62 65	3 A 4 A 5 A 4 A 5 A	73.5 73 72.5 76 74.5	73 67 67 75 90 78 89	10.2 10.1 12.8 10.0 1.2	N W W S N	60 42 45 76 79 76 64	11 5 7 10 7 7 20	5 A 3 P	9 A 8 P	.32 .40	
20 21 22 23 24 25 26 27 28 29 30 31	76 80 80 86 84 87 91 88 76 78 90	3 P 3 P 3 P 3 P 3 P 3 P 3 P 2 P 4 P 3 P 3 P	55 50 66 69 63 55 58 67 66 63 67	5 A 4 A 4 A 0 A 4 A 6 A 5 A 3 A 4 A	70 65.5 65 73 77.5 73.5 71 74.5 77.5 71 70.5 78 78	64 66 84 73 73 67 74 70 87 88 78	2.0 12.5 12.0 1.5 8.2 10.4 11.6 9.7 9.1 0.1 0 7.4 6.7	SW NW S S NW W SW W N SE N W SE	176 91 142 71 46 63 88 64 98 61 57	22 12 12 10 6 7 10 10 8 5 8	6 A 12 N	8 A 7 P	.01	

^{*} Based on least time required to blow one mile.

MONTHLI	SUMIN	** *** *
JULY 1951	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.119, 18th Minimum 29.436, 5th Mean scmi-daily 29.962	30.27 29.59 29.96	30.50, 1892 29.27, 1932
Rang	0.68	0.97, 1892; 0.47, 1938
Air Temperature, in Degrees F.	,	
Highest 91, 16th, 26th Lowest 48, 7th Mean 72, 3 Range 43 Highest mean daily 78, 30th, 31st Lowest mean daily 65, 21st Mean maximum 83, 5 Mean minimum 61, 0 Greatest daily range 41, 7th Least daily range 10, 28th Degree-days of heating load for month — Degree-days cumulative for season —	93,9 46.4 70.8 47.7	104, 1911 40, 1890, 1898 66.3, 1891; 76.0, 1949
Precipitation, in inches		
Precipitation	4.10	14.51, 1897; 0.70, 1929
Snow	11	20, 1915; 4. 1924
Wind, in miles		
Total movement2609Greatest daily movement157, 6thLeast daily movement38, 12thMean hourly velocity3,5Maximum velocity22, 6th, 20th	3422 4.6 28.7	5097, 1909; 1109, 1894
Wind direction		
Prevailing direction SW North, days 4 Northeast, days 0 East, days 0 Southeast, days 4 South, days 8 Southwest, days 3 West, days 8 Northwest, days 4	SW	
Weather		
Mean relative humidity, percent	68.4 50.9 10 14 7 268	70, 1943; 31, 1924 22, 1923; 0, 1915 24, 1909, 1921; 4, 1946 18, 1889; 0, 1910 371, 1910; 180, 1931

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1950.

REMARKS

The July weather was near normal in nearly all respects. No extremes were recorded but the high mean relative humidity of previous months was continued as was also the more than normal amount of cloudiness. The mean relative humidity of 77.1 was considerably above the normal of 68.4. The mean cloudiness at 60.3 per cent was also above the normal of 50.9 The per cent of possible hours of bright sunshine recorded at 51.1 was below the normal of 58.

The readings for air temperature and rainfall were close to normal. The rainfall was slightly above normal for the month. It has been above normal for every month except June during 1951. This has resulted in an accumulated precipitation of 26.8 inches to the end of July as compared to a normal of 25.3.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 752

August 1951

Meteorological Observations

FOR

AUGUST

EARLE F. COX

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

		Te	mpera	ature		ty	ø,		Wind		Precipitation			
	Maximum		mum Minimum			Humidi	of Hour nshine	2	i,	es F S				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1	84	5 P	56	12 P 5 A	70	85 73	4.6 13.1	SE	131	20	9 A	12 N	.05	
2 3 4 5	84 72	4 P 2 P	48 57	5 A 4 A	66 64.5	81	2.7	NW S	113	17	7 P	8 P	.14	l
4	74	4 P	55	4 A	64.5	70	11.4	N	7.3	7				
$\begin{bmatrix} 5 \\ 6 \end{bmatrix}$	74 76	4 P 5 P	48 45	4 A 6 A	61 60.5	71 70	12.7 12.5	N NE	74	12				
7	79	1 P	48	5 A	63.5	65	9.4	S	66	12				
8	67	3 P	59	5 A	63	90	0	E	67	9	2 A	11 A	.56	
9	84 85	3 P 5 P	58 71	5 A 4 A	71 78	87 86	9.8 4.1	SW	71 7C	9	9 A	10 A	.07	
11	88	4 P	68	6 A	78	75	10.0	NW	73	6	/ A	IOA	.07	
12	81	2 P	62	6 A	71.5	80	2.5	N	39	9	2 P	6 P	.04	
13 14	86 87	4 P 4 P	65 57	5 A 5 A	75.5 72	79 79	9.6	NW S	87 59	12				
15	87	1 P	69	3 A	78	88	4.6	SE	97	10	4 P	6 P	.03	
16	75	2 P	63	12 P	69	92	0.3	N	62	32	6 P	11 P	1.11	
17	77	4 P	62	7 A	69.5	92	0	N	59	13	6 P	7 P	.87	
18 19	73 79	6 P 5 P	60 50	6 A 6 A	66.5 64.5	79	10.2	NW NW	70 28	11 6				
20	78	3 P	52	6 A	65	94	6.2	NW	37	14	10 A	5 P	.24	İ
21	72	5 P	59	0 A	65.5	95	0	S	64	1.3	12 N	5 P	.19	
22 23	77	11 A	64	6 A 5 A	70.5	64	9.6	NW	177	18				1
23	72 72	4 P 12 N	50 49	5 A 5 A	61 60.5	80	11.0	NW N	56	10				
25	75	2 P	56	5 A	65.5	87	4.9	NW	61	11	9 A	6 P	.26	
26	77	6 P	55	2 A 5 A	66	86	8.1	NW	65	9				1
27 28	76 82	6 P	57 59		66.5	87 89	7.6	SW	33	10			1	
28	82 87	4 P	59	2 A 5 A	73	87	7.4	SW SW	39	6				
30	90	3 P	59	5 A	74.5	84	8.3	sw	36	3				
31	89	2 P	62	6 A	75.5	85	7.3	s	92	13		1	1	1

^{*} Based on least time required to blow one mile.

WALTER S. FOSTER, Observer

MONTHE		17 11 (1
AUGUST 1951	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.186, 24th Minimum 29.700, 3rd Mean semi-daily 30.044 Rang .486	30.32 29.61 30.00 0.71	30.50, 1934 28.87, 1930
Air Temperature, in Degrees F.		
Highest 90, 30th Lowest 45, 6th Mean 68.4 Range 45 Highest mean daily 78, 10th, 11th, 15th Lowest mean daily 60.5, 6th, 24th Mean maximum 79,3 Mean minimum 57.8 Greatest daily range 36, 2nd Least daily range 8, 8th Degree-days of heating load for month Degree-days cumulative for season —	91.6 43.4 68.6 48.1	100, 1918, 1948 34, 1940 62.4, 1903
Precipitation, in inches		
Precipitation	4.08	8.40, 1948; .31, 1894
Maximum precip. in 24 hrs 1.11, 16th Number of days with .01 or more 11	11	16, 1892, 1933; 4, 1899
Wind, in miles	ļ	
Total movement 2133 Greatest daily movement 177, 22nd Least daily movement 28, 19th Mean hourly velocity 2,86 Maximum velocity 32, 16th	3127 4.2 22.7	4,217, 1910;1,920, 1894
Wind, direction		,
Prevailing direction WNW North, days 6 Northeast, days 1 East, days 1 Southeast, days 2 South days 5 Southwest, days 6 West, days 0 Northwest, days 10	SW	
Weather		
Mean relative humidity, percent	70.2 49.7 9 13 9 237	67, 1901, 1946; 27, 1923 23, 1941; 0, 1915, 1929 25, 1912; 3, 1939 18, 1901, 1928; 2, 1910, 1923 332, 1941; 152, 1915, 1929

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1950.

The month of August was near normal in most respects. The mean daily temperature was 68.4 while the normal for this month is 68.6. The highest for the month was 90.0 as compared to a normal of 91.6.

The precipitation during the month was 3.56 inches which compares to a normal 4.08. Of the total amount during the month 1.98 inches fell on the 16th and 17th. There were eleven days during the month when a total of 0.01 of an inch or more fell. This is equal to the number of days during the month on which we can expect 0.01 of an inch or more of rain. While the precipitation during the month was below normal, the accumulated precipitation for the year is still approximately one inch above the normal.

The mean relative humidity for the month was 82.3 which compares to a normal of 70.2 for the month. The mean relative humidity has been running far above normal and this is still the case during the month of August. In spite of the high mean relative humidity the cloudiness and hours of bright sunshine for the month were about normal.

The total wind movement for the month was 2133 miles which compares to a normal of 3127. There were no wind storms during the month and the maximum velocity was 32 miles per hour which was reached on the 16th.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series Bulletin No. 753

September 1951

Meteorological Observations

FOR

SEPTEMBER 1951

EARLE F. COX

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

> Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Те	mpera	ture		, i	vo.		Wind			Precip	itation	
	Max	imum	Mir	imum		fumidi	f Hournshine			ý				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hr.*	Began	Ended	Water	Degree Days
1 2	61 58 67	8 A 2 P 3 P	55 54 54	12 p 6 a 6 a	58 56 60.5	89 94 92	$0.0 \\ 0.0 \\ 0.0$	N N W	113 55 53	9 7 8	5 A 1 A	6 A 11 P	.02 .87	7 9 4.5
1 2 3 4 5 6 7 8 9	76 78 66 78 72 79	2 P 1 P 2 P 2 P 4 P	50 46 50 61 49 43	5 A 5 A 5 A 5 A	63 62 58 69.5 60.5	75 79 91 82 66 75	11.8 10.4 2.7 8.2 12.1 11.3	NW S NW NW	100 20 61 123 141 42	12 4 12 12 12 7 7	12 N	4 P	.10	2 3 7 4.5
10 11 12 13 14	82 71 85 81 84	4 P 3 P 5 P 3 P 3 P 1 P	46 59 61 64 67	6 A 6 A 3 A 6 A 7 A 5 A	64 65 73 72.5 75.5	74 91 88 83 85 94	10.3 0.5 5.9 4.8 6.0	S SE SE SE SE	54 102 57 99 160	7 9 13		10 р	1.03	4 1
15 16 17 18 19 20	67 79 74 73 77 78	6 P 4 P 3 P 4 P 4 P 5 P	65 52 54 50 47 46	8 A 6 A 7 A 7 A 7 A	66 65.5 64 61.5 62 62	89 82 85 80 88	0.5 7.0 10.8 8.8 9.2 10.0	NE NE	101 51 53 39 66 25	13 7 8 8 6 7	6 л	10 P	1.03	1 3.5 3
21 22 23 24 25 26 27 28 29 30	82 80 74 74 69 62 70 61 56 49	3 P 2 P 2 P 1 P 10 A 3 P 8 P 12 A 2 P 4 P	56 59 50 43 47 44 36 49 38 28	3 A 6 A 12 P 6 A 6 A 6 A 6 A 6 A	69 69.5 62 58.5 58 53 53 53 55 47 38.5	87 75 81 74 88 69 79 67 66 71	6.0 8.2 5.0 9.0 5.0 11.1 0.6 10.6 11.2 6.9	W S S S N W S W N W SE	67 178 196 52 68 129 80 229 213 42	10 12 18 7 13 11 13 25 25 5	4 A 8 A 7 P	8 A 8 P 12 P	.05	6.5 7 12 12 10 18 26.5

^{*} Based on least time required to blow one mile.

WALTER S. FOSTER, Observer

MONTHEL		
SEPTEMBER 1951	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)	20.45	20.65 1021
Maximum 30.407 Minimum 29.722	30.45	30.65, 1924 28.41, 1938
Mean semi-daily 30.095 Range 0.685	30.00 .88	1.99, 1938; .57, 1910
Air Temperature, in Degrees F.		
Highest 85, 12th Lowest 28, 30th Mean 61,4 Range 57 Highest mean daily 75.5, 14th Lowest mean daily 38.5, 30th Mean maximum 72.1 Mean minimum 50.8 Greatest daily range 36, 9th, 10th Least daily range 2, 15th Degree-days of heating load for month, 147 Degree-days cumulative for season .147	87.7 33.2 61.7 54.5	97, 1895, 1929 24.5, 1914
Precipitation, in inches		
Precipitation 2.63 Snow	4.24	14.55, 1938; .52, 1914
Maximum precip. in 24 hrs1.03, 15th Number of days with .01 or more7	10	16, 1933; 3, 1903
Wind, in miles		
Total movement 2769 Greatest daily movement 229, 28th Least daily movement 20, 5th Mean hourly velocity 3.9 Maximum velocity 25, 28th, 29th	3271 4.5 25.9	4,686, 1896; 1,414, 1894 80, 1938
Wind, direction		
Prevailing direction WSW North, days	WSW	
Weather		
Mean relative humidity, percent	73.6 49.9 10 10 10 201 54.9	70, 1934; 23, 1947 19, 1932; 2, 1907, 1928 19, 1908; 3, 1889 20, 1934; 3, 1941, 1948 255, 1916; 106, 1934
Thunder and lightning	Sept. 21	Oct. 13, 1909; Aug. 22, 1894-95

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1950.

The weather for the month of September was mild and dry. The mean daily temperatures for the month averaged 61.4, only 0.3 degrees below the normal mean daily temperature of September. Although the low precipitation of 2.63 inches was considerably under the 4.24 inch average for the month, higher than normal relative humidity helped offset the effect of the light rainfall. Nearly 40 percent of the month's rainfall was accounted for by the storm of the 15th.

Heating load for the month was 147 degree-days, slightly more than normal, but one-sixth of the total came on the 30th, accompanying the first frost of the season.

While the first frost of the season occurred on the 30th, 9 days later than normal, it may be noted that the records show no first frost to have occurred on the average date, the 21st, in this century.

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series Bulletin No. 754

October 1951

Meteorological Observations

FOR

OCTOBER 1951

EARLE F. COX

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

> Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

-		Te	mpera	ture		y.	8		Wind			Precip	itation	
	Max	timum	Mir	imum		Humidity	Hour Ishine			ý				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative H	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hr.*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 7 8 8 9 10 11 12 13 11 15 116 17 18 19 22 22 23 4 25 26 27 7 8 29	71 70 64 75 77 56 64 63 55 52 65 72 73 75 66 68 63 66 71 72 59 74 69 42	2 P 12 A 12 A 1 P 2 P 0 A 2 P 4 P 2 P 12 A 2 P 4 P 2 P 3 P 2 P 4 P 2 P 2 P 1 P 2 P 1 P 2 P 1 P 2 P 1 P 2 P 1 P 2 P 1 P 2 P 1 P 2 P 1 P 2 P 1 P 2 P 1 P 2 P 1 P 2 P 1 P 2 P 1 P 2 P 1 P 2 P 1 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 3 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 3 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2	45 54 55 55 55 63 55 46 45 45 33 41 41 28 30 51 52 54 40 47 40 40 40 40 40 40 40 40 40 40 40 40 40	6 A 4 A 4 A 6 A 6 A 7 A 12 P 7 A 7 A A 7 A A 7 A A 7 A A 7 A A 7 A A 12 P 7 A 7 A A 12 P 7 A 12 P 7 A 14 A 17 A 17 A 18 A 18 A 18 A 18 A 18 A 18	N 58 62 61 65 72.5 66 51 54.5 46.5 53 49.5 61 58 62 57.5 46.5 58.5 56.5 57 58 46.5 57 58 40 41.5	77 84 87 88 88 85 73 76 65 68 82 75 70 88 81 81 64 69 71 70 60 60 57 81 85 81	7.5 3.2 0.0 2.4 7.2 8.4 0.0 8.1 10.7 8.5 1.0 8.4 10.5 9.7 3.6 6.1 0.0 9.0	SW S N N N N N N N N N N N N N N N N N N	52 23 42 122 122 143 144 113 154 182 190 140 61 56 91 44 58 80 94 126 114 213 88 80 114 214 214 315 315 315 315 315 315 315 315 315 315	5 5 4 10 12 20 20 14 20 20 10 20 20 10 8 13 4 13 14 17 7 10 8 8 8 10 12 10 10 10 10 10 10 10 10 10 10 10 10 10	11 P 4 A 0 A 2 A 0 A 2 P 0 A 8 A	12 P 2 P 3 A	1.66 .36 .93 .03	S
30 31	54 64	2 P 2 P	30 45	2 A 2 A	42 54.5	63 67	4.0 3.0	s ssw	63 108	8 28				

^{*} Based on least time required to blow one mile.

MARIUS P. VAROUDES, Observer

MONTHLI	30 WIWIZ	XICI
OCTOBER 1951	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.674, 14th Minimum 29.552, 28th Mean semi-daily 30.142	30.53 29.42 30.06	30.68, 1929 29.00, 1926
Range 1.122	1.11	1.47, 1926; .76, 1899
Air Temperature, in Degrees F.		
Highest 82, 5th Lowest 27, 14th Mean 54.8 Range 55 Highest mean daily 72.5, 5th Lowest mean daily 40, 28th Mean maximum 65.3 Mean minimum 44.2 Greatest daily range 45, 14th Least daily range 4, 28th Degree-days of heating load for month 328.5 Degree-days cumulative for season 483	79.4 23.4 50.5 56.0	90.5, 1908 17, 1936 57.4, 1947; 43.2, 1890
Precipitation, in inches		
Precipitation 4.66	3.29	8.81, 1911; .01, 1924
Snow None Maximum precip. in 24 hrs. 1.66, 7th Number of days with .01 or more 8	9	15, 1913; 1, 1897, 1924
Wind, in miles		
Total movement 3216 Greatest daily movement 231, 20th Least daily movement 23, 2nd	4074	5,467, 1910; 2,540, 1894
Mean hourly velocity 4.32 Maximum velocity 28, 27th, 29th, 31st	5.4 29.5	42, 1937
Wind, direction		
Prevailing direction WNW North, days 7 Northeast, days 1 East, days 1 Southeast, days 3 South, days 8 Southwest, days 3 West, days 0 Northwest, days 8	W	
Weather		
Mean relative humidity, percent 74.8 Mean cloudiness, percent	69.0 48 10 10 11 175.0	67, 1932; 18, 1924 21, 1938; 1, 1911 17, 1924; 3, 1938 19, 1896; 3, 1924 232, 1923, 1938; 91, 1913
bright sunshine 56.1 Thunder and lightning None	51.3	

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1950.

In general, pleasant weather prevailed throughout the month of October, providing an opportunity for the observance of brilliant foliage brought out by the frost during the last week of September. Although the precipitation was above normal with a recording of 4.66 compared to a normal of 3.29, the mean cloudiness was far below normal with a recording of 35.5 percent compared to a normal of 48 percent. The percent of possible hours of bright sunshine was also somewhat above normal with a reading of 56.1 percent compared to a normal of 51.3 percent.

The mean daily temperature was above normal with a reading of 54.8 compared to a normal of 50.5. The mean relative humidity continued to be considerably above normal. The figure for the month of October was 74.8 percent compared to a normal of 69.0 percent. The mean relative humidity has been above normal every month for over a year. In some months this figure has been as high as 12 percent above normal. The relative humidity of 74.8 for the month of October is closer to the normal than the relative humidity in most of the months during the previous year.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series Bulletin No. 755

November 1951

Meteorological Observations

FOR

NOVEMBER 1951

EARLE F. COX

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

> Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Ter	mpera	ture		33	s ty		Wind			Precipi	tation	
	Max	imum	Min	imum		umidit	Hours							
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8	42 36 46 43 45	0 A 12 M 12 M 2 P 2 P	31 27 30 22 23	12 P 6 A 12 P 6 A 7 A	36.5 31.5 38 32.5 34	80 78 83 69 71	$0.0 \\ 0.0 \\ 0.0 \\ 7.5 \\ 6.2$	N N N SE	225 133 119 110 90	15 11 12 7 13	5 A 7 P 0 A	1 P 12 P 8 A	.57 .37 2.04	.5
9	38 47 49 49	1 P 1 P 12 M 4 P	20 28 35 33	7 A 11 P 7 A 8 A	29 37.5 42 41	70 86 79 66	7.5 0.0 4.6 9.2	N N W N W	99 99 129 269	8 13 32 22	8 A	11 Р	1.75	
10 11 12 13 14	59 51 50 66 65	2 P 2 P 2 P 2 P 11 A	26 29 25 30 51	6 A 6 A 6 A 2 A 0 A	42.5 40 37.5 48 58	78 71 80 82 87	7.1 7.5 9.0 6.1 2.6	S W N W S E S E	36 149 35 74 93	7 20 5 10 7	11 A	11 г	.46	:
15 16 17 18	63 47 46 36	1 P 11 A 10 A 2 P	52 38 39 27	7 A -1 A 7 A 6 A	57.5 42.5 42.5 31.5	83 85 80 71	0.0 0.0 3.3 8.6	SE SW N W	163 149 168 253	$ \begin{array}{c c} 22 \\ 10 \\ 17 \\ 20 \end{array} $	8 A	9 Р	.44 T	
19 20 21 22 23	36 32 43 56 59	1 P 11 A 3 P 4 P 2 P	25 20 16 18 40	3 A 2 A 7 A 5 A 9 P	30.5 26 29.5 37 49.5	71 71 70 70 75	$ \begin{array}{c c} 6.7 \\ 3.8 \\ 8.6 \\ 1.7 \\ 0.6 \end{array} $	N W N W N W S E S E	196 216 46 110 179	$ \begin{array}{c c} 20 \\ 20 \\ 8 \\ 11 \\ 13 \end{array} $	5 P	12 в	.05	
24 25 26 27	48 42 44 30	8 A 2 P 12 M 12 M	$ \begin{array}{r} 44 \\ 23 \\ 27 \\ 24 \end{array} $	7 A 7 A 3 A 8 A	46 32.5 35.5 27	85 73 90 78	0.0 8.3 2.1 8.7	SE NW N	164 100 155 413	28 10 32 28	0 A 2 A	9 A 10 A	.63	2.5
28 29 30	26 46 49	7 P 3 P 2 P	8 17 16	7 A 7 A	31.5 32.5	79 85 88	2.8 8.0 8.0	N N E N	49 17 20	4 5 2				

^{*}Based on least time required to blow one mile.

Roger F. Johnson, Observer

NOVEMBER 1951	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum	30.59 29.30	30.87, 1932 28.73, 1904
Mean semi-daily	30.05 1.29	1.84, 1904; 0.89, 1943
Air Temperature, in Degrees F.		
Highest 66, 13th Lowest 8, 28th Mean 37.2 Range 58 Highest mean daily 58, 14th Lowest mean daily 17, 28th Mean maximum 46.3 Mean minimum 28.1 Control of the billion 28.21	66.0 12.9 38.9 53.0	82, 1950 4, 1938 44.1, 1931; 33.6, 1901
Greatest daily range	783	
Precipitation, in inches		
Precipitation 6.39 Snow 3.0	3.41 2.34	8.64, 1927; 0.63, 1917 13.50, 1938
Maximum precip. in 24 hrs2.04, 3rd Number of days with .01 or more9	9	22, 1921; 2, 1904
Wind, in miles		, , ,
Total movement 4058 Greatest daily movement 413, 27th Least daily movement 17, 29th	4531	5,978, 1906; 2,589, 1889
Mean hourly velocity	6.3 30.3	48, 1948, 1950
Wind, direction		
Prevailing direction N North, days 11 Northeast, days 2 East, days 0 Southeast, days 7 South, days 0 Southwest, days 2 West, days 3 Northwest, days 5	WNW	
Weather		
Mean relative humidity, percent	70.6 55.1 6 9 15 121 41.3 Nov. 6	78, 1947; 34, 1917 15, 1903; 0, 1948 16, 1912; 4, 1889, 1930, '32 '35 24, 1927; 9, 1905, 1917, 1949 182, 1903; 44, 1948 Oct. 10, 1925; Dec. 13, 1941

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1950.

The mean daily air temperature during the month of November was slightly below normal, and the rainfall for the month was far above normal. The lowest temperature recorded during the month was 8; this was recorded on November 28. The highest temperature was 66, and the mean daily temperature during the month was 37.2. The normal mean daily temperature for the month is 38.9.

As a result of the low mean daily temperature for the month, the number of degree-days of heating load for the month was slightly above normal. The reading for this was 835 compared to a normal of 783. The accumulative degree-days for the season stand at 1310.5 which is very close to normal.

The total precipitation for the month was 6.39 inches. This is far above the normal rainfall of 3.41 inches for the month. The first snowfall occurred on November 3, and the total snowfall for the month was 3 inches. The maximum precipitation in 24 hours during the month occurred on the 3rd when 2.04 inches of rainfall was recorded. On nine days of the month the rainfall exceeded .01 inches. The wind movement during the month was not excessive, the highest velocities being recorded on the 8th and 26th; the velocity reached 32 miles per hour.

The mean relative humidity for the month continued far above normal with a reading of 77.9 compared to a normal of 70.6. This tendency has continued for over a year. The mean cloudiness for the month was slightly below normal, and the percent of possible hours of bright sunshine was slightly above normal.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series Bulletin No. 756

December 1951

Meteorological Observations

FOR

DECEMBER 1951

EARLE F. COX

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W.

Height of barometer above ground, 36 ft. Above sea level, 253.5 ft.

Height of wind instruments, 67 ft. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Ter	mpera	ture		8 tt							oitation		
	Max	imum	Min	imum		umidit Hours sline									
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max, Miles per Hour*	Began	Ended	Water	Snow	
$\begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \end{bmatrix}$	46 51 50 42	2 P 3 P 3 P 2 P	22 33 22 25	7 A 7 A 8 A 4 A	34 42 36 33,5	92 91 92 92	6.7 7.6 7.9 0.7	N N N	$\frac{4}{102}$ $\frac{24}{12}$	7 13 6 1					
5 6 7	59 57	8 Р 3 Р	38 51	12 P 7 A	$\frac{48.5}{54}$	9 7 98	$0.0 \\ 0.0$	S E	$\frac{53}{142}$	20 9	3 A	11 Р	.61		
7 8	59 61	3 р 12 м	48 54	8 A 1 A	53.5 57.5	96 7 2	$\frac{2.1}{6.6}$	S N E	83 182	$\frac{13}{25}$	8 P	9 P	.13		
9	47 43	5 Р 12 м	$\frac{39}{34}$	8 A 8 A	$\frac{43}{38.5}$	87 75	0.0	W N	$\frac{70}{280}$	$\frac{25}{22}$	2 A	2 P	.56		
$\frac{11}{12}$	35 38	1 P 1 P	$\frac{18}{23}$	5 A 6 A	$\frac{26.5}{30.5}$	86 85	$\frac{4.2}{7.0}$	N W W	$\frac{72}{100}$	10 20					
13 14	$\frac{26}{28}$	2 P 3 P	19 15	7 A 7 A	$22.5 \\ 21.5$	69 81	3.5 4.0	N N W	$\frac{222}{115}$	20 13	5 P	12 р	.20	3.0	
$\frac{15}{16}$	28 15	11 A 2 P	$\frac{20}{5}$	2 P 7 A	24 10	89 70	6.2 8.3	M.	$ \begin{array}{r} 224 \\ 224 \end{array} $	$\frac{40}{20}$	0 A	10 A	.21	4.0	
17 18	$\begin{array}{c c} 18 \\ 26 \end{array}$	3 P 8 P	-8 7	6 A 2 A 7 A	5 16.5	78 85	7.6 0.0	S E N W	$\frac{63}{107}$	$\frac{7}{32}$	10 A	12 Р	.20	2.0	
$\frac{19}{20}$	$\frac{20}{25}$	3 Р 12 м	13 -9	6 A	16.5 8	74 84	7.2 3.5	N W N	$\begin{array}{r} 143 \\ 27 \end{array}$	17 5	0 a 1 p	2 A 12 P	.03 .31	3.5	
$\frac{21}{22}$	53 40	3 P 0 A	$\frac{25}{23}$	0 A 12 P	39 31.5	95 73	$\frac{6.0}{7.3}$	S E W	85 199	13 20	0 A	10 A	1.39	1.1	
$\frac{23}{24}$	32 38	$\begin{array}{cc} 4 & P \\ 2 & P \end{array}$	18 23	12 P	$\frac{25}{30.5}$	$\begin{array}{ c c c }\hline 67\\ 72\\ \end{array}$	7.1 5.4	S E N W	69 76	$\frac{15}{20}$					
$\frac{25}{26}$	24 36	3 P 2 P	11 23	6 A 0 A	$\frac{17.5}{29.5}$	87 89	$0.0 \\ 4.6$	N W N W	31 108	$\frac{2}{22}$	4 P 0 A	12 P 11 A	.18 .32	$\frac{2.0}{4.1}$	
$\frac{27}{28}$	28 22	2 P 3 P	12 -9	2 A 7 A	$\frac{20}{6.5}$	76 77	8.0 6.6	N W	$156 \\ 53$	32					
29 30	34 42	2 Р 12 м	18 31	5 A 10 P	26 36.5	87 95	0.9	S E S E	91 76	10 10	4 P	8 Р	.16		
31	44	2 P	36	7 A	40	94	3.8	N W	88	7				<u> </u>	

^{*}Based on least time required to blow one mile.

ROGER F. JOHNSON, Observer

DECEMBER 1951	Normal l	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30,569, 20th Minimum 29,322, 15th Mean semi-daily 30,091	30 66 29.27 30.05	31.104, 1949 28.85, 1915
Range	1.38	1.78, 1895, 1.01, 1892
Air Temperature, in degrees F.	5.1.3	
Highest .61, 8th Lowest .9, 20th, 28th Mean .29,8 Range .70 Highest mean daily .57.5, 8th Lowest mean daily .5, 17th Mean maximum .37.6 Mean minimum .21.9 Greatest daily range .34, 20th Least daily range .6, 6th Degree-days of heating load for month .1091.5 Degree-days cumulative for season .2419.5	54.3 1.09 27.5 56.2	65.5, 1908 —22.5, 1917 36.9, 1891; 17.1, 1917
Precipitation, in inches		
Precipitation 4.35 Snow 19.7 Maximum precip. in 24 hrs. 1.39, 21st	3.39 8.50	7.77, 1901; .58, 1943 26.50, 1902, T, 1891, 1943
Number of days with .01 or more12	10	17, 1902; 4, 1892, 1943
Wind, in miles		
Total movement 3281 Greatest daily movement 280, 10th Least daily movement 4, 1st Mean hourly velocity 4.4	6.3 31.4	6,694, 1925; 3,239, 1918
Maximum velocity40, 15th Wind, direction	31.4	66, 1949
Wind, threeton Prevailing direction NW North, days 6 Northeast, days 1 East, days 0 Southeast, days 6 South, days 3 Southwest, days 0 West, days 5 Northwest, days 10	WNW	
Weather		
Mean relative humidity, percent 84.0 Mean cloudiness, percent 51.6 Number of clear days 10 Number of fair days 10 Number of cloudy days 11 Number of lours bright sunshine 136.8 Percent of possible hours of bright sunshine 46.8 Thunder and lightning 18th First snow	69.4 54.9 8 9 14 128 45.2 Nov. 6	71, 1929; 39, 1919 15, 1890, 0. 1948 16, '09; 4, '89, '30, '31, '36, '38 25, 1948; 7, '09, '23 172, 1896; 27, 1948

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1950.

ANNUAL SUMMARY

ANNUAL	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.850, Feb. 10th Minimum 28.996, Jan. 15th	30.81 28.95	31.10, 1949 28.41, 1938
Mean semi-daily	30.01 1.85	2.47, 1938; 1.38, 1933
Air Temperature, in degrees F.		
Highest 91, July 16th, 26th Lowest	95.7 -12.2 47.4 107.8	104, 1911 —26, 1902 56.9, 1949; 44, 1902
Lowest mean daily	57.8 36.5	
Least daily range2, Feb. 8th, Sept 15th Degree-days cumulative for year6237.0	6698.	
Precipitation, in inches		
Precipitation	43.70 47.78	59.00, 1938; 30.68, 1908 89.00, 1893; 24.50, 1919
Maximum precip. in 24 hrs. 2.04, Nov. 3rd Number of days with .01 or more122	124	146, 1945; 96, 192
Wind, in miles		
Total movement	52,223	63,571, 1908; 36,257, 189- 705, 1909
Mean hourly velocity47 Maximum velocity40, April 15th, May	5.8	
2nd, Dec. 15th	39.5	80, 193
Wind, direction		
Prevailing direction NW North, days .89 Northeast, days .12 East, days .8 Southeast, days .40 South, days .61 Southwest, days .38 West, days .36 Northwest, days .81	W	
Weather		
Mean relative humidity, percent	67.6 51.7 116 123 126 2.353	60, '98, '01, '02; 41, '08, '24 217, 1941; 59, 192; 182, 1912; 64, 1936 179, '01, '02; 71, 1910, 1941 3038, 1941; 1864, 1902
bright sunshine .43.5 Last snow . March 23rd First snow . Nov. 3rd Last frost . May 14th First frost . Sept. 30th	52.8 April 15 Nov. 6 May 14 Sept. 21	Mar. 14 '10; May 11, '07, '4; Oct. 10, '25; Dec. 13, '41 Apr. 14, '43; June 8, '3; Aug. 22, '94, '95; Oct. 13, '09

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series Bulletin No. 757 January 1952

Meteorological Observations

FOR

JANUARY 1952

EARLE F. COX

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Te	Temperature				Wind			Precipi	tation			
	Max	imum	Min	inium		umidit	Iumidit F Hours							
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4	43 42 36 37	2 P 2 P 1 P	35 34 31 23	3 A 6 A 8 A 7 A	39 38 33.5 30	98 88 83 78	0.0 0.0 6.1 5.0	SE NW N SW	70 36 151 95	8 8 28 12	10 P 0 A	12 P 8 A	.06 .29	.6 2.9
5 6 7	$\frac{30}{31}$	4 P 2 P	$\frac{23}{28}$	8 P 11 P	$\frac{26.5}{29.5}$	99 91	$\frac{1.4}{3.5}$	N N	140 89	15 17	4 A	2 Р	.47	8
7 8 9 10 11 12	21 35 27 34 34 38	5 P 3 P 2 P 2 P 3 P 1 P	16 12 14 22 17 14	8 A 7 A 3 A 9 P 7 A 2 A	$ \begin{array}{c c} 18.5 \\ 23.5 \\ 20.5 \\ 28 \\ 25.5 \\ 26 \end{array} $	87 69 73 87 75 79	6.0 8.1 2.8 0.0 8.1 4.0	N E N W W N S W	262 245 4 86 243 33	20 13 3 28 17 4	8 A	12 a	.03	.6
13 14 15 16 17	45 38 41 42 40	2 P 3 P 8 P 2 P 12 P	23 32 33 35 17	8 A 1 A 8 A 1 A 8 A	$ \begin{array}{r} 34 \\ 35 \\ 37 \\ 38.5 \\ 28.5 \end{array} $	84 90 96 77 90	5.9 1.0 0.0 6.6 0.0	S N E N E	$\begin{array}{c c} & 6 \\ & 34 \\ & 24 \\ & 231 \\ & 49 \\ \end{array}$	5 3 28 28 28 32	9 A	2 P	.14	
18 19 20 21	45 39 42 38	6 A 2 P 2 P 0 A	33 24 29 17	8 P 8 A 0 A 12 P	39 31.5 35.5 27.5	74 68 90 69	5.0 6.4 0.0 7.6	N W S N W N	348 122 212 368	40 15 28 28	0 A 11 P 0 A	4 A 12 P 10 A	.23 .02 .45	0.3 1.4
22 23 24 25	30 42 30 28	12 P 10 A 0 A 12 P	10 28 16 12	5 A 12 P 12 P 7 A	$\begin{vmatrix} 20 \\ 35 \\ 23 \\ 20 \end{vmatrix}$	79 88 73 71	0.0 2.0 0.3 5.4	N N N W S W	78 73 202 68	10 20 15 13	5 P 0 A	12 P 4 A	,43 ,39	1.1
$\frac{26}{27}$	48 46	4 P	26 35	4 A 12 P	37 40.5	100	0.0	S N W	$\frac{139}{32}$	15 7	$\begin{cases} 2 & A \\ 5 & P \end{cases}$ $0 A$	{11 A 12 P 4 A	{.64 .27 .07	0.8
28	38	0 A	35	8 A	36.5	86	0.0	N W	69	14	0 A 0 A 3 P	6 A 9 P	.07 {.16 {.12	2.5
29 30 31	25 20 28	0 A 4 P 4 P	8 -7 -3	12 P 7 A 7 A	$\begin{array}{c} 16.5 \\ 6.5 \\ 12.5 \end{array}$	76 72 75	6.7 7.8 3.0	N W N W N W	191 56 18	17 8 6				

^{*}Based on least time required to blow one mile.

ROGER F. JOHNSON, Observer

MONTHLI	SUMIN	IARY
JANUARY, 1952	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum	30.70 29.20	31.00, 1927 28.55, 1913
Mean semi-daily 30.065 Range 1.259	30.07 1.50	2.18, 1913; 0.97, 1896
Air Temperature, in degrees F.		
Highest 48, 26th Lowest -7, 30th Mean 28.8 Range 55 Highest mean daily 40.5, 27th Lowest mean daily 6.5, 30th Mean maximum 35.9 Mean minimum 21.7 Greatest daily range 31, 31st Least daily range 3, 6th, 28th Degree-days of heating load for 11235	51.1 -7.1 24.2 58.2	66.0, 1932 —26.0, 1904 34.2, 1913; 13.9, 1918
month	1265	
Precipitation, in inches		
Precipitation 4.02	3.61 13.32	7.15, 1898; 1.07, 1896
Snow	13.32	33, 1898, 1923; 1.50, '08 17, '50, '20; 4, 1901
Wind, in miles	1	17, 30, 20, 4, 1901
Total movement	5055	7770, 1908; 2896, 1895
Mean hourly velocity	6.8 32.7	47, 1938
Wind, direction	i Mile # 1 s.	
Prevailing direction NNW North, days 9 Northeast, days 2 East, days 2 Southeast, days 1 South, days 3 Southwest, days 3 West, days 10 Northwest, days 10	WNW	
Weather Mean relative humidity, percent82.7	70.1	
Mean cloudiness, percent	55 9 9 13 137 46.7	78, 1932; 37, 1908 20, 1944; 1, 1949 18, 1890, 1926; 3, 1916, 1944 22, 1923, '31, '49; 4, 1920, '39 214, 1920; 50.4, 1949

Note—The first column in the above summary gives observations made during the month. The second column gives the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1951.

The mean daily temperature during the month of January was considerably above the normal, the reading being 28.8 degrees, as compared to a normal of 24.2 degrees. This resulted in a reduction in the degree days so that up to the middle of the heating season on January 22 the degree days accumulated were 3184.5, as compared to a normal of 3352.

The precipitation during the month was 4.02 inches; there was also 18.2 inches of snowfall. The normal of these two for the month is 3.61 and 13.32 inches, respectively. This excess of precipitation over the normal has continued for some time. During the month there were 14 days which had a daily total precipitation of 0.01 inches or more.

The wind movement during the month was considerably below normal. The total movement was 3774 miles as compared to a normal of 5055 miles. However, the winds during the month were not entirely low in velocity, as a recording of 40 miles per hour was taken on January 18th.

The mean relative humidity for the month was 82.7 per cent, as compared to a normal of 70.1 per cent. This is in line with readings which for over a year have continued to be 10 points above normal.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series Bulletin No. 758

February 1952

Meteorological Observations

FOR

FEBRUARY 1952

EARLE F. COX

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

> Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Te	mpera	ture		h	on.		Wind			Precipi	tation	
	Maximum		Minimum			Humidir	of Hour Inshine	£0	.	s *	20 LU X			
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max, Miles per Hour*	Began	Ended	Water	Snow
1	47	4 P	15	7 A	31	78	5.5	s E	88	9				
2	48	3 Р	31	12 P	39.5	86	2.2	N	116	12				
3	48	2 Р	25	7 A	36.5	88	7.3	s	72	14				
4	42	10 A	37	3 A	39.5	93	0.0	NE	109	10	$\begin{cases} 2 & A \\ 4 & P \end{cases}$	[7 A [11 P	{.22 .32	
5	44	2 P	33	9 P	38.5	86	1.8	N	73	20				
6	40	3 Р	29	12 P	34.5	69	5.7	N W	200	13				
7	34	1 Р	24	12 P	29	87	4.0	N	91	15	3 A	5 A	.03	0.6
8	40	12 M	21	4 A	30.5	76	7.0	S	77	15	11 P	12 P	.02	
9	37	12 M	25	12 P	31	75	6.7	N W	268	28	0 A	2 A	.09	
10	38	1 P	24	10 P	31	78	0.3	s w	129	17				
11	45	3 Р	24	4 A	34.5	87	4.2	N W	60	25	8 A	11 A	.28	
12	30	0 A	15	12 P	22.5	62	9.5	N W	461	-10		İ		
13	22	4 P	12	8 A	17	62	9.4	N W	3-14	20				
14	27	4 P	9	7 A	18	68	9.5	N W	287	25				
15	39	2 P	12	7 A	25.5	62	9.3	N.W	121	13				
16	38	2 P	13	7 A	25.5	67	6.1	N W	67	13				
17	36	2 P	26	12 P	31	96	0.0	N E	125	25	1 A	12 P	.61	8.0
18	35	3 P	25	12 P	30	100	0.0	N E	455	32	0 A	10 A	.31	2.7
19	34	4 P	18	12 P	26	88	6.8	N W	253	20				
20	39	4 P	10	6 а	24.5	88	9.7	N E	73	10				
21	37	2 P	23	12 P	30	98	3.6	N E	84	9	7 A	10 A	.06	0.4
22	37	12 M	24	6 A	30.5	93	0.3	N E	61	11		-		
23	35	12 M	11	12 P	23	81	6.2	N W	51	8				
24	38	4 P	8	7 A	23	86	9.5	N E	18	5			ļ	
25	40	3 Р	10	4 A	25	87	8.0	S W	20	8				
26	43	4 P	14	6 а	28.5	79	9.3	s w	30	5				
27	34	2 P	19	5 A	26.5	87	0.0	N W	114	20				
28	41	3 р	27	6 A	34	73	9.0	N W	285	20				
29	41	12 м	22	12 р	31.5	69	9.7	W	83	17				

^{*}Based on least time required to blow one mile.

MONTHLY	SUMM	IAKI
FEBRUARY, 1952	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum .30.277, 24th Minimum .29.433, 11th Mean semi-daily .29.908	30.63 29.24 30.04	31.05, 1920 28.56, 1895
Range	1.38	1.89, '00, '08; .88, '13, '31
Air Temperature, in degrees F.		
Highest .48, 2nd, 3rd Lowest .8, 24th Mean .29,2 Range .40 Highest mean daily .39.5, 2nd, 4th	50.5 -7.4 23.7 58	65.0, 1930 —23, 1943 32.6, 1890; 11.6, 1934
Lowest mean daily17, 13thMean maximum38.2Mean minimum20.2Greatest daily range32, 1stLeast daily range5, 4th	32.8 14.5	
Degree-days of heating load for month	1167 4927	
Precipitation, in inches		
Precipitation 1.97 Snow 11.7 Maximum precip. in 24 hrs	3.19 13.99	8.12, 1900; 0.62, 1901 48.75, 1893; 0.50, 1937
Number of days with .01 or more8	10	15, 1893, 1920; 4, 1901
Wind, in miles		
Total movement 4215 Greatest daily movement 461, 12th Least daily movement 18, 24th Mean hourly velocity 6.1	4776 7.1	6445, 1896; 3402, 1949
Maximum velocity	31.7	50, 1946
Wind, direction		
Prevailing direction NW North, days	WNW	
Weather		
Mean relative humidity, percent	66.6 50.9 10 8 10 158 53.4	66, 1890, 1927; 31, 1905 19, 1941; 2, 1927 16, 1920; 2, 1920 16, 1894; 2, 1920 221, 1924; 110, '27, '38
	t	

Note—The first column in the above summary lists observations made during the month. The second column lists the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1951.

The mean daily temperature during the month of February was 5.5 degrees above normal. The reading was 29.2 as compared to a normal of 23.7. The degree days for the month were 1037.5 as compared to a normal of 1167. The accumulative degree days for the season at the end of February were 4579.5 as compared to a normal of 4927. The accumulative degree days are slightly above 1950 and also above 1951, but the heating load has still been considerably below normal.

The precipitation for the month was 1.97 inches, which is far below the normal of 3.19 inches. This is the first month for several months that the precipitation has been below normal. The snowfall to the end of February was 42.6 inches, which is 14.45 inches above normal, but warm weather and a certain amount of rainfall have reduced the amount of snow on the ground so that at present there appears to be little danger of flood in the Connecticut Valley from an accumulation of snow. This statement, of course, must be qualified, according to the conditions that prevail in the Connecticut Valley above Amherst. The excess moisture in the soil which has accumulated over a period of several months because of the more than normal rainfall could result in a flood if sudden and heavy rainfall should occur. The soil has little ability, in its present condition, to absorb further moisture.

The mean relative humidity was far above normal for the month, as it has been for many previous months. The mean relative humidity for the month was 81.0 as compared to a normal of 66.6.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 759

March 1952

Meteorological Observations

FOR

MARCH 1952

EARLE F. COX

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Те	mpera	ture		r _z	ø		Wind			Precipi	tation	
	Maximum		Minimum			Humidir	of Hournshine	50		es *				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1	29	5 Р	12	12 P	20.5	83	1.5	N E	140	17	1 P	2 P	.02	.4
2	38	3 Р	3	6 A	20.5	73	10.2	N W	36	3				
3	37	4 P	12	4 A	24.5	72	7.0	N W	60	8				
4	40	3 Р	27	4 A	33.5	78	1.3	SE	51	11	8 P	12 P	.32	1.0
5	40	2 P	29	7 A	34.5	97	4.5	N E	266	17	0 A	10 A	.50	1.1
6	38	2 P	29	6 A	33.5	100	2.3	N E	197	12	7 A	10 A	.08	.60
7	38	3 р	33	· 6 A	35.5	100	0.0	N E	36	5	6 P	7 P	.01	
8	39	1 P	29	6 а	34.0	97	6.8	N W	52	8				
9	41	12 м	28	6 A	34.5	87	7.5	SE	51	5				
10	43	1 P	31	6 A	37	77	7.0	S E	107	13				
11	48	4 P	32	6 A	40	99	1.5	SE	93	32	3 A	7 P	1.37	
12	44	2 P	34	6 A	39	59	10.5	N W	456	25		!		
13	44	3 Р	26	7 A	35	71	8.3	N W	96	20			ļ	
14	39	2 P	27	6 A	33	61	11.0	N W	315	25				
15	35	3 р	21	6 A	28	59	10.3	N W	365	32				
16	29	11 P	20	8 A	24.5	69	9.5	N	285	20			į.	
17	44	3 Р	28	0 A	36	73	9.8	N W	296	25				
18	50	5 P	28	7 A	39	66	9.0	N W	144	9				
19	39	12 м	32	4 A	35.5	88	3.2	NE	147	14	1 P	12 P	.39	4.3
20	51	6 P	30	6 A	40.5	85	10.0	N W	100	7	0 A	2 A	.04	
21	54	5 P	27	6 A	40.5	95	10.2	N W	48	5	9 A	10 A	.03	
22	52	12 M	30	7 A	41	84	3.5	N W	59	7	6 A	7 A	.01	
23	40	2 P	35	8 A	37.5	99	0.0	N W	57	8	8 A	1 P	.40	
24	42	4 P	33	2 A	37.5	95	0.0	SE	93	6				
25	50	3 Р	27	6 A	38.5	66	10.9	S E	122	14				
26	66	3 Р	28	6 A	47	71	11.7	SE	133	12				
27	53	2 P	31	6 A	42	65	12.1	N W	141	12				
28	47	4 P	25	7 A	36	73	11.8	N W	120	14				
29	49	2 P	24	6 A	36.5	70	11.5	N E	91	11				
30	51	3 Р	24	6 A	37.5	62	11.5	ΝE	99	17				
31	55	4 P	31	6 A	43	74	11.8	SE	106	8]

^{*}Based on least time required to blow one mile.

WIOTITIET		
MARCH, 1952	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.381 Minimum 29.162 Mean semi-daily 29.818	30.57 29.24 30.04	31.05, 1943 28.47, 1914
Range 2.3016	1.34	2.10, 1914; 0.85, 1915
Air Temperature, in degrees F.		
Highest 66, 26th Lowest 3, 2nd Mean 35.3 Range 63 Highest mean daily 47, 26th Lowest mean daily 20.5 1st, 2nd Mean maximum 44.0 Mean minimum 26.6 Greatest daily range 38, 26th	63.8 6.1 34.4 57.6 43.2 25.4	85, 1929 -7.5, 1906
Least daily range	949 5876	
Precipitation, in inches		
Precipitation 3.17 Snow 7.4 Maximum precip. in 24 hrs. 1.37, 11th	3.70 7.47	7.89, 1942; 0.12, 1915 27, 1899; 0, 1921
Number of days with .01 or more11	11	17, 1890, 1913, 1936; 3, 1915
Wind, in miles		
Total movement	7.7 31.2	8182, 1896; 3006, 1905 48, 1932, 1939, 1941, 1942
Wind, direction		
Prevailing direction NW North, days 1 Northeast, days 7 East, days 0 Southeast, days 8 South, days 0 Southwest, days 0 West, days 0 Northwest, days 15	WNW	
Weather		
Mean relative humidity, percent	64.2 51.8 11 10 10 199 53.6	68, 1901; 27, 1915 22, 1924; 3, 1901 17, 1900; 1, 1943 21, 1901; 1, 1915 292, 1924; 93, 1901

Note—The first column in the above summary lists observations made during the month. The second column lists the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1951.

The mean daily temperature during the month of March was only slightly above normal with a reading of 35.3 compared to a normal of 34.4. The degree days of 919.5 for the month brought the accumulative degree days for the season to the end of March to a total of 5499. The normal for this period is 5876.

The precipitation for the month was 3.17 compared to a normal of 3.70. Even though this amount of rainfall was below normal, a slight increase in rainfall in the Connecticut Valley could have resulted in a flood inasmuch as the river was close to flood stage. The snowfall during the month was 7.4 inches compared to a normal of 7.47 inches. The total snowfall for the season to the end of March was 50.0 inches.

The mean relative humidity continued to be far above normal with a recording of 79.0 as compared to a normal of 64.2. The number of hours of bright sunshine, which may have an effect upon the 1952 crops, was considerably above normal with a recording of 61.2 compared to a normal of 53.6. This is far in excess of the recording in March of 1951 when a reading of 32.2 was recorded.

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 760

April 1952

Meteorological Observations

FOR

APRIL

1952

EARLE F. COX

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W.

Height of barometer above ground, 36 ft. Above sea level, 253.5 ft.

Height of wind instruments, 67 ft. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Te	mpera	ture		ž.	gr.		Wind			Precipi	tation	
	Maximum		Minimum			Tumidi	of Hourn	b0	د	87 *				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1	67	2 P	34	6 а	50.5	75	9.8	SE	106	17	11 Р	12 Р	.10	
2	48	6 Р	37	7 A	42.5	96	11.3	ΝE	106	7	0 а	δа	.32	
3	56	2 P	37	6 а	46.5	74	12.5	N W	114	14	5 A	6 A	.09	
4	63	4 P	34	6 а	48.5	66	8.0	N W	102	14				
5	52	6 P	35	8 A	43.5	99	0.0	N E	95	13	{1P 7P	{2 P 9 P	.22	
6	52	12 м	41	8 A	46.5	75	4.7	s w	185	17	(,,	(
7	47	2 Р	36	7 A	41.5	79	4.2	w	109	11				
8	54	1 P	36	6 A	45	75	7.4	N W	80	10				
9	56	6 P	38	6 A	47	71	10.3	s w	66	6				
0	63	4 P	36	4 A	49.5	74	8.3	S	164	20				
1	48	5 P	35	8 A	41.5	61	12.2	N W	295	20				
12	54	3 Р	27	6 а	40.5	62	12.5	s w	46	5				
13	51	4 P	39	6 A	45	65	8.4	N E	71	9				
14	49	3 Р	39	6 а	44	99	0.0	N E	107	13	1 A	6 A	.58	
15	51	4 P	41	4 A	46	75	2.5	N E	215	20				
16	66	5 P	32	6 A	49	55	12.5	N E	153	14				
7	73	5 P	30	5 A	51.5	57	12.1	N W	75	11				
8	72	4 P	44	7 A	58	55	12.1	N W	171	17			,	
9	81	2 P	38	6 а	59.5	65	11.3	s w	106	14				
0:	82	1 P	60	4 A	71	54	12.0	N W	310	25				
21	71	3 Р	42	6 а	56.5	58	12.2	N	136	14				
22	79	4 P	34	6 A	56.5	67	9.2	SE	82	14				
23	72	9 A	54	3 A	63	71	4.8	N W	128	20				
24	67	4 P	40	5 A	53.5	53	9.8	S E	98	7				
25	53	11 A	43	6 а	48	93	0.0	ΝE	52	5	10 A	12 P	.70	
26	59	3 P	42	0 а	50.5	93	4.8	N W	35	4	∫0a 4p	{6A 5₽	.20	
27	71	3 Р	48	6 A	59.5	74	7.9	S E	74	11		(33		
28	58	1 P	49	6 а	53.5	99	0.0	ΝE	165	13	0 л	7 P	1.19	
29	70	6 Р	48	3 A	59	86	9.5	N E	139	10				
30	67	4 P	45	6 а	56	53	7.5	ΝE	203	25				

^{*}Based on least time required to blow one mile.

Normal	Extremes
30.48 29.38	30.71, 1911 28.98, 1943
1.11	1.52, 1930; 0.65, 1950
79.4 22.0 45.7 57.4	90, 1941 8.5, 1923 52, 1921; 41, 1943
56.8 34.8	
579 6455	
3.35	6.89, 1929 ; .55, 1941 11, 1891 ; 0, 1910, 1934
11	18, 1909; 3, 1892
5404	8208, 1908 ; 3853, 1917
7.5 31.4	40, 1935, 1938, 1940, 1945
WNW	
61.6 51.8 11 9 10 220 54.7	75, 1901; 34, 1925, 1927 23, 1941; 1, 1949 18, 1915; 2, 1894, 1901 22, 1901; 1, 1941 329, 1941; 103, 1901
	30.48 29.38 29.99 1.11 79.4 22.0 45.7 57.4 56.8 34.8 579 6455 3.35 2.17 11 5404 7.5 31.4 WNW

Note—The first column in the above summary lists observations made during the month. The second column lists the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1951.

The mean daily temperature for the month of April was well above normal with a reading of 50.8 as compared to a normal of 45.7. This resulted in less degree days of heating load for the month than is normal. The reading for this was 433 as compared to a normal of 579. The accumulative degree days for the season were 5932 as compared to a normal of 6455.

The precipitation for the month was 3.40 as compared to a normal of 3.35. A great increase in this normal amount of rainfall could have had serious consequences in the Connecticut Valley because the river was nearly at flood stage for the greater part of the month.

The mean relative humidity was much lower than for the month of March but it was still considerably above normal for the month of April. The reading was 72.6 as compared to a normal of 61.6. The increase in the percentage of possible hours of bright sunshine begun in March continued through April. This reading was 59.4 as compared to a normal of 54.7.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series Bulletin No. 761

May, 1952

Meteorological Observations

FOR

MAY

1952

EARLE F. COX

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W.

Height of barometer above ground, 36 ft. Above sea level, 253.5 ft.

Height of wind instruments, 67 ft. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Te	mpera	ture		ty	ga		Wind		Precipitation				
	Maxi	inum	Mini	imum		midi	Hour nine						1		
Dav	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow	
1	64	4 P	45	7 A	54.5	54	0.0	N E	159	20					
$\stackrel{\cdot}{2}$	65	6 Р	34	5 л	49.5	67	0.8	N E	118	14					
3	60	5 P	32	6 а	46.0	62	13.2	N W	157	17					
4	67	4 P	40	6 а	53.5	59	12.8	NW	208	25					
5	68	7 P	33	6 а	50.5	66	12.5	s w	78	9					
6	60	6 Р	36	6 A	48.0	84	5.5	s	100	17	6 Р	7 P	.05		
7	60	4 P	47	6 A	53.5	66	10.3	N W	288	20					
8	70	3 Р	35	6 A	52.5	54	11.9	N W	174	17					
9	73	5 Р	43	6 A	58.0	60	12.8	N W	117	12					
10	80	3 Р	40	7 A	0.00	59	9.0	SE	64	6					
11	62	0 A	49	9 A	55.5	100	0.0	ΝE	89	8	6 A	12 P	.67		
12	65	12 м	45	4 A	55.0	80	4.2	s w	229	25	} 0a }12m	{6A 1P	{1.01 {.06		
13	60	11 A	46	6 A	53.0	71	3.1	s w	191	14		-			
14	67	6 P	46	6 A	56.5	64	7.4	N W	240	20		1			
15	54	6 Р	44	5 A	49.0	96	2.0	N E	49	9	6 A	3 Р	.26		
16	62	12 M	47	7 A	54.5	58	13.3	N W	186	25					
17	65	4 P	32	7 A	48.5	65	9,9	N W	10	5	f 0A	(8A	\(\) .24		
18	68	6 Р	47	6 л	57.5	88	8.9	N W	56	10	{ 0A { 11A	12M	1.01		
19	73	2 P	39	5 A	56.0	66	9.3	S	18	8					
20	60	12 M	52	6 A	56.0	95	0.0	SE	52	5	1 P ∫1A	12 P	.44 \$.06		
21	65	3 P	49	5 A	57.0	88	4.6	W	53	13	{5P	{6P	{.04		
22	68	3 P	51	5 A	59.5	72	10.9	N W	223	17					
23	76	4 P	43	4 A	59.5	67	10.3	N	94	11				ļ	
24	80	3 P	45	5 A	62.5	69	8.0	s w	29	5		10-	.84		
25	59	3 P	49	8 P	54.0	100	0.0	SE	102	$\begin{vmatrix} 10 \\ 7 \end{vmatrix}$	4 A	12 Р	.84		
26	69	6 P	48	0 A	58.5	94	7.3 13.7	N	65 158	20					
27	77	6 P	49	6 A	63.0	59 72		N	$\frac{158}{42}$	8					
28	82	4 P	41	6 A	61.5	85	9.8 6.0	NW	84	11	{12M 7₽	} 2P }10P	5.09		
29	80 73	1 P	58	6 A 7 A	63.0	71	10.1	S W	58	5	1 7P	(10p	1.05		
30 31	72	5 Р 6 Р	53 55	7 A	63.5	97	3.3	SE	69	$\frac{3}{9}$	{ 3A	∫ 9A	§.12		
91	'-	l or	3.3	I A	00.0	''	5.5		"		\10P	13P	1.06	<u> </u>	

^{*}Based on least time required to blow one mile.

MONTILL		
MAY 1952	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.056, 19th Minimum 29,200, 12th Mean semi-daily 29,903 Range 0.856	30.39 29.51 29.96 0.88	30.62, 1936 29.10, 1938
Air Temperature, in degrees F.		
Highest 82, 28th Lowest 32, 3rd, 17th Mean 56.1 Range 50 Highest mean daily .69, 29th	86.1 31.2 57.1 54.9	94.5, 1896, 1911 24.0, 1900
Lowest mean daily 46, 3rd Mean maximum 67.9 Mean minimum 44,3 Greatest daily range 41, 28th Least daily range 8, 20th Degree-days of heating load for month 280.5	56.8 34.8	
Degree-days cumulative for season6212.5		
Precipitation, in inches Precipitation	3.60	7.44, 1931; .48, 1903
Maximum precip. in 24 hrs1.07, 12th Number of days with .01 or more10	12	20, 1901, 1945; 5, 1903
Wind, in miles Total movement	4504	5946, 1907; 2180, 1894
Least daily movement10, 17thMean hourly velocity4.8Maximum velocity25, 4th, 12th, 16th	6.1 28.8	45, 1935
Wind, direction		
Prevailing direction NW North, days 3 Northeast, days 4 East, days 0 Southeast, days 4 South, days 2 Southwest, days 5 West, days 1 Northwest, days 12	W	
Weather		
Mean relative humidity, percent	60.7 52.1 11 11 9 252	72, 1948; 30, 1923 21, 1944; 0, 1927 18, 1949; 5, 1923 20, 1927; 2, 1941 350, 1944; 137, 1927
Number of hours bright sunshine230,9		20, 1927; 2, 19- 350, 1944; 137, 192

Note—The first column in the above summary lists observations made during the month. The second column lists the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column lists extremes observed from 1889 to 1951.

REMARKS

The weather during the month of May was unusual because the mean daily temperature was slightly below normal; the mean daily temperature was 56.1 compared to a normal of 57.1. This resulted in having the degree days for the month above normal. This reading was 280.5 and brought the accumulative degree days for the season to 6212.5.

Rainfall occurred chiefly on weekends. The total rainfall for the month was 4.00 inches as compared to a normal of 3.60 inches.

The mean relative humidity continued to be far above normal with a reading of 74.0 as compared to a normal of 60.7. The mean cloudiness was near normal, and the percent of possible hours of sunshine with the reading of 51.1 was slightly below the normal of 55.6.

Meteorological Series Bulletin No. 762

June, 1952

Meteorological Observations

FOR

JUNE 1952

ROBERT K. PATTERSON

OBSERVATORY

Latitude, 42° 23′ 31″ N.

Longitude, 72° 31′ 48" W.

Height of barometer above ground, 36 ft. Above sea level, 253.5 ft.

Height of wind instruments, 67 ft. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Te	mpera	ture		rt.	ø		Wind			Precipi	tation	
	Max	imum	Min	imum		lumidi	f Hours shine		43					
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1	68	2 P	60	5 а	64.0	99	0.0	s	99	20	0 a	10 р	2.70	
2	79	4 P	58	6 а	68.5	68	8.1	N	159	20	8 A	9 a	0.03	
3	83	5 P	48	6 а	65.5	69	8.7	s w	65	9				
4	86	1 P	56	6 A	71.0	70	6.5	s	98	14	11 P	12 p	0.09	
5	85	5 P	56	7 A	70.5	79	9.8	w	69	12				
6	81	12	56	7 A	68.5	81	9.1	s w	97	13	3 р	4 P	0.01	
7	79	6 P	58	3 A	68.5	60	10.7	N	297	28				
8	83	6 P	51	6 A	67.0	58	10.1	\mathbf{w}	115	14				
9	81	1 P	62	6 A	71.5	62	3.7	N W	122	15				
10	83	5 P	51	6 A	67.0	64	10.9	w	152	17	ļ			
11	68	12	57	6 A	62.5	68	6.5	N W	166	25				
12	74	5 P	57	0 A	65.5	69	9.4	N	257	20		ŀ		
13	81	4 Р	50	5 A	65.5	64	12.3	N	67	10		į .		
14	83	2 P	50	4 A	66.5	65	12.6	s	91	11	l			
15	88	5 P	51	5 A	69.5	72	10.3	N W	168	17	4 A	5 A	0.01	1
16	84	4 P	58	6 A	71.0	58	11.2	w	115	13				
17	87	4 P	60	4 A	73.5	85	8.6	s w	107	10	4 A	8 P	0.86	1
18	81	1 P	59	5 A	70.0	60	11.7	N W	202	25	1		1	
19	80	2 P	5 3	5 A	66.5	66	8.2	N W	175	32				
20	72	3 Р	52	4 A	62.0	58	12.3	N W	253	25				
21	72	5 P	52	5 A	62.0	65	8.7	s w	69	7				
22	66	1 P	58	6 A	62.0	82	1.9	w	49	5				
23	79	3 Р	48	4 A	63.5	72	8.8	s w	93	10				
24	88	5 P	55	6 A	71.5	76	8.7	s w	73	17		4	0.40	
25	91	5 P	66	6 P	78.5	76	10.8	N	127	14	1 A	4 A	0.49	1
26	98	5 P	71	5 A	84.5	74	11.7	N W	117	10	9 Р	10 P	0.26	
27	88	3 P	68	6 A	78.0	74	11.8	NE	83 98	9 7				
28	83	4 P	65	6 A	74.0	$\begin{array}{ c c } 76 \\ 92 \end{array}$	11.6	NE	102	9	10 A	2 P	0.52	
29	77	8 A 3 P	65 58	5 A	69.0 67.5	68	0.6 10.3	s	102	11	10 A	2 P	0.02	
30	''	5 P	00	6 A	07.5	00	10.3	NE	143	11				

^{*}Based on least time required to blow one mile.

137 11	
Normal	Extremes
30.30 29.55 29.95 0.75	30.70, 1947 29.24, 1902
91.2 40.0 65.7 51.3	101, 1919 34, 1891
3.75	9.68, 1922; 0.76, 1908
11	17, 1922, 1945; 4, 1908
	4571, 1908; 1409, 1906 48, 1939
	,
51.1 10 12 8 257 54.0	71, 1903; 28, 1908 22, 1908, 1941, 1943; 0, 1949 23, 1912; 3, 1941 22, 1903; 1, 1923 365, 1947; 102, 1903
	29.55 29.95 0.75 91.2 40.0 65.7 51.3 3.75 11 3585 5.0 24.6 WSW

Note—The first column in the above summary lists observations made during the month. The second column lists the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column lists extremes observed from 1889 to 1951.

REMARKS

Total precipitation for the month was 4.97 inches of which 2.70 inches occurred on the first day. The storm starting May 31 accumulated 2.76 inches of rain in less than 12 hours.

Good weather prevailed throughout most of the month with only four cloudy days. Bright sunshine occurred 58.1 percent of the possible hours, the normal for the month being 54 percent.

As in the preceding month, the mean relative humidity remained high. A high temperature of 98° occurred on the 26th with a mean of 84.5° for the day.

The total wind movement was high with 3834 miles of total movement compared to a normal of 3,585.

Meteorological Series

Bulletin No. 763

July 1952

Meteorological Observations

FOR

JULY

1952

ROBERT K. PATTERSON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Te	mpera	ture		ty.	ø		Wind			Precipi	tation	
	Max	imum	Min	imum		Humidi	f Hour nshine		4	82 *				
Dav	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1	83	5 P	50	5 A	66.5	62	12.1	N	65	9				
2	86	2 P	51	5 A	68.5	61	11.2	W.	57	8				
3	90	3 р	58	5 а	74.0	70	11.1	s w	134	25				
4	89	1 P	68	2 A	78.5	81	9.2	s	122	17	4 P	6 P	0.73	
5	86	7 P	69	7 A	77.5	72	11.1	N	72	20				
6	91	4 P	58	6 а	74.5	66	11.6	W.	38	6				
7	91	3 Р	60	4 A	75.5	67	11.4	W	44	11				
8	86	4 P	62	7 A	74.0	70	8.7	s w	106	11				
9	78	2 P	68	4 A	73.0	74		S E	131	20				
10	69	1 P	66	6 A	67.5	93		N E	124	32	7 A	5 P	2.01	
11	85	7 P	64	6 A	74.5	73	8.7	N	88	7	6 а	7 A	0.03	
12	90	5 P	59	7 A	74.5	81	10.8	s w	33	4				
13	91	3 Р	55	5 A	73.0	70	11.3	N W	44	5				
14	95	4 P	62	6 A	78.5	73	11.7	s	66	9				
15	92	2 P	70	6 A	81.0	70	11.1	s	154	11				
16	89	3 Р	72	5 A	80.5	-66	10.9	N W	144	11				
17	88	6Р	64	6 A	76.0	66	11.8	s	66	6				
18	93	2 P	66	6 A	79.5	69	10.6	W	37	14				
19	88	4 P	73	6 A	80.5	70	9.2	W	157	25	0 A	2 A	0.03	
20	90	2 P	71	5 A	80.5	66	10.1	W	106	14				
21	82	6 Р	68	5 а	75.0	82	5.7	SE	39	4				
22	93	6 P	72	6 A	82.5	75	10.4	s w	137	17				
23	95	4 P	64	5 A	79.5	67	11.0	SE	94	9				
24	82	4 P	71	7 A	76.5	60	10.4	W	128	14				
25	79	6 Р	52	4 A	65.5	65	11.7	s	113	7				
26	88	5 Р	51	6 а	69.5	65	11.5	SE	111	10				
27	87	4 P	66	6 а	76.5	87	8.3	s	116	32	8 a	5 P	1.21	
28	87	2 Р	59	5 A	73.0	82	5.1	s	40	6				
2 9 .	86	2 P	68	6 A	77.0	75	մ.4	N W	129	20				
30	79	4 P	53	6 а	66.0	70	7.0	s	90	5				
31	85	1 Р	62	3 A	73.5	88	3.2	N W	74	25	1 P	G P	1.01	

^{*}Based on least time required to blow one mile.

JULY 1952	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)	20.27	
Maximum 30.353, 1st Minimum 29.750, 19th Mean semi-daily 30.062	30.27 29.59 29.96	30,50, 1892 29,27, 1932
Range	0.68	0.97, 1892; 0.47, 1938
Air Temperature, in degrees F.		
Highest .95, 14th, 23rd Lowest .50, 1st Mean .75 Range .45 Highest mean daily .82,5, 22nd Lowest mean daily .65,5, 25th Mean maximum .86,9 Mean minimum .63 Greatest daily range .37, 26th Least daily range .3, 10th Degree-days of heating load for month Degree-days cumulative for season	93.9 46.4 70.8 47.7	104, 1911 40, 1890, 1898 66.3, 1891; 76.0, 1949
Precipitation, in inches		
Precipitation	4.10	14.51, 1897; 0.70, 1929
Maximum precip. in 24 hrs2.01, 10th Number of days with .01 or more5	11	20, 1915; 4, 1924
Wind, in miles		
Total movement	3422 4.6 28.7	5097, 1909; 1109, 1894
Maximum velocity32, 10th	۵۰.7	44, 1936
Wind, direction Prevailing direction S North, days 3 Northeast, days 1 East, days - Southeast, days 4 South, days 8 Sonthwest, days 4 West, days 7 Northwest, days 4	SW	
Weather		
Mean relative humidity, percent	68.4 50.9 10 14 7 268	70, 1943; 31, 1924 22, 1923; 0, 1915 24, 1909, 1921; 4, 1946 18, 1889; 0, 1910 371, 1910; 180, 1931

Note—The first column in the above summary lists observations made during the month. The second column lists the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column gives extremes observed from 1889 to 1951.

WEATHER SUMMARY

The highest temperature for the month was 95°which occurred on the 14th and 23rd, and the lowest was 50°F, on the first. The mean temperature was 75°, the normal for the month being 70.8°. The highest mean daily temperature was 82.5°F, and the lowest mean daily 65.5°F. The greatest daily temperature was 37°F, on the 26th, and the least daily range was 3°F, on the tenth.

The rainfall was slightly above normal for the month with 4.99 inches of precipitation, the normal for the month being 4.10 inches. Of the 4.99 inches of rainfall for the month, 2.01 inches fell in an eleven-hour period on the 10th.

A wind velocity of 32 MPH occurred on the 10th, and the total wind movement for the month was 2919 miles. The mean relative humidity of 72 was above the normal of 68.4. The per cent of possible hours of bright sunshine recorded 61.3 was above the normal of 58. The number of clear days for the month was 16, the normal being 10. Five thunder and lightning storms occurred during the month.

Meteorological Series

Bulletin No. 764

August 1952

Meteorological Observations

FOR

AUGUST 1952

ROBERT K. PATTERSON

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W.

Height of barometer above ground, 36 ft. Above sea level, 253.5 ft.

Height of wind instruments, 67 ft. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Те	mpera	ture		ž.			Wind			Precipi	tation	
	Max	imum	Min	imum		umídií	Houn shine							
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Ведяп	Ended	Water	Snow
1	82	5 р	57	4 A	69.5	75.7	5.1	W	42	8				
2	75	4 P	56	5 а	65.5	86.0	0	SE	38	6				
3	86	3 р	62	6 а	74.0	76.0	7.9	s w	34	6				
4	86	3 р	62	5 A	74.0	82.5	4.5	SE	92	17				
5	87	2 P	64	0 л	75.5	86.3	1.1	SE	149	10	2 P	3 Р	0.12	
6	75	3 р	56	6 A	65.5	98.0	0	N W	57	5	6 A	8 Р	0.13	
7	82	3 Р	65	6 а	73.5	80.0	5.5	s	38	8	0 л	1 A	0.03	
8	77	3 Р	59	6 а	68.0	83.7	0	E	25	5				
9	82	4 P	52	3 A	67.0	76.3	6.2	E	57	17				
10	70	8 P	62	9 A	66.0	96.7	0	N W	49	6	4 A	5 P	0.57	
11	86	4 P	58	3 A	72.0	82.3	7.7	w	49	7	0 а	1 A	0.03	
12	79	3 Р	54	5 A	66.5	94.3	0	w	38	5	6 A	12 P	0.80	i
13	84	4 P	55	5 а	69.5	81.0	7.6	w	71	17	0 а	5 A	0.96	
14	79	6 Р	56	6 A	67.5	80.3	7.4	s w	58	14				
15	86	4 P	54	6 A	70.0	78.7	6.7	SE	64	20				
16	76	1 P	52	0 л	64.0	90.7	0	SE	144	17	1 P	8 P	0.38	
17	84	2 P	69	5 A	76.5	81.0	8.1	w	201	18	0 л	4 A	0.74	Ì
18	80	4 P	58	7 A	69.0	75.3	9.1	w	93	11				
19	79	5 P	53	6 A	66.0	73.3	8.2	N W	52	10				
20	81	4 P	49	6 A	65.0	74.0	7.8	s	28	5			l	
21	76	1 P	54	6 A	65.0	84.0	1.3	SE	98	10	9 P	11 P	0.21	
22	78	1 P	60	3 A	69.0	77.7	6.1	w	159	25				
23	72	5 P	45	5 A	58.5	63.0	6.4	w	148	20				
24	75	5 P	43	7 A	59.0	69.3	6.4	N W	133	18				
25	81	3 P	46	6 A	63.5	71.6	4.1	W.	43	8				
26	89	4 P	52	6 A	70.5	78.6	1.7	5	37	5				
27	91	4 P	60	6 A	75.5	78.6	4.7	s	34	5				
28	86	3 P	64	7 A	75.0	77.7	2.6	s	73	6				
29	89	3 P	68	3 A	78.5	80.3	5.6	s w	124	11				
30	77	12 N	70	6 A	73.5	81.0	0.3	Е	113	13	6 A	7 A	0.01	
31	71	1 P	59	6 A	65.0	77,3	0	Е	57	8				1

^{*}Based on least time required to blow one mile.

Normal	Extremes
30.32 29.61 30.00 0.71	30.50, 1934 28.87, 1936
91.6 43.4 68.6 48.1	100, 1918, 1948 34, 1940 62.4, 1903
4.08	8.40, 1948; 0.51, 1894
11	16, 1892, 1933 ; 4, 1899
3127 4.2 22.7	4,217, 1910; 1,920, 189 ² 40, 194)
	,
237	67, 1901, 1946; 27, 192, 23, 1941; 0, 1915, 192 25, 1912; 3, 193; 18, 1901, 1928; 2, 1910, 192, 332, 1941; 152, 1915, 192
	30.32 29.61 30.00 0.71 91.6 43.4 68.6 48.1 4.08 11 3127 4.2 22.7 SW

Note—The first column in the above summary lists observations made during the month. The second column lists the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column lists extremes observed from 1889 to 1951.

REMARKS

The weather for August was normal. The mean temperature was 68.95° with a precipitation of 3.98 inches. Although the weather conditions were normal for the month, the serious summer drought experienced by eastern and southern states carried into the early part of August and was ended by heavy rains, wind storms, and hail between the 10th and the 17th of the month. The storms were localized and did a great deal of damage to the farm crops and buildings in the areas affected.

Meteorological Series

Bulletin No. 765

September 1952

Meteorological Observations

FOR

SEPTEMBER

1952

I. J. PFLUG

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Te	mpera	ture		ίν	S		Wind			Precipita	ation	
	Max	imum	Min	imum		Iumidi	f Hour nshine		t	9				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 1 2 3 4 5 1 6 7 8 8 9 10 11 12 13 11 14 15 16 17 18 19 20 21 1 22 2 3 2 4 2 5 5 2 6 7 2 8	73 88 87 74 76 81 84 68 68 73 84 68 73 87 72 61 77 77 74 77 77 77 68 77 77 77 77 77 77 77 77 77 77 77 77 77	8 P 4 P 4 P 4 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 4 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2	61 63 69 49 49 52 52 52 43 43 47 54 60 62 60 62 60 46 45 55 55 48 48 50 48 49 49 49 49 49 49 49 49 49 49 49 49 49	6 A 6 A 7 A 7 A 7 A 7 A 7 A 6 A A 4 A 11 A 8 A A 7 A A 7 A A 7 A A 6 A A 7 A A A 6 A A 7 A A 7 A A A 7 A A A 7 A A A 7 A A A 7 A A 7 A A 7 A A 7 A A 7 A A 7 A A 7 A A 7 A A 7 A A A 7 A A A 7 A A A 7 A A A 7 A A A 7 A A A 7 A A A A 7 A A A A A 7 A	67.0 75.5 71.5 62.5 65.0 68.0 68.0 65.5 58.0 65.5 71.5 66.0 58.0 61.5 71.5 66.5 59.0 56.5 54.0 59.5 63.0 59.5 63.0 59.5 63.0 65.5	93.3 79.7 85.0 76.3 78.7 76.0 69.3 78.0 79.0 82.3 79.3 77.0 76.3 84.7 93.3 94.7 82.3 94.7 88.0 83.3 79.7 88.0		E SE SE NW W W N NW SW SW NW NW NW NW	90 149 90 37 86 192 68 78 33 78 88 91 76 61 63 146 70 34 57 72 33 43 126 88 83 83 83 84 86 87 87 88 88 89 80 80 80 80 80 80 80 80 80 80	17 12 20 14 6 11 11 14 5 3 9 7 7 12 12 11 13 14 6 5 7 7 12 12 13 14 14 15 16 17 18 18 18 18 18 18 18 18 18 18 18 18 18	2P 11A 6 P 4 A 10 A	10 P 12 N 10 P 6 P 12 N 7 P	1.43 0.02 10 2.13 0.12 0.25	
29 30	80 75	3 P 1 P	42 46	7 A 7 A	61.0	79.3 81.0	2.4	S NE	40 76	5 14				

^{*} Based on least time required to blow one mile.

HANS JOA, Observer

Normal	Extremes
30.45 29.57 30.00	30.65, 192- 28.41, 193
.88	1.99, 1938; 57, 191
	97, 1895, 192 24.5, 191
4.24	14.55, 1938; 0.52, 191
10	16, 1933; 3, 190
3271 4.5 25.9	4.680, 1896; 1414, 189 80, 193
73.6 49.9 10 10	70, 1934; 23, 194 19, 1932; 2, 1907, 192 19, 1908, 3, 188 20, 1934; 3, 1941, 194 255, 1916; 106, 193
54.9	
	30.45 29.57 30.00 .88 87.7 33.2 61.7 54.5 4.24 10 3271 4.5 25.9 WSW

Note—The first column in the above summary lists observations made during the month. The second column lists the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column lists extremes observed from 1889 to 1951.

BEMARKS

The mean daily temperature for the month of September was 1.5 degrees above normal and the mean relative humidity 8.6 percent above normal. The number of hours of bright sunshine was 84 compared to a normal of 201. All these factors made the month of September seem unusually warm and in many cases uncomfortable. There was no frost during the month of September.

Meteorological Series

Bulletin No. 766

October 1952

Meteorological Observations

FOR

OCTOBER

1952

I. J. PFLUG

OBSERVATORY

Latitude, 42° 23′ 31″ N. Longitude, 72° 31′ 48″ W Height of barometer above ground, 36 ft. Above sea level, 253.5 ft. Height of wind instruments, 67 ft. Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Te	mpera	ature		ž.			Wind	1		Precipi	tation	
	Max	imum	Mir	imum		fumidi	f Hour nshine			9				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8 9 10	73 71 53 64 73 76 65 59 61 59	4 P 4 P 2 P 4 P 4 P 3 P 3 P 3 P 2 P 4 P	40 55 48 38 45 56 58 33 32 38	7 A 1 A 8 A 7 A 7 A 8 A 7 A 8 A	56.5 63.0 50.5 51.0 59.0 66.0 61.5 46.0 46.5 48.5	80.7 93.3 85.0 72.0 77.3 79.3 88.3 75.3 79.3 71.3		N SE NW SW SE S NW NW NW	54 102 160 139 122 80 120 63 14	5 25 20 10 13 5 20 14 4 14	6 P 0 A	12 P 8 A	0.50 0.18 0.39	
11 12 13 14 15 16 17 18	57 64 70 71 67 64 60	2 P 4 P 4 P 4 P 3 P 4 P 2 P	31 31 38 53 37 49 33 35	8 A 8 A 6 A 8 A 8 A 8 A 7 A	44.0 47.5 54.0 62.0 52.0 56.5 46.5 47.0	78.0 71.7 74.3 67.0 80.0 68.3 68.0 70.3		E E S W N W	62 - 83 118 - - 66 40	8 5 10 20 - 25 13				
19 20 21 22 23 24 25 26 27 28 29 30	59 48 53 64 65 65 53 56 65 58 42 52	2 P 2 P 3 P 3 P 4 P 2 P 3 P 3 P 3 P 1 P 0 P 1 P	45 28 27 34 31 44 38 21 40 54 32 32	7 A 7 A 6 A 7 A 7 A 8 A 8 A 7 A 2 A 1 A 6 A	52.0 38.0 40.0 49.0 48.0 54.5 45.5 38.5 52.5 56.0 37.0 42.0	61.3 63.7 65.0 61.3 57.7 57.7 61.0 70.7 73.0 79.7 64.7 59.0		W NW NW SS S W NW SE SE SE NW	118 174 163 84 90 194 215 61 193 211 242 139	25 25 20 10 9 32 32 8 14 20 25				

^{*} Based on least time required to blow one mile.

Hans Joa, Observer

[†] Sunshine Indicator not working.

1201(11121	20111	
OCTOBER 1952	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum .30.375, 26th Minimum .29.614 28th, 29th Mean semi-daily .30.024	30.53 29.42 30.06	30.68, 1929 29.00, 1926
Range0.761	1.11	1.47, 1926; 0.76, 1899
Air Temperature, in Degrees F.		
Highest 76.0, 6th Lowest 21.0, 26th Mean 50.6 Range 55.0 Highest mean daily 66.0, 6th Lowest mean daily 37.0, 29th Mean maximum 61.7 Mean minimum 39.2 Greatest daily range 35.0, 26th Least daily range 40, 28th Degree-days of heating load for month 446 Degree-days cumulative for season 557	79.4 23.4 50.5 56.0	90.5, 1908 17, 1936 57.4, 1947; 43.2, 1890
Precipitation, in inches		
Precipitation	3.29	8.81, 1911; 0.01, 1924
Snow — Maximum precip. in 24 hrs 0.50 Number of days with .01 or more	9	15, 1913; 1, 1897, 1924
Wind, in miles		
Total movement. 3344 28 days Greatest daily movement 242, 29th Least daily movement 14, 9th Mean hourly velocity 4.98 Maximum velocity 32, mph. 24th	4074 5.4 29.5	5467, 1910; 2540, 1894
	29.5	42, 1937
Wind, direction		
Prevailing direction NW North, days 3 Northeast, days - East, days 2 Southeast, days 6 South, days 4 Southwest, days 3 West, days 5 Northwest, days 8	11.	
Weather		
Mean relative humidity, percent 71.0 Mean cloudiness, percent 31.9 Number of clear days 19 Number of fair days 5 Number of cloudy days 7 Number of hours bright sunshine - Percent of possible hours of bright sunshine - Thunder and lightning 2nd First Frost 11th	69.0 48 10 10 11 175.0 51.3	67, 1932; 18, 1924 21, 1938; 1, 1911 17, 1924; 3, 1938 19, 1896, 3, 1924 232, 1923, 1938; 91, 1913

Note—The first column in the above summary lists observations made during the month. The second column lists the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column lists extremes observed from 1889 to 1951.

REMARKS

The outstanding feature of the October weather was the small amount of rainfall, all of which occurred in the first seven days of the month.

The lack of rainfall created very dry woodland conditions, resulting in many forest fires.

The first general frost occurred on October 11. Temperatures for the month were normal; the number of degree days for the month was 446, and the total accumulative degree days for the heating season was 557.

Meteorological Series

Bulletin No. 767

November 1952

Meteorological Observations

FOR

NOVEMBER

1952

L. J. PFLUG

OBSERVATORY

Latitude, 42° 23′ 31″ N.

Longitude, 72° 31′ 48″ W.

Height of barometer above ground, 36 ft. Above sea level, 253.5 ft.

Height of wind instruments, 67 ft.

Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Te	mpera	ature		<u> </u>	, m		Wind	l 		Precipit	ation	
	Max	kimum	Mir	nimum	imum		of Hour inshine	F0	ļ,	es.				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5	69 68 54 48 51	3 P 3 P 2 P 3 P 2 P	41 33 46 26 20	7 A 7 A 12 P 12 P 7 A	55.0 50.5 50.0 37.0 35.5	43.7 63.3 75.0 57.0 66.3	7.0 5.9 0.0 7.7 6.8	W W S NW	142 54 95 163 64	17 8 14 17 8	1 P	3 р	0.02	
6 7 8 9 10	57 44 44 32 46 43	1 P 2 P 12 M 4 P 3 P 4 P	40 34 24 18 30 26	3 A 12 P 12 P 6 A 12 P 4 A	48.5 39.0 34.0 25.0 38.0 34.5	62.3 63.7 89.7 100.0 83.7 75.3	5.2 1.6 5.8 0.4 2.4 4.1	W W NW NW NW	173 194 120 3 29 7	25 25 28 1 15 4	1 P	7 р	0.13 1.0	0
12 13 14 15 16 17 18	43 47 52 47 56 50 49	11 A 3 P 4 P 8 P 3 P 0 A 3 P	24 24 20 34 44 37 29	7 A 5 A 7 A 1 A 4 A 12 P 12 P	33.5 35.5 36.0 40.5 50.0 43.5 39.0	68.7 81.0 78.0 95.0 86.3 82.3 83.7	5.7 5.6 6.3 0.0 2.3 0.1 3.9	NW NW NW NW NW NE NE	172 16 17 17 50 61 39	32 5 7 1 6 9 7	5 р	10 р	0.10	
19 20 21	41 45 55	3 P 8 P 12 M	24 38 43	1 A 2 A 0 A	32.5 41.5 49.0	91.0 91.7 79.7	1.9 0.0 1.1	NE NE E	24 39 125	2 5 13	9 a	12 м	0.07	
22 23 24	52 55 53	12 P 12 M 3 P	47 40 32	0 A 12 P 8 A	49.5 47.5 42.5	89.0 80.3 76.3	0.0 3.4 5.7	E S S	195 39 73	25 9 20	11 A 1 A	9 p 2 a	0.54	
25 26 27 28 29 30	45 50 56 41 40 38	2 P 3 P 4 P 0 A 2 P 2 P	27 34 41 30 23 25	7 A 1 A 12 P 12 P 7 A 0 A	36.0 42.0 48.5 35.5 31.5 31.5	81.0 83.0 73.7 65.3 70.0 73.7	3.4 0.0 4.4 5.6 6.6 5.3	NW W NW NW NW	11 60 96 154 81 96	2 12 10 25 9 25	0 а	1 A	0.01	

^{*} Based on least time required to blow one mile.

HANS JOA, Observer

MONTILI	SUMM	IAICI
NOVEMBER 1952	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.46 Minimum 29.43 Mean semi-daily 30.10	30.59 29.30 30.05	30.87, 1932 28.73, 1904
Range	1.29	1.84, 1904; 0.89, 1943
Air Temperature, in Degrees F.		
Highest 69, 1st Lowest 18, 9th Mean 40.4 Range 51.0 Highest mean daily 25.0, 9th Mean maximum 49.0 Mean minimum 31.8 Greatest daily range 35, 2nd Least daily range 5, 22nd Degree-days of heating load for month 737 Degree-days cumulative for season 1294	66.0 12.9 38.9 53.0	82, 1950 -4, 1938 44.1, 1931; 33.6, 1901
Precipitation, in inches		
Precipitation 0.89 Snow 1.00 Maximum precip, in 24 hrs. 0.54	3.41 2.34	8.64, 1927; 0.63, 1917 13.50, 1938
Number of days with .01 or more 7	9	22, 1921; 2, 1904
Wind, in miles		
Total movement. 2409 Greatest daily movement 195, 22nd Least daily movement 3, 9th Mean hourly velocity 3.3 Maximum velocity 32, 12th	4531 6.3 30.3	5,978, 1906; 2,589, 1889 48, 1948, 1950
Wind, direction	30.3	2", 2 2", 2 2"
Prevailing direction NW North, days - Northeast, days 4 East, days 2 Southeast, days 3 South, days 3 Southwest, days 5 Northwest, days 16	WNW	
Weather		
Mean relative humidity, percent	70.6 55.1 6 9 15 121 41.3	78, 1947; 34, 1917 15, 1903; 0, 1948 16, 1912; 4, 1889, 1930, '32, '35 24, 1927; 9, 1905, 1917, 1949 182, 1903; 44, 1948
First Snow	Nov. 6	Oct. 10, 1925; Nov. 27, 1931

Note—The first column in the above summary lists observations made during the month. The second column lists the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column lists extremes observed from 1889 to 1951.

REMARKS

The mean daily air temperature during the month of November was above normal; the degree days for the month were 737 compared to the normal of 783.

Precipitation for the month was 0.89 inches compared to the normal of 3.41 inches.

The mean relative humidity for the month was 77.0 per cent, mean cloudiness 56.7 per cent; there were 108.2 hours of bright sunshine during November.

Meteorological Series

Bulletin No. 768

December 1952

Meteorological Observations

FOR

DECEMBER

1952

f. J. PFLUG

OBSERVATORY

Latitude, 42° 23′ 31″ N.

Height of barometer above ground, 36 ft.

Height of wind instruments, 67 ft.

Longitude, 72° 31′ 48″ W

Above sea level, 253.5 ft.

Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS

	1 cmportar a. v						Wind	l Precipitation						
Day	Degrees Ma	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hour- Bright Sunspine	Prevailing Direction	Tota! Movement	Max Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 27 28 29 29 20 20 20 20 20 20 20 20 20 20 20 20 20	228 42 45 44 46 38 44 46 40 40 40 40 41 35 41 48 43 43 44 48 41 48 41 48 48 49 40 40 40 40 40 40 40 40 40 40 40 40 40	3 P 3 P 3 P 3 P 1 P 2 P 1 P 0 A 2 P 3 P 2 P M 2 P 3 P 2 P M 2 P M 2 P M 2 P M 4 P 2 P M 4 P 2 P M 4 P 2 P M 4 P 4 P 4 P 4 P 4 P 4 P 4 P 4 P	18 16 22 19 25 29 22 27 17 25 32 27 17 25 32 29 29 38 32 29 29 38 32 29 29 38 32 27 17 18 18 18 18 18 18 1	8 A 6 A 1 A 7 A 8 A 6 A 1 2 P 8 A 1 2 P 7 A 2 A 8 A 1 2 P 10 P A 1 2 P 10 P A 8 A 8 A 8 A 8 A	25.0 22.0 32.0 32.0 32.0 32.0 32.0 32.0 36.0 37.5 30.0 34.0 50.0 36.5 34.5 26.0 33.0 40.0 33.0 40.0 33.0 40.0 33.0 40.0 33.0 40.0 31.5 34.0 34.5 34.0 34.5 34.0 35.0 36.5 34.0 36.5 34.0 36.5 34.0 36.5 34.0 36.5 34.0 36.5 34.0 36.5 34.0 36.5 34.0 36.5 34.0 36.5 34.0 36.5 34.0 36.5 34.0 36.5 34.0 36.5 34.0 36.5	70.0 89.0 77.3 77.0 95.7 77.3 83.0 94.3 85.3 85.3 85.3 85.3 84.7 74.7 62.7 73.0 65.0 65.0 65.0 75.3 60.7 79.0 795.0 75.3 60.7 75.3 75.3 75.3 75.3 75.3 75.3 75.3 75	6.8 0.0 5.4 6.6 0.0 0.0 3.2 5.8 1.9 0.0 0.7 5.6 2.3 5.8 4.0 0.0 0.7 5.4 2.5 5.8 4.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0	NW N N N N N N N N N N N N N N N N N N	133 355 1300 811 744 1355 61 107 71 249 140 134 102 53 141 144 125 74 36 56 180 199 186 202 202 202 366 114	17 7 11 8 13 17 17 4 4 4 4 14 25 25 25 17 7 20 20 20 20 22 11 22 11 22 12 13 14 25 15 16 17 17 17 18 18 18 18 18 18 18 18 18 18 18 18 18	1 A O A 4 A O A	12 P 1 A	1.12 0.09 2.10 0.76 T	T

^{*} Based on least time required to blow one mile.

llans Joa, Observer

MONTILI		IAIXI
DECEMBER 1952	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.595, 4th Minimum 29.431, 11th Mean scmi-daily 30.056	30.66 29.27 30.06 1.38	31.104, 1949 28.85, 1915
Range	1.58	1.78, 1895, 1.01, 1892
Air Temperature, in degrees F.		, f f 1000
Highest 60, 11th Lowest 5, 28th Mean 32.6 Range 55 Highest mean daily 50.0	54.3 1.09 27.5 56.2	65.5, 1908 —22.5, 1917 36.9, 1891; 17.1, 1917
Lowest mean daily 14.5 Mean maximum 39.7 Mean minimum 25.5 Greatest daily range 26.4th Least daily range 5, 23rd Degree-days of heating load for month 998 Degree-days cumulative for season 2292	1162.5 2494.0	
Precipitation, in inches		
Precipitation 4.10 Snow T Maximum precip. in 24 hrs. 2.10, 11th	3.39 8.50	7.77, 1901; .58, 1943 26.50, 1902, T. 1891, 1943
Number of days with .01 or more 5	10	17, 1902; 4, 1892, 1943
Wind, in miles		
Total movement	4710	6694, 1925; 3239, 1918
Mean hourly velocity	6.3	66, 1949
Wind, direction		
Prevailing direction WNW North, days	WNW	
Weather		
Mean relative humidity, percent	69.4 54.9 8 9 14 128	71, 1929; 39, 1919 15, 1890; 0, 1948 16, '09; 4, '89, '30, '31, '36, '38 25, 1948; 7, '09, '23 172, 1896; 27, 1948
Percent of possible hours of bright sunshine	45.2 Nov. 6	

Note—The first column in the above summary lists observations made during the month. The second column lists the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column lists extremes observed from 1889 to 1951.

ANNUAL SUMMARY

ANNUAL	Normal	Extremes
Barometer, in inches (Readings reduced to sea level) Maximum	30.81 28.95	31.10, 1949
Minimum 29.122, Mar. 11th Mean semi-daily 29.979 Range 1.538	30.01 1.85	28.41, 1938 2.47, 1938; 1.38, 1933
Air Temperature, in degrees F. Highest 98, June 26th Lowest -7, Jan. 30th Mean 49.9 Range 105 Highest mean daily 84.5, June 26th	95.7 —12.2 47.4 107.8	104, 1911 —26, 1904 51.7, 1949; 44, 1904
Lowest mean daily 6.5 Jan. 30th Mean maximum 60.1 Mean minimum 39.7 Greatest daily range 45, April 22nd Least daily range 3, Jan. 6, 28, July 10	57.8 36.5	
Degree-days 1951-52 heating season6223	6698	
Precipitation, in inches Precipitation 40.61 Snow 38.3	43.70 47.78	59.00, 1938; 30.68, 1908 89.00, 1893; 24.50, 1919
Maximum precip. in 24 hrs2.70, June 1st Number of days with .01 or more97	124	146, 1945; 96, 1924
Wind, in miles		
Total movement	52,223	63,571, 1908; 36,257, 1894 705, 1909
Mean hourly velocity	5.8 39.5	80, 1938
Wind, direction Prevailing direction NNW North, days 38 Northeast, days 41 East, days 13 Southeast, days 40 South, days 41 Southwest, days 37 West, days 44 Northwest, days 112	W	
Weather		
Mean relative humidity, percent 76.6 Mean cloudiness, percent 49.7 Number of clear days 92 Number of fair days 146 Number of cloudy days 127 Number of hours bright sunshine 2134 Percent of possible hours of	67.6 51.7 116 123 126 2,353	60, '98, '01, '02; 41, '08, '24 217, 1941; 59, 1927 182, 1912; 64, 1936 179, '01, '02; 71, 1910, 1941 3038, 1941; 1864, 1902
bright sunshine 46.6 Last snow Mar 19th First snow Nov. 8th Last frost May 17th First frost Oct. 11th	52.8 April 15 Nov. 6 May 14 Sept. 21	Mar. 14 '10; May 11, '07, '45 Oct. 10, '25; Dec. 13, '41 Apr. 14, '43; June 8, '32 Aug. 22, '94, '95; Oct. 13, '09

Meteorological Series

Bulletin No. 769

January 1953

Meteorological Observations

FOR

JANUARY

1953

1. J. PFLUG

OBSERVATORY

Latitude, 42° 23′ 31″ N.

Longitude, 72° 31′ 48" W.

Height of barometer above ground, 36 ft. Above sea level, 253.5 ft.

Height of wind instruments, 67 ft.

Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS

		Te	mpera	ture		ty	or,		Wind]	Precipita	ation	
	Max	imum	Mir	imum		Humidi	of Hour inshine	50 _	at	*es				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5	40 38 39 36	4 P 8 P 1 P 3 P	21 16 33 24	8 A 7 A 8 A 12 P	30.5 27.0 36.0 30.0	80.0 92.3 90.0 66.0	6.1 1.6 1.3 6.3	N N NW NW	114 19 111 206	10 6 17 25	0 а	М	1.01	Т
6 7 8	36 32 31 24	3 P 4 P 3 P 4 P	17 18 19 19	8 A 11 P 8 A 8 A	26.5 25.0 25.0 21.5	78.0 74.7 78.7 94.0	6.6 2.7 62. 1.8	NW S NW NW	73 118 52 60	28 5 8	5 A 7 P	8 A 12 P	T 0.04 0.13	0.1 1.8 0.8
9 10 11 12 13	28 33 37 37 42	11 P 8 P 1 P 4 A 4 P	20 28 32 24 24	0 A 0 A 2 A 12 P 0 A	24.0 30.5 34.5 30.5 33.0	91.7 95.0 95.0 60.3 68.3	0 0.8 3.9 2.0	N N N NW SE	207 198 145 203 113	17 17 8 32 13	0 A 0 A 3 A	12 P 10 P 7 A 8 A	1.09 0.93 0.04 T 0.02	6.3 0.4 0.1 0.4
13 14 15 16 17	44 42 47 28	3 P 3 P 2 P 7 P	24 22 24 26 13	7 A 8 A 12 P 8 A	33.0 33.0 36.5 20.5	86.0 93.7 85.7 75.3	5.6 6.2 0.9 9.4	SE S S NW NW	17 12 141 126	7 3 32 12	2.1	OA	0.02	0.4
18 19 20 21	49 42 40 39	2 P 4 P 2 P 2 P	24 28 31 30	0 A 7 A 4 A 12 P	36.5 35.0 35.5 34.5	97.0 78.3 80.7 79.7	3.6 3.2 1.5 0.4	NW NW NW	38 32 11 75	12 11 5 10	0 A 5 P	8 A	0.70	1.4
22 23 24 25	42 35 52 48	3 P 2 P 8 P 0 A	26 16 33 21	9 A 7 A 0 A 12 P	34.0 25.5 42.5 34.5	81.7 88.3 97.3 69.7	7.4 1.5 0 2.1	N N S	109 14 94 274	11 5 20 39	2 A 2 A	7 P	1.36 0.19	
26 27 28 29	30 22 41 39	2 P 12 P 3 P 3 P	15 11 22 26	12 P 6 A 0 A 7 A	22.5 16.5 31.5 32.5	58.0 83.3 89.7 73.3	7.0 1.6 1.1 7.0	NW NW NW NW	327 25 107 159	28 3 25 20	7 Р 1 А	12 P 11 A	0.05 0.51	0.5
30 31	32 48	12 P 3 P	20 32	8 A 0 A	26.0 40.0	67.7 80.6	6.6 4.8	NW S	127 184	14 20			T	

^{*} Based on least time required to blow one mile.

HANS JOA, Observer

MOMINE	OUI-II-	
JANUARY 1953	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.605, 17th Minimum 29.039, 25th Mean semi-daily 29.994 Range 1.566	30.70 29.20 30.07 1.50	31.00, 1927 28.55, 1913 2.18, 1913; 0.97, 1896
Air Temperature, in degrees F.	1.50	2.16, 1713, 6.77, 1076
Highest 52, 24th Lowest 11, 27th Mean 30.4 Range 41 Highest mean daily 42.5, 24th Lowest mean daily 16.5, 27th	51.1 -7.1 24.2 58.2	66.0, 1932 —26.0, 1904 34.2, 1913; 13.9, 1918
Mean maximum 37.8 Mean minimum 23.1 Greatest daily range 27, 25th Least daily range 5, 8th, 10th, 11th Degree-days of heating load for month 1071 Degree-days cumulative for season 3363	1265	
Precipitation, in inches		
Precipitation 6.24 Snow 11.8	3.61 13.32	7.15, 1898; 1.07, 1896 33, 1898, 1923 1.50, '08
Maximum precip. in 24 hrs 1.36 Number of days with .01 or more 13	11	17, '50, '20; 4, 1901
Wind, in miles		
Total movement3491Greatest daily movement327, 26thLeast daily movement11 20thMean hourly velocity4.07Maximum velocity39, 25th	5055 6.8 32.7	7770, 1908; 28%, 1895 47, 1938
Wind, direction		
Prevailing direction NNW North, days	WNW	
Weather		
Mean relative humidity, percent \$1.6 Mean cloudiness, percent 59.2 Number of clear days 10 Number of fair days 6 Number of cloudy days 15 Number of hours bright sunshine 109.2 Percent of possible hours of bright sunshine 37.1 Thunder and lightning	70.1 55 9 9 13 137 46.7	78, 1932; 37, 1908 20, 1944; 1, 1948 18, 1890, 1926; 3, 1916, 1949 22, 1923, '31, '49; 4, 1920, '39 214, 1920; 50.4, 1949

Note—The first column in the above summary lists observations made during the month. The second column lists the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column lists extremes observed from 1889 to 1952.

REMARKS

The weather during the month of January was exceptionally mild. This abnormal weather is reflected in the mean daily temperature, which is 6.2 degrees above normal, and the degree days for the month which are 1071, compared to a normal of 1265.

The precipitation for January was 6.24 inches, compared to the normal of 3.61 inches.

The mean relative humidity was 81.6 percent, mean cloudiness 59.2 percent. Snowfall and snow accumulation were both below normal.

Meteorological Series

Bulletin No. 770

February 1953

Meteorological Observations

FOR

FEBRUARY

1953

I. J. PFLUG

OBSERVATORY

Latitude, 42° 23′ 31″ N.

Longitude, 72° 31′ 48″ W

Height of barometer above ground, 36 ft. Above sea level, 253.5 ft.

Height of wind instruments, 67 ft.

Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS

		Te	mpera	iture		>	un.		Wind]		Precipita	ation	
У	Degrees Way	Time	Degrees W	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max Miles per Hour*	Began	Ended	Water	Snow
Day						<u> </u>					Be	뗩		
1 2 3 4	46 24 29 36	0 A 8 P 8 P	10 7 21 17	12 P 6 A 7 A 12 P	28.0 15.5 25.0 26.5	59.7 58.7 95.3 75.3	7.0 6.0 0.0 6.2	NW NW SE N	411 146 49 182	48 25 8 20	1 A	11 A	0.02	т 3.6
5 6 7 8	38 40 46 36	3 P 8 P 3 P 3 P	10 12 36 31	7 A 7 A 8 A 2 A	24.0 26.0 41.0 33.5	80.3 88.7 93.7 74.7	7.5 3.4 0.0 0.0	N N NW N	31 42 40 165	10 6 6 17	10 p 2 A	12 P 8 P	0.06 0.79	
9 10 11 12	38 3 38 30	3 P 3 P 3 P 0 A	30 21 19 26	12 P 8 A 8 A	34.0 27.0 28.5 28.0	60.0 56.0 60.7 95.0	5.6 7.8 7.5 0.4	NW NW NW NE	291 468 59 97	32 32 10 9	Р 0 А	12 P 11 A	0.03 0.49	т 7.4
13 14 15 16	40 44 39 33	3 P 5 P 5 P	26 25 24 25	0 A 1 A 7 A 8 A	33.0 34.5 31.5 29.0	82.0 67.7 100.0 56.3	5.4 7.5 0.0 8.0	E SW NW N	32 107 80 358	11 10 48 48	9 A	— 9 р	0.01	0.2 T
17 18 19 20 21 22	33 48 52 57 36	2 P 3 P 4 P 4 P 2 P	24 20 22 24 36 24	3 A 7 A 7 A 7 A 12 P 12 P	28.5 26.5 35.0 38.0 46.5 30.0	63.0 59.7 78.7 84.0 82.0 59.0	7.8 8.4 8.0 5.4 0.5 7.1	N N NW S NW	250 264 14 45 307 378	25 25 4 13 28 25	6 A	10 A	0.01	0.2
23 24 25 26 27	38 54 46 44 48	3 P 5 P 3 P 5 P 2 P	21 29 27 26 34	8 A 0 A 12 P 3 A 6 A	29.5 41.5 36.5 35.0 41.0	67.7 69.0 66.0 96.0 75.7	7.1 8.0 8.1 2.4 5.6	NW NW NW W SW	130 97 141 40 190	20 17 14 17 39	7 A	9 A	0.03	0.1
28	37	М	22	12 Р	29.5	63.0	7.2	NW	209	32	5 A	6 A	0.01	0.2

^{*} Based on least time required to blow one mile.

HANS JOA, Observer

MONTILL	SOITH.	
FEBRUARY 1953	Normal	Extremes
Barometer, in inches		
(Readings reduced to sea level)		
Maximum	30.63	31.05, 192
Minimum	29.24 30.04	28.56, 189
Range	1.38	1.89, '00, '08; .88, '13, '3
Air Temperature, in degrees F.		
Highest	50.5	65.0, 193
Lowest	-7.4 23.7	23, 194 32.6, 1890; 11.6, 193
Range	58	32W, 1370, 11W, 37V
Mean maximum	32.8	
Mean minimum	14.5	
Least daily range		
Degree-days of heating load for month 937.5 Degree-days cumulative for season. 4300.5	1167 4927	
Precipitation, in inches		
Precipitation	3.19	8.12, 1900; 0.62, 190
Snow	13.99	48.75, 1893; 0.50, 193
Maximum precip. in 24 hrs 1.08, 15th Number of days with .01 or more12	10	15, 1893, 1920; 4, 190
Wind, in miles		
Total movement	4776	6445, 1896; 3402, 19-
Least daily movement	7.1	
Maximum velocity	31.7	50, 19-
Wind. direction		
Prevailing direction NNW North, days 8 Northeast, days 1	WNW	
East, days		
South, days1		
Southwest, days	!	
Northwest, days		
Weather		
Mean relative humidity, percent73.9	06.6	// 1000 1007 21 10
Mean cloudiness, percent	50.9	66, 1890, 1927; 31, 19 19, 1941; 2, 19
Number of fair days8	8	19, 1941; 2, 19 16,1920; 2, 19
Number of cloudy days	10 158	16, 1894; 2, 19 221, 1924; 110, '27, '
Percent of possible hours of bright sunshine	53.4	
Digit sunsinc	7.7.1	

Note—The first column in the above summary lists observations made during the month. The second column lists the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column lists extremes observed from 1889 to 1952.

The mean daily temperature for the month of February was 7.8 degrees (F) above normal. This is the highest February mean daily temperature since 1925. February is the fourth consecutive month having a mean daily temperature above normal.

The degree days accumulative for the 1952-53 heating season are 4300.5 compared to a normal of 4927. The mild winter is reflected in the low number of degree days.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No 771

March 1953

Meteorological Observations

FOR

MARCH

1953

I. J. PFLUG

OBSERVATORY

Latitude, 42° 23' 31" N.

Longitude, 72° 31′ 48″ W

Height of barometer above ground, 36 ft. Above sea level, 253.5 ft.

Height of wind instruments, 67 ft.

Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS

		Te	трега	ture		r,			Wind		P	recipita	tion
Day	Degrees	Time mnmi	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water
1 2 3 4 5 6 7 8 9 10 11 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 20 20 20 20 20 20 20 20 20 20 20 20 20	30 32 38 53 40 40 31 28 34 27 52 51 55 52 39 45 52 39 45 57 57 57 57 57 57 57 57 57 57 57 57 57	3 P 4 P 5 P 3 P 2 P 2 P 4 P 4 P 2 P 3 P 4 P 4 P 3 P 1 A 1 P 4 P 3 P 1 A 1 P 4 P 3 P 1 P 4 P 3 P 1 P 4 P 3 P 1 P 4 P 3 P 1 P 1 P 1 P 1 P 1 P 1 P 1 P 1	16 10 14 32 30 27 19 15 14 14 14 8 28 42 36 32 32 32 32 51 41 40 34 40 34 40 34	7 A 5 A 7 A 0 A 12 P 7 A 12 P 7 A 12 P 7 A 8 A 12 P 7 A 8 A 12 P 6 A 12 P 7 A 6 A 12 P 7 A 6 A 12 P 7 A 6 A 12 P 7 A 8 A 12 P 7 A 8 A	23.0 21.0 26.0 42.5 35.0 33.5 25.0 20.5 20.5 22.5 40.0 46.5 45.5 40.0 36.5 39.5 41.5 49.5 55.5 49.0 53.0 47.5 45.0 49.5 45.0 49.5 45.0 49.5 45.0 49.5 45.0 49.5 45.0 49.5 45.0 49.5 45.0 46.0 46.0 46.0 49.0 49.0 49.0 49.0 49.0 49.0 49.0 49	62.3 57.7 72.7 94.0 72.0 72.0 64.3 51.3 46.3 58.0 92.0 96.3 72.0 98.3 79.7 70.3 63.0 81.7 65.7 70.7 73.0 88.7 84.0 82.7 92.7 86.0 82.7 92.7 86.0 87.0	8,6 8,3 6,8 6,8 6,3 7,8 9,8 8,4 9,0 2,8 0,0 4,6 2,0 8,6 9,6 9,6 9,6 9,6 9,6 9,6 1,6 1,6 1,6 1,6 1,6 1,6 1,6 1,6 1,6 1	NW N N N N N N N N N N N N N N N N N N	242 188 115 129 292 277 349 102 187 108 93 135 28 42 40 69 118 138 216 101 142 149 94 437 250	32 13 13 13 15 48 32 25 28 13 8 20 9 5 17 7 9 10 16 6 10 10 10 10 10 10 10 10	9 P 0 A 8 P 0 A 8 A 0 A 1 P 0 A 7 A 7 A	12 P 8 A 12 P 3 P 12 P 3 A (12P 3 P 11 P 12 P 8 P 12 P 8 A	0.07 0.91 T T 1.58 0.19 1.58 0.19 0.20 0.04 T 1.19 0.37 0.32 0.21 0.32 0.91 0.01

^{*} Based on least time required to blow one mile.

HANS JOA, Observer

MONTHLI		IAKI
MARCH 1953	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30,690, 11th Minimum 29,365, 30th Mean semi-daily 30,021	30.57 29.24 30.04	31.05, 1943 28.47, 1914
Range	1.34	2.10, 1914; 0.85, 1915
Air Temperature, in degrees F.		05.10.1
Highest 67, 23rd Lowest 8, 11th Mean 38.2 Range 59	63.8 6.1 34.4 57.6	85, 1945 14, 1946
Highest mean daily 55.5, 24th Lowest mean daily 20.5, 10th Mean maximum 45.9 Mean minimum 30.5 Greatest daily range 35.0, 23rd	43.2 25.4	
Least daily range	949 58 7 6	
Precipitation, in inches		
Precipitation	3.70 7.47	7.89, 1942; 0.12, 191 27, 1899; 0, 192
Number of days with .01 or more15	11	17, 1890, 1913, 1936; 3, 191
Wind, in miles		
Total movement 4829 Greatest daily movement 437, 30th Least daily movement 13, 15th		8182, 1896; 3006, 190
Mean hourly velocity	1 7.7	48, 1932, 1939, 1941, 194
Wind, direction		
Prevailing direction NW North, days 3 Northeast, days East, days 1 Southeast, days 5 South, days 5 Southwest, days 1 West, days 5 Northwest, days 1		
Weather		
Mean relative humidity, percent. 74.7 Mean cloudiness, percent	51.8 11 10 10 10 199	68, 1901; 27, 191 22, 1924; 3, 190 17, 1900; 1, 194 21, 1901; 1, 191 292, 1924; 93, 191

Note—The first column in the above summary lists observations made during the month. The second column lists the averages based on observations made from 1889 to 1938, except that humidity seconds are based on observations made from 1929 to 1938. The third column lists extremes observed from 1889 to 1952.

The mean daily temperature during March was 3.3°F, above normal. This is the fifth consecutive month that the mean temperature has been above normal. The cumulative degree days for the heating season through March is 5132 degree days compared to the normal of 5876.

The rainfall during March was 8.28 inches; this is 4.58 inches above normal and is the most rainfall recorded in March since the start of our records in 1889. The rainfall occurred on 15 days; the maximum in 24 hours was 1.77 inches. The rainfall during March was the most for any month since the September 1938 hurricane. Dr. Ives of Amherst College points out that water accumulation since January 1 is about 70 percent above normal.

Dr. Frank Southwick of the Pomology Department makes the following remarks:

"Mild temperatures were experienced during January, February, and March. Fruit plants appear to have come through the winter with virtually no signs of winter injury. In McIntosh apple blocks where the crop was light in 1952, it can now be seen that a very heavy bloom is going to occur. Peaches apparently will bloom heavily, also. At the end of March, apple flower buds had commenced to swell but were still in suitable condition for a dormant spray application. The almost continuous rain period during the latter part of March made spraying and crop planting difficult or impossible, however."

UNIVERSITY OF MASSACHUSE LIBRARY

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 772

April 1953

Meteorological Observations

FOR

APRIL.

1953

I. J. PFLUG

OBSERVATORY

Latitude, 42° 23′ 31″ N.

Longitude, 72° 31′ 48″ W.

Height of barometer above ground, 36 ft. Above sea level, 253.5 ft.

Height of wind instruments, 67 ft.

Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Te	mpera	ture		£3	ø		Wind]	Precipita	ation
	Max	imum	Mir	imum		Humidi	of Hour inshine	p.c.	Ħ	*es			
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18	60 63 65 62 57 59 64 52 47 59 47 59 44	1 P 1 P 4 P 2 P 1 P 4 P 3 P 5 P 10 A 5 P 11 A 3 P 5 P 11 A 7 P 10 P 11 A	40 41 33 33 42 32 41 37 44 41 32 33 33 33 33 33 42 37 44 41 33 33 33 33 33 34 34 35 35 36 37 37 37 37 37 37 37 37 37 37 37 37 37	6 A 12 P 5 A 12 P 6 A 12 P 12 P 12 P 14 A 4 A 9 P 1 A 4 A 6 A 3 A 4 A 12 P	50.0 50.5 49.0 49.0 52.0 44.5 44.5 50.0 50.5 49.0 50.5 49.0 50.5 49.0 50.5 49.0 50.5 49.0 50.5 49.0 50.5 49.0 50.5 49.0 50.5 49.0 50.5 49.0 50.5 49.0 50.5 49.0 50.5 49.0 50.5 49.0 50.5 49.0 50.5 49.0 49.0 50.5 49.0 49.0 49.0 50.5 49.0 49.0 49.0 50.5 49.0	74.0 79.7 68.7 60.7 63.3 60.0 87.0 73.3 66.7 90.0 71.3 94.3 64.3 45.0 89.3 61.0 72.3 85.7	7.2 0.8 10.2 9.4 8.8 9.0 0.0 8.7 9.6 0.0 8.7 3.2 0.3 6.6 10.9 0.0 9.0 1.6	NW NW SE NW NW SE SE W SW NW W W W W W W W NW	42 61 48 89 91 60 69 128 46 149 162 395 302 82 235 75 123	5 20 5 14 17 7 10 8 11 18 32 32 32 32 32 31	9 P 2 A 11 P 0 A 1 A 1 A 2 P 4 P 0 A 5 P M	11 P 6 P 12 P 2 A 11 P 12 P 12 P 12 P 8 A 6 P 12 P 12 P	0.03 0.25 0.03 0.02 0.85 0.37 0.71 1.05 0.36 1.4 0.61 T 0.15 0.29
19 20 21 22 23 24 25 26 27 28 29 30	41 44 47 65 69 69 63 56 63 51 61 67	11 A 3 P 1 P 2 P 1 P 4 P 3 P 1 P 1 P 1 P 5 P 3 P	33 36 34 49 46 48 50 46 40 38 35	12 P 0 A 3 A 5 A 4 A 5 A 3 A 12 P 12 P 9 P 4 A 6 A	37.0 38.5 41.5 49.5 59.0 57.5 55.5 53.0 54.5 45.5 49.5 51.0	85.7 69.3 64.0 51.3 44.7 46.0 68.7 88.0 73.0 62.3 53.0 50.0	0.0 6.5 6.3 9.8 10.3 10.9 5.9 0.8 7.5 7.2 10.3 10.3	W W S SE W W W	185 233 182 341 187 65 36 138 291 226 68	25 28 25 39 20 10 28 20 32 28 11	6 A 0 A	8 A 12 P	0.29 0.07 0.4 T 0.02 0.55

^{*} Based on least time required to blow one mile.

HANS JOA, Observer

30.48 29.38 29.99 1.11 79.4 22.0	30.71, 191 28.98, 194 1.52, 1930; 0.65, 1950
29.38 29.99 1.11 79.4 22.0	28.98, 1943
29.38 29.99 1.11 79.4 22.0	28.98, 1943
79.4 22.0	1.52, 1930; 0.65, 1950
22.0	
22.0	1
45.7 57.4	90, 1941 8.5, 1923 52.1, 1941; 41, 1943
56.8 34.8	
579 6455	
3.35 2.17	6.89, 1929; .55, 1941 11, 1891; 0, 1910, 1934
11	18, 1909; 3, 1892
7.5	8208, 1908; 3853, 1917
31.+	40, 1935, 1938, 1940, 1945
WNW ·	
61.6 51.8 11 9 10 220 54.7	75, 1901; 34, 1925, 1927 23, 1941; 1, 1949 18, 1915; 2, 1894, 1901 22, 1901; 1, 1941 329, 1941; 103, 1901
	56.8 34.8 579 6455 3.35 2.17 11 5404 7.5 31.4 WNW

Note—The first column in the above summary lists observations made during the month. The second column lists the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column lists extremes observed from 1889 to 1952.

The mean daily temperature during April was 1.9 degrees above normal. The precipitation was 5.36 inches, 2.01 inches above normal; the precipitation since January 1 was 22.85 inches.

Dr. Frank Southwick of the Pomology Department made the following observations:

"Although April averaged 1.9 degrees above normal, bud development of fruit plants at the end of the month was about average. Apparently the maximum daily temperatures were sufficiently low to impede growth even though the average mean temperatures were above normal. McIntosh apples reached delayed dormant about April 10. They remained dormant about two weeks and were not in a full pink stage at the end of the month.

"About April 25, plums, cherries, and peaches commenced to bloom, but are not expected to reach full bloom until early May. Since the blossom period will probably occur about the usual time, the chances of loss of fruit from spring frosts do not appear to be great. Excessive rainfall in April has made scab control a difficult problem for apple growers and has delayed the planting of annual crops."

Publication of This Document Approved by George J. Cronin, State Purchasing Agent.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 773

May 1953

Meteorological Observations

FOR

MAY

1953

I. J. PFLUG

OBSERVATORY

Latitude, 42° 23′ 31″ N.

Longitude, 72° 31′ 48″ W.

Height of barometer above ground, 36 ft. Above sea level, 253.5 ft.

Height of wind instruments, 67 ft.

Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Te	mpera	ature		ty	ø		Wind			Precipita	tion	
	Max	imum	Mir	nimum		Humidi	of Hour inshine	8 _	ij	les **				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1	47	2 P	38	12 P	42.5	82.0	0.6	N	109	13	1 A 3 P	8 A 9 P	1.33	
2	44	2 P	38	6 а	41.0	81.7	0.0	N	75	10	0 A	12 P	0.24	
3	54	3 P	37	4 A	45.5	78.0	8.4	Е	70	4	0 A	1 A	0.02	
4	73	5 P	33	5 A	53.0	63.0	11.6	NW	44	5				
5	64	5 P	51	7 A	57.5	89.7	6.2	s	72	4	5 A	10 A	0.28	
6	7 9	3 Р	47	5 A	63.0	68.0	10.0	s	7 9	12				
7	69	1 P	58	4 A	63.5	89.7	0.2	s	70	4	4 P	9 P	0.19	
8	72	3 P	58	6 A	65.0	88.3	5.6	SE	80	13	4 A 6 P	6 A 10 P	0.59	
9	79	2 Р	54	5 A	66.5	78.3	8.4	w	59	5	2 P	3 Р	0.01	
10	87	4 P	50	5 A	68.5	65.7	11.0	sw	37	4				
11	90	4 P	51	5 A	70.5	56.3	9.0	s	24	5		ł		
12	88	4 P	56	5 Λ	72.0	62.0	10.5	SE	49	4				
13	7 9	3 P	61	5 A	70.0	73.3	7.7	SE	106	10	7 P	10 P	0.18	
14	72	2 Р	57	12 P	64.5	82.7	4.8	SE	65	9				
15	57	5 P	54	5 A	55.5	93.3	0.0	SE	73	-8	9 A	4 P	0.44	
16	74	6 P	55	0 A	64.5	73.3	11.8	N	57	10				
17	60	3 P	53	12 P	56.5	93.0	0.0	SE	73	8	1 A	8 A	0.73	
18	7 9	4 P	52	3 A	65.5	78.7	6.5	SE	95	25	1 A	5 A	0.28	
19	77	2 P	55	5 A	66.0	64.7	10.1	w	58	7				
20	80	2 P	51	5 A	65.5	68.3	10.5	sw	39	7				
21	76	3 P	52	5 A	64.0	65.3	11.4	E	85	15				
22	75	6 P	51	1 A	63.0	85.0	4.3	E	106	10	11 P	12 P	0.37	
23	69	2 P	52	12 P	60.5	61.7	11.2	W	241	25	0 A	2 A	0.66	
24	73	4 P	43	4 A	58.0	55.3	11.8	W	188	22				
25	74	2 P	46	4 A	60.0	63.0	11.8	NW	100	20				
26	70	11 A	46	5 A	58.0	65.0	6.0	NW	54	15	2 P	9 P	0.83	
27	69	M	46	3 A	57.5	72.7	7.7	NW	94	20	3 P	12 P	0.54	
28	59	M	47	3 A	53.0	68.7	10.2	NW	245	25	0 A	2 A	0.07	
29	66	3 r	41	5 A	53.5	54.0	11.3	NW	128	15				
30	59	M	43	2 A	51.0	74.7	6.9	NW	41	5				
31	70	4 P	40	5 A	55.0	63.3	10.2	NW	42	8				

^{*} Based on least time required to blow one mile.

MONTHLI	SUMME	11/1
MAY, 1953	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.293, 4th Minimum 29.513, 18th Mean scmi-daily 29.911 Range 0.780	30.39 29.51 29.96 0.88	30.62, 1936 29.10, 1938
Air Temperature, in degrees F.		
Highest 90, 11th Lowest 33, 4th Mean 59,7 Range .57 Highest mean daily 72.0, 12th Lowest mean daily 110, 22th	86.1 31.2 57.1 54.9	94.5, 1896, 1911 24.0, 1900
Lowest mean daily 41.0, 2nd Mean maximum 70.5 Mean minimum 48.9 Greatest daily range 40, 4th Least daily range 3, 15th Degree-days of heating load for month 189.5 Degree-days cumulative for season 5841.5	56.8 34.8	
Precipitation, in inches		
Precipitation	3.60	7.44, 1931; .48, 1903
Snow	12	20, 1901, 1945; 5, 1903
Wind, in miles		
Total movement	6.1	5946, 1907; 2180, 1894
Maximum velocity 25, 18th, 23rd, 28th	28.8	45, 1935
Wind, direction Prevailing direction WNW North, days 3 Northeast, days - East, days 3 Southeast, days 7 South, days 4 Southwest, days 2 West, days 4 Northwest, days 8	W	
Weather		
Mean relative humidity, percent. 72.9 Mean cloudiness, percent 62.6 Number of clear days 12 Number of fair days 13 Number of cloudy days	60.7 52.1 11 11 9 252 55.6	72, 1948; 30, 1923 21, 1944; 0, 1927 18, 1949; 5, 1923 20, 1927; 2, 1941 350, 1944; 137, 1927
26th, and 27th		

Note — The first column in the above summary lists observations made during the month. The second column lists the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column lists extremes observed from 1889 to 1952.

The mean daily temperature for the month of May was 2.6 degrees above normal. The rainfall for the month was 6.76 inches compared to the normal of 3.60 inches.

Dr. Frank Southwick of the Pomology Department makes the following remarks:

"May was another very wet month with total rainfall about double the average amount. Temperatures for the month were near the seasonal average so that fruit plants generally blossomed at about the expected time. Peaches were in full bloom about May, and apples were in the pink stage of development on the same date. All apple varieties blossomed from May 9 to 13. Weather conditions during the apple blossoming period were excellent for pollination and fertilization. A good set of all fruits is anticipated. No killing frosts were experienced in the fruit-producing areas of the State during May."

Publication of This Document Approved by George J. Cronin, State Purchasing Agent.

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 774

June 1953

Meteorological Observations

FOR

JUNE

1953

L. J. PFLUG

OBSERVATORY

Latitude, 42° 23′ 31″ N.

Height of barometer above ground, 36 ft.

Height of wind instruments, 67 ft.

Longitude, 72° 31′ 48″ W.

Above sea level, 253.5 ft.

Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Te	mpera	iture		ž.	gs.		Wind		I	Precipita	ıtion
	Max	kimum	Mir	imum		Humidi	of Hour inshine	20	ıt	*es			
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water
1 2 3 4 5	75 74 72 78	3 P 4 P 5 P 3 P	43 45 41 48	5 A 5 A 5 A 3 A	59.0 59.5 56.5 63.0	50.7 43.3 45.7 57.3	11.6 12.0 12.5 6.4	NW NW NW	174 142 27	25 22 4			т
6 7 8	89 92 80 79	3 P 3 P 5 P 5 P	59 69 57 50	6 A 5 A 12 P 5 A	74.0 80.5 68.5 64.5	66.0 60.0 71.7 58.3	10.4 11.1 5.6 11.6	NE SW SW W	72 156 92 69	17 17 15 10	10 р 11 л	11 P 1 P	0.01
10 11 12	90 76 74 77	4 P 4 P 3 P	60 54 47 43	4 A 12 P 4 A 5 A	75.0 65.0 60.5 60.0	68.0 46.3 38.7 57.0	8.8 12.2 12.2 10.7	SE NW NW NW	215 281 192 59 57	36 25 25 13	3 P	7 р	
13 14 15 16 17	70 71 83 87 85	11 A 4 P 4 P 1 P	49 49 43 51 53	6 A 12 P 5 A 5 A 5 A	59.5 60.0 63.0 69.0 69.0	61.7 63.0 60.0 55.0	7.5 10.6 9.0 11.3 8.9	SE NE NE NW	106 53 38 67	10 12 4 8	,5 P	/ P	0.04
18 19 20	82 90 98	4 P 5 P 4 P 4 P	59 58 63	5 A 5 A 4 A	70.5 74.0 80.5	60.7 62.0 66.7 66.7	10.9 10.6 10.8	SE SE W	70 47 54	5 5 10		12 B	
21 22 23	97 88 81 79	1 P 2 P 4 P 5 P	71 66 59 55	5 A 12 P 12 P 3 A	84.0 77.0 70.0 67.0	78.0 82.7 53.7 48.7	5.8 8.7 11.6 11.9	W NE W	69 170 161	32 10 22 18	11 P 1 P 3 A	12 P 4 P 7 A	0.68 0.55
24 25 26 27 28	84 95 87 88	2 P 3 P 6 P 3 P	63 65 56	5 A 5 A 12 P 4 A	66.5 79.0 76.0 72.0	56.7 71.0 60.7 63.3	11.1 9.1 11.4 11.7	SE SE NW SE	71 148 94 93	15 15 12 13	4 P 2 A 8 P	5 P 4 A 12 P	0.22 0.07 0.58
29 30	88 88	4 P 3 P	65 57	12 P 5 A	76.5 72.5	69.7 63.3	11.4	SE W	67 53	12	0 A	4 A	0.20

^{*} Based on least time required to blow one mile.

HANS JOA, Observer

MONTHLI	JOININ	7 11 (1
JUNE, 1953	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.191, 28th Minimum 29.654, 9th Mean semi-daily 29.957 Range 0.537	30.30 29.55 29.95 0.75	30.70, 1947 29.24, 1902
Air Temperature, in degrees F.		
Highest 98, 20th Lowest 41, 3rd Mean 69.1 Range 57 Highest mean daily 84.0, 21st Lowest mean daily 56.5, 3rd Mean maximum 83.2 Mean minimum 54.9 Greatest daily range 40, 15th Least daily range 21, 13th Degree-days of heating load for month. 44.5 Degree-days cumulative for season 5886.0	91.2 40.0 65.7 51.3	101, 1919 34, 1891 71.9, 1949; 49.5, 1917
Precipitation, in inches		
Precipitation	3.75	9.68, 1922; 0.72, 1949
Snow Maximum precip. in 24 hrs. 0.68 Number of days with .01 or more. 0	11	17, 1922, 1945: 4, 1908
Wind, in miles		
Total movement2939Greatest daily movement281, 10thLeast daily movement27, 4thMean hourly velocity4.1Maximum velocity36, 9th	3585 5.0 24.6	4571, 1908; 1409, 1906 48, 1939
Wind, direction		
Prevailing direction WNW North, days — Northeast, days .4 East, days — Southeast, days .8 South, days .1 Southwest, days .2 West, days .6 Northwest, days .9	WSW	
Weather		
Mean relative humidity, percent	54.0	71, 1903; 28, 1908 22, 1908, 1941 1943; 0, 1949 23, 1912; 3, 1941 22, 1903; 1, 1923 365, 1947, 102, 1903

Note—The first column in the above summary lists observations made during the month. The second column lists the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column lists extremes observed from 1889 to 1952.

The mean daily temperature for the month of June was 69.1 degrees. This is 3.4 degrees above normal. The precipitation for the month was 2.41 inches compared to the normal of 3.75 inches. During the period May 28 to June 21 there was only 0.11 inch of rainfall, thereby creating mild drought conditions.

Dr. Frank Southwick of the Pomology Department makes the following remarks:

"The month of June was below normal in rainfall but above average in sunlight and average mean temperature. Dry weather noticeably influenced the size and yield of unirrigated strawberries. It does not appear that dry weather has influenced growth of tree fruits yet. Now that the June drop is over, it is apparent that most apple varieties which bloomed, with the possible exception of Delicious, have set enough fruit for a full crop. The above normal amount of sunlight is expected to favor flower bud formation for the 1954 apple crop and help return many alternating McIntosh blocks to an annual bearing condition. As one might expect, following the excessively wet months of March, April, and May, apple scab is a serious problem in many orchards. Peaches have set very heavily, and an above average crop of this fruit is anticipated."

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No 775

July 1953

Meteorological Observations

FOR

JULY

1953

I. J. PFLUG

OBSERVATORY

Latitude, 42° 23′ 31″ N.

Longitude, 72° 31′ 48″ W.

Height of barometer above ground, 36 ft. Above sea level, 253.5 ft.

Height of wind instruments, 67 ft.

Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

-		Te	mpera	ature		Ly.	S		Wind		F	Precipita	tion	
	Max	imum	Min	imum		fumidi	f Hour nshine			\$				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4	86 89 82 82	4 P 4 P 2 P 4 P	66 70 60 51	5 A 4 A 11 P 5 A	76.0 79.5 71.0 66.5	67.7 78.0 51.3 54.3	9.8 10.1 11.5 12.0	W NW NW	71 93 239 59	10 20 32 7	3 A	4 A	0.01	
5 6 7	86 82 85	4 P 2 P 2 P	51 61 62	4 A 4 A 12 P	68.5 71.5 73.5	58.7 70.7 54.0	11.7 8.7 11.7	SW SE W	58 145 150	8 13 17	6 A	7 A	T 0.01	
8 9 10 11	84 74 78 81	1 P M 2 P 3 P	54 59 49 50	5 A 5 A 5 A 5 A	69.0 66.5 63.5 65.5	71.0 60.0 61.0 63.7	8.0 10.8 11.1 11.2	SW W W	158 92 33	22 22 14 12	2 P	12 P	0.84	
12 13 14 15	82 72 76 88	3 P 6 P 3 P	55 58 54 58	5 A 11 A 4 A	68.5 65.0 65.0 75.0	67.3 81.3 79.3 79.3	10.2 6.3 7.9 8.4	W NW NW	43 141 50	7 18 5 8	6 A	М	0.45	
16 17 18	92 96 97	4 P 2 P 4 P 3 P	61 62 67	4 A 5 A 5 A 4 A	76.5 79.0 82.0	68.0 65.7 60.0	10.5 11.5 11.3	NW NW W NW	44 75 46 47	17 13 10				
19 20 21 22	90 89 89 80	4 P 4 P 1 P 2 P	65 70 69 68	5 A 6 A 5 A 5 A	77.5 79.5 79.0 74.0	67.3 82.3 75.0 75.7	9.9 7.4 8.1 9.2	SE SE NE	56 63 49 97	10 7 10 10	4 P 2 P	5 P 6 P	0.03 0.07	
23 24 25	70 78 76	1 P 1 P 5 P	65 56 4 7	12 P 12 P 5 A	67.5 67.0 61.5	87.0 62.3 60.0	0.0 11.0 11.6	NE NW NW	47 165 60	9 18 7	10 A	5 P	0.52	
26 27 28 29	84 88 86 77	2 P 4 P 3 P 6 P	54 65 59 55	4 A 4 A 5 A 4 A	69.0 76.5 72.5 66.0	64.3 76.7 64.0 75.7	10.8 6.9 10.8 3.4	S SE SW NW	109 96 72 35	9 12 17 8	2 Р	3 P	0.02	
30	91 84	3 P 3 P	70 59	4 A 5 P	80.5 71.5	57.3 49.3	10.7 10.9	W NW	105 187	11 25				

^{*} Based on least time required to blow one mile.

HANS JOA Observer

SUMIN	IAINI
Normal	Extremes
30.27 29.59 29.96	30.50, 1892 29.27, 1932
0.68	0.97, 1892; 0.47, 1938
93.9 46.4 70.8 47.7	104, 1911 40, 1890, 1898 66.3, 1891; 76.0, 1949
4.10	14.51, 1897; 0.70. 1929
11	20, 1915; 4, 1924
3422 4.6 28.7	5097, 1909; 1109, 1894 44, 1930
SW	
68.4 50.9 10 14 7 268	70, 1943; 31, 1924 22, 1923; 0, 1915 24, 1909, 1921; 4, 1946 18, 1889; 0, 1910 371, 1910; 180, 1931
	30.27 29.59 29.96 0.68 93.9 46.4 70.8 47.7 4.10 11 3422 4.6 28.7 SW

Note—The first column in the above summary lists observations made during the month. The second column lists the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column lists extremes observed from 1889 to 1952.

The small amount of rainfall, 1.95 inches, compared to the normal of 4.10 inches, was the outstanding feature of the July weather. The mean daily temperature and the mean relative humidity were both very near normal.

The temperature was in the "90's" for four consecutive days, July 16. 17. 18. and 19. According to Professor Kightlinger of the Department of Agronomy this hot spell caused considerable damage to tobacco.

Publication of This Document Approved by George J. Cronin, State Purchasing Agent

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 776

August 1953

Meteorological Observations

FOR

AUGUST

1953

I. J. PFLUG

OBSERVATORY

Latitude, 42° 23′ 31″ N.

Height of barometer above ground, 36 ft.

Height of wind instruments, 67 ft.

Longitude, 72° 31′ 48″ W.

Above sea level, 253.5 ft.

Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Te	mpera	iture		Ly.	g a		Wind		I	recipita	tion	
	Ma	ximum	Mir	imum		Humidi	f Hour nsbine		1	SS.				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3	79 71	3 P 4 P 5 P	48 54	5 A 12 P 2 A	63.5 62.5 63.5	47.3 75.7	11.2 4.4 7.7	W	107 28 69	17 4 9	7 A	9 A	0.03	
4 5	78 77 78	5 P M 10 A	49 52 57	2 A 4 A 12 P	64.5 67.5	56.3 66.0 75.3	11.7	NW SE SE	104 188	13 20			Т	
7	80 81	4 P	52 52	6 A 6 A	66.0	60.3 64.7	11.1	N E	74 63	10 11				
8 -	84 77	4 P 2 P	52 66	3 A 4 A	68.0 71.5	62.7 83.3	108. 3.4	E SE	43 58	7 14	10 р	12 р	0.83	
10 11 12	84 79 84	4 P 3 P 5 P	67 62 55	6 A 12 P 4 A	75.5 70.5 69.5	85.0 79.0 77.7	7.9 9.9 9.4	W	106 103 33	12 17 4	0 A	5 P	1.01 T	
12 13 14	87 79	5 P 3 P 1 P	65 70	3 A 6 A	76.0 74.5	76.3 79.0	8.2 7.2	NW S S	95 52	11 6				
15 16	84 80	3 P 2 P	65 55	12 P 5 A	74.5 67.5	59.7 58.7	10.6 11.0	W W	170 106	25 13				
17 18 19	77 76 77	3 P	57 47 45	6 A 6 A 5 A	67.0 61.5 61.0	72.7 67.0 63.3	8.6 10.5 9.8	W	73 60 79	11 15				
20 21	76 77	3 P 4 P 3 P	49 50	5 A 6 A 6 A	62.5 63.5	59.0 64.7	10.8	NW NW	50 37	11 7				
22 23	80 84	3 P 3 P	50 54	7 A 3 A	65.0 69.0	68.3 69.0	9.8 8.2	NE N	20 45	6 5				
24 25	87 88	2 P 4 P	55 58	4 A 5 A	71.0	71.7 73.0 69.7	9.5 9.0 7.2	N NW	55 58 70	11 10 11			T	
26 27 28	86 90 94	2 P 3 P 3 P	67 60 66	6 A 6 A 4 A	76.5 75.0 80.0	70.0 66.7	9.7 8.6	SE W S	15 82	6			т	
29 30	96 96	3 P 2 P	62 61	6 A 6 A	79.0 78.5	61.0 60.7	10.0 8.7	NW NW	68 31	12 10				
31	95	3 P	65	4 A	80.0	52.3	9.8	s	133	18				

^{*} Based on least time required to blow one mile.

HANS JOA, Observer

PIONTIE		/ 11 C I
AUGUST, 1953	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.300, 12th Minimum 29.558, 5th Mean semi-daily 30.078 Range 0.742	30.32 29.61 30.00 0.71	30,50, 1934 28,87, 1930
Air Temperature, in degrees F.		
Highest 96, 29th, 30th Lowest 45, 19th Mean 69,8 Range 51,0 Highest mean daily 80,0, 28th, 31st Lowest mean daily 61,0, 19th Mean maximum 82,6 Mean minimum 57,0 Greatest daily range 35,0, 30th Least daily range 9,0, 14th	91.6 43.4 68.6 48.1	100, 1918, 1948 34, 1940 62.4, 1903;73.5, 1939
Precipitation, in inches		
Precipitation 1.87 Snow — Maximum precip. in 24 hrs. 1.01, 10th	4.08	8.40, 1948; 0.31, 1894
Maximum precip. in 24 hrs 1.01, 10th Number of days with .01 or more3	11	16, 1892, 1933; 4, 1899
Wind, in miles		
Total movement. 2275 Greatest daily movement 188, 5th Least daily movement 15, 27th Mean hourly velocity 3.1	3127	4.217, 1910; 1,920, 1894 40, 1941
Maximum velocity	22.7	30, 1241
Wind, direction Prevailing direction WNW North, days 3 Northeast, days 1 East, days 3 Southeast, days 4 South, days 4 Southwest, days - West, days 9 Northwest, days 7	SW	
Weather		
Mean relative humidity, percent. 67,3 Mean cloudiness, percent 40.5 Number of clear days 16 Number of fair days 14 Number of cloudy days 15 Number of hours bright sunshine 278.3 Percent of possible hours of bright sunshine 64.6 Thunder and lightning	70.2 49.7 9 13 9 237	67, 1901, 1946; 27, 1923 23, 1941; 0, 1915, 1929 25, 1912; 3, 1939 18, 1901, 1928; 2, 1910, 1923 332, 1941;152, 1915, 1929

Note — The first column in the above summary lists observations made during the month. The second column lists the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column lists extremes observed from 1889 to 1952.

The weather for the month of August was unusual. The early part of the month was cool with the last week in the month reaching record highs. The mean daily temperature for the month was 1.2 degrees above normal. Although the record high temperature for August was not broken, record high temperatures were recorded for August 29, 30 and 31.

The rainfall for the month was 1.87 inches compared to the normal of 4.08 inches.

Dr. Ives of South Amherst reported that August was the tenth consecutive month with above normal temperatures. He also observed that this was the fifth driest summer (least rain) since 1929.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 777

September 1953

Meteorological Observations

FOR

SEPTEMBER

1953

I. J. PFLUG

OBSERVATORY

Latitude, 42° 23′ 31″ N.

Height of barometer above ground, 36 ft.

Height of wind instruments, 67 ft.

Longitude, 72° 31′ 48″ W.

Above sea level, 253.5 ft.

Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Te	mpera	ture		k:		Wind Precipitation						
	Max	imum	Min	imum		Humidi	of Hour Inshine	200	ıt	s.*				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3	95 99 96	3 P 3 P 2 P 3 P	57 68 68	6 A 5 A 6 A	76.0 83.5 82.0	53.7 57.0 62.3	10.1 9.4 9.3	NW S NW	36 109 42	7 20 10				
4 5	92 90	3 P 1 P	67 73	3 A 5 P	79.5 81.5	76.3 73.3	6.7 7.9	SE SE	85 213	13 20			Т	
6 · 7	78 80	3 P M	65	12 P 12 P	71.5 68.0	75.3 78.3	5.3 5.0	NW NW	115 157	11 18			T	
8	76 73	2 P 2 P 3 P	50 40	5 A 5 A	63.0 56.5	56.0 61.7	9.3 9.7	W	129 94	20				
10 11	76 81	3 P 3 P	38 43	6 A 6 A	57.0 62.0	64.0 64.0	10.0	W	47 49	9 8				
12	74 73	М 1 Р	49 53	6 A 12 P	61.5	78.7 68.3	4.9 8.1	SE SW	118 193	13 20	10 p 0 a	12 P 3 A	0.08	
14 15	65 66	3 P 4 P	50 44	5 A 6 A	57.5 55.0	55.7	10.0	W	216	22	0	7.4	1.12	
16 17	77 70	3 P 4 P	47 47	5 A	62.0	75.0	7.6	s	66	13	8 P	10 р	0.12	
18	70	4 P	44	6 A 4 A	57.0	70.0	9.4 9.6	NW NW	83	10				
19 20	71 76	3 P 11 A	44 59	7 A 0 A	57.5 67.5	75.3 70.7	8.7	SE SE	128 141	13				
21	77	4 P	56	12 P 12 P	54.5	77.7	6.3	SE W	113	20	3 A 9 P	7 A 12 P	0.26	
22 23	61 68	2 P 3 P	34	6 A	47.5	67.0	9.4	W	61 69	14				
24 25	70	4 P 1 P	30	6 A	49.0 55.5	69.0	8.8 7.3	NW SE	94	11				
26 27	78 78	3 P M	57 60	0 A 3 A	67.5 69.0	72.3 81.0	7.9 6.8	SE SE	138 80	10				
28 29	73 75	3 P 2 P	50 43	12 P 5 A	61.5	63.0 68.3	9.0	NW NW	85 63	13				
30	69	4 P	144	5 A	56.5	80.7	7.5	SE	118	22				

^{*} Based on least time required to blow one mile.

Hans Joa, Observer

SEPTEMBER 1953	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.567, 24th Minimum 29.623, 13th	30.45 29.57	30.65, 1924 28.41, 1938
Mean scmi-daily	30.00	1.99, 1938; 57, 1910
Air Temperature, in degrees F.		
Highest 99. 2nd Lowest 30. 24th Mean 63.5 Range 49 Highest mean daily 83.5, 2nd Lowest mean daily 47.5, 23rd Mean maximum 76.4 Mean minimum 50.6 Greatest daily range 38, 1, 10, 11, 24 Least daily range 13, 6th Degree-days of heating load for month 141 Degree-days cumulative for season 141	87.7 33.2 61.7 54.5	97,1895, 1929 24.5, 1914
Precipitation, in inches		
Precipitation 1.42 Snow —	4.24	14.55, 1938; 0.52, 1914
Maximum precip. in 24 hrs	10	16, 1933; 3, 1903
Wind, in miles		
Total movement. 3100 Greatest daily movement 216, 14th Least daily movement 23, 15th Mean hourly velocity 4.3 Maximum velocity 22, 30th	4.5	4,680, 1896; 1414, 189- 80, 1938
	20.7	00, 120
Wind, direction	MACAN	
Prevailing direction. WNW North, days — Northeast, days — East, days — Southeast, days 10 South, days 2 Southwest, days 2 Northwest, days 7 Northwest, days 9		
Weather		
Mean relative humidity, percent. 69.3 Mean cloudiness, percent 38.2 Number of clear days 15 Number of fair days 15 Number of cloudy days Number of hours bright sunshine 239.6 Percent of possible hours of bright sunshine 63.8	49.9 10 10 10 201	70,1934; 23, 194 19, 1932; 2, 1907, 192 19, 1908; 3, 188 20, 1934; 3, 1941, 194 255, 1916; 106, 193
Thunder and lightning		Oct. 13, 1909; Aug. 22, 1894-9

Note—The first column in the above summary lists observations made during the month. The second column lists the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column lists extremes observed from 1889 to 1952.

The outstanding feature of the September weather was the low rainfall. The precipitation was 1.88 inches compared to the normal of 4.24 inches.

A new record temperature for the month was established September 2 when the mercury reached 99 degrees.

Publication of This Document Approved by George J. Cronin, State Purchasing Agent.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 778

October 1953

Meteorological Observations

FOR

OCTOBER

1953

I. J. PFLUG

OBSERVATORY

Latitude, 42° 23′ 31″ N.

Longitude, 72° 31′ 48″ W.

Height of barometer above ground, 36 ft. Above sea level, 253.5 ft.

Height of wind instruments, 67 ft.

Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Tei	mpera	ture		ń.	m	Wind Precipitat					tion	
	Maximum		Min	Minimum		Iumidi	f Hour nsbine			20				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunsbine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 20 21 22 23 24 25 27 28 29 30 30 31 31 31 31 31 31 31 31 31 31 31 31 31	75 76 81 80 63 52 52 57 66 68 63 61 61 74 74 77 77 58 62 63 73 57 62 63 63 67 67 67 67 67 67 67 67 67 67 67 67 67	2 P 4 P 3 P 1 P 0 A 2 P 1 P 2 P 2 P 4 P 4 P 4 P 4 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2	48 40 48 48 48 44 38 33 37 42 29 25 32 37 33 36 40 40 40 40 40 40 40 40 40 40 40 40 40	12 P 5 A 2 A 12 P 12 P 12 P 12 P 6 A 6 A 4 A 12 P 5 A 6 A 7 A 7 A 7 A 12 P 6 A 6 A 12 P 12 P 5 A 6 A 7 A 7 A 7 A 12 P 6 A 12 P 12 P 12 P 12 P 12 P 12 P 13 A 14 A 15 A 16 A 17 A 18 A	61.5 57.0 60.5 64.0 55.5 48.0 45.0 45.0 50.5 51.5 45.0 46.0 53.0 56.5 54.5 55.0 64.0 57.5 51.5 51.5 55.0 64.0 57.5 51.5 51.5 55.0 64.0 57.5 57.6 67.5 57.5 57.6 67.5 57.6 67.5 57.6 67.5 57.6 67.5 57.6 67.5 57.6 67.5 57.6 67.5 57.6 67.5 57.6 67.5 57.6 67.5 57.6 67.6	56.3 66.7 64.7 67.3 85.0 62.0 63.7 77.7 68.3 77.7 68.3 62.7 72.0 66.3 67.7 64.3 69.7 70.0 77.7 58.3 60.0 87.7 80.0 93.0 91.0 95.0 77.3 73.3	9.0 8.9 8.7 8.4 0.0 6.1 9.0 6.1 9.0 8.4 4.2 4.9 8.7 8.4 7.5 8.0 8.1 7.9 6.2 0.0 6.2 0.0 6.2 0.0 6.2 0.0 7.1 7.2 0.0 7.1 7.2 0.0 0.0 7.2 0.0 0.0 7.2 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0	NW NW NW S N NW NW NW NW NW NW NW NW NW NW NW NW N	102 55 50 137 95 92 205 121 45 57 62 122 32 44 49 30 25 29 73 62 76 117 129 62 92 33 80 209 52	11 28 15 7 7 8 20 4 8 8 8 10 12 10 9 9 9 11 10 13 14 4 5 5 10 10 10 10 10 10 10 10 10 10 10 10 10	2 P 0 A 3 P 1 A 3 A 0 A	11 P 12 P 3 A	0.46 0.77 0.16 T T 1.02 0.15 0.71 1.26 0.48	

^{*} Based on least time required to blow one mile

OCTOBER 1953	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30.462 Minimum 29.522 Mean scmi-daily 30.061	30.53 29.42 30.06	30.68, 1929 29.00, 1926
Range	1.11	1.47, 1926; 0.76, 1899
Air Temperature, in degrees F.		
Highest 81, 3rd Lowest 25, 14th Mean 53,6 Range 56 Highest mean daily 64, 4th, 20th Lowest mean daily 45, 7th, 8th, 13th Mean maximum 66,5 Mean minimum 40,7 Greatest daily range 42, 14th, 15th Least daily range 8, 6th, 29th Degree-days of heating load for month 353 Degree-days cumulative for season 494	79.4 23.4 50.5 56.0	90.5, 1908 17, 1936 57.6, 1949; 43.2, 1890
Precipitation, in inches		
Precipitation	3.29	8.81. 1911; 0.01, 1924
Snow — Maximum precip, in 24 hrs. 1.26 Number of days with .01 or more9	9	15, 1913; 1, 1897, 1924
Wind, in miles		
Total movement	5.4	5467, 1910; 2540, 1894
Mean hourly velocity 3.2 Maximum velocity	29.5	42, 1937
Wind, direction		
Prevailing direction NNW North, days .7 Northeast, days .3 East, days Southeast, days South, days Southwest, days West, days Northwest, days 15	W	
Weather		
Mean relative humidity, percent. 72.6 Mean cloudiness, percent	69.0 48 10 10 11 175.0 51.3	67, 1932; 18, 192- 21, 1938; 1, 191 17, 1924; 3, 193 19, 1896; 3, 192- 232, 1923, 1938; 91, 191

Note—The first column in the above summary lists observations made during the month. The second column lists the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column lists extremes observed from 1889 to 1952.

The outstanding feature of the October weather was the rainfall of 5.15 inches compared to the normal of 3.29 inches. This was the first month since May that the rainfall was above normal. Rainfall for the first 10 months of 1953 was 42.41 inches compared to the normal of 36.91 inches.

The mean daily temperature for October was 3.1 degrees above normal.

PUBLICATION OF THIS DOCUMENT APPROVED BY GEORGE J CRONIN, STATE PURCHASING AGENT.

MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No. 779

November 1953

Meteorological Observations

FOR

NOVEMBER

1953

I. J. PFLUG

OBSERVATORY

Latitude, 42° 23′ 31″ N. Height of barometer above ground, 36 ft. Above sea level, 253.5 ft.

Longitude, 72° 31′ 48″ W.

Height of wind instruments, 67 ft.

Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

		Те	empera	ature		ty	s.	Wind Precipita					ation		
	Maximum		mum Minimum		num		f Hour nshine		_	1 2					
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow	
1 2 3 3 4 4 5 6 7 7 8 9 100 111 122 133 144 15 16 17 18 19 20 21 22 23 24 25 27 28 29 30	64 65 72 72 41 35 49 49 53 52 49 40 57 57 63 70 64 66 55 56 48 45 41 43 66 45 45 45 45 45 45 45 45 45 45 45 45 45	3 P 2 P 2 P M 1 P 9 P 2 P 2 P 2 P 3 P 3 P 3 P 10 P 12 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P 2 P	35 39 42 34 26 30 36 32 29 31 31 34 36 31 31 34 36 31 31 34 36 31 31 31 31 31 31 31 31 31 31 31 31 31	6 A A 6 A A 12 P 12 P 6 A A 1 A 3 A A 7 A A 8 A A 12 P 8 A A 12 P 7 A 8 A A 12 P 7 A	49.5 52.0 57.0 57.0 39.5 42.5 39.5 45.0 42.5 41.5 37.5 45.0 44.0 52.0 47.0 52.0 47.0 52.0 45.5 57.5 57.5 57.5 57.5 57.5 57.5 57.5	79.0 74.0 66.3 54.7 75.3 66.0 77.7 75.3 81.7 87.7 85.0 74.0 75.0 74.0 74.0 75.0 74.0 75.0 74.0 78.0 74.0 78.0 78.0 74.0 78.0 78.0 78.0 78.0 78.0 78.0 78.0 78	6.9 6.5 5.1 6.7 6.9 0.0 0.6 6.8 3.3 6.3 5.8 3.6 6.1 0.0 2.6 6.0 3.8 3.6 5.9 5.5 5.8 3.6 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0	NW NW S W NW NW N N N N N N N N N N N N	31 71 97 2268 147 2100 353 113 1101 9 17 11 12 28 9 29 34 12 12 17 162 17 53 108 14 32 108 166 166 166	10 20 25 32 20 48 25 14 13 3 5 10 — 5.5 4 43 25 9 13 17 2.5 7 7 20 8	2 P 6 A 11 P 0 A 5 A 8 P	6 P 1 P 12 P M 6 P	0.16 0.10 1.08 0.91	1	

^{*} Based on least time required to blow one mile.

MONTHLY SUMMARY

).87, 1932 3.73, 1904
0.89. 1943
82, 1956 4, 1933 33,6, 190
0.63. 1917 3.50, 1938
1; 2, 1904
589, 1889 948, 1950
748, 1730
; 34, 1917 3; 0, 1948 0, '32, '35 917, 1949 ; 44, 1948

Note—The first column in the above summary lists observations made during the month. The second column lists the averages based on observations made from 1889 to 1938 except that humidity records are based on observations made from 1929 to 1938. The third column lists extremes observed from 1889 to 1952.

REMARKS

The mean daily temperature for November was 43.2 degrees compared to the normal of 38.9 degrees Fahrenheit. The rainfall for November was 2.36 inches compared to the normal of 3.41 inches.

The first snowfall of the season occurred on November 7.

Publication of This Document Approved by George J. Cronin, State Purchasing Agent

INVERSITY OF MARSON ANDSELLIS

MASSACHUSETTS

AGRICULTURAL EXPERIMENT STATION

Meteorological Series

Bulletin No 780

December 1953

Meteorological Observations

FOR

DECEMBER

1953

1. J. PFLUG

OBSERVATORY

Latitude, 42° 23′ 31″ N.

Longitude, 72° 31′ 48″ W.

Height of barometer above ground, 36 ft. Above sea level, 253.5 ft.

Height of wind instruments, 67 ft.

Time used, E. S. T.

Requests for bulletins should be addressed to the AGRICULTURAL EXPERIMENT STATION AMHERST, MASS.

DAILY RECORDS

	Temperature				,	· va	Wind			Precipitation				
	Maximum		Min	nimum		Humidi	of Hour Inshine	bs.	ä	*es				
Day	Degrees	Time	Degrees	Time	Mean	Mean Relative Humidity	Number of Hours Bright Sunshine	Prevailing Direction	Total Movement	Max. Miles per Hour*	Began	Ended	Water	Snow
1	37 44	2 P 1 P	31	0 A 12 р	34.0 34.5	91.7 66.3	0.1 5.9	N NW	120 198	12.5 12	0 A	11 A	0.09	1
3	44 50	3 P	21 25	4 A 4 A	32.5 40.5	77.0 85.3	3.6 0.0	NW NW	14 34	18 18			T	
1 5	56	1 A	34	12 P	45.0	76.7	5.2	W	216	28	0 A	4 A	0.71	
6	45 59	12 P 1 A	31 35	7 A 12 P	38.0 47.0	92.0 70.0	0.0	M.	24 226	32	7 p 0 a	12 P 4 A	0.55	
7 8	59 59	2 P	34	7 A	46.5	56.7	4.2 5.5	S	130	12.5	•			
9	54	12 P	30	6 A	42.0	87.7	0.0	SE	50	11	7 P	12 P	0.20	
10	58	10 A 3 P	38 31	12 P 8 A	48.0 39.5	74.0 68.0	5.7	W	277 190	32 22	() A	8 A	0.69	
11 12	48 48	2 P	38	2 A	43.0	77.7	0.0	SE	135	8	7 P	11 Р	0.18	
13	49	2 P	34	9 P	41.5	78.7	5.5	W	35	9		_	1.00	
14	46	11 Р	32	0 A	39.0 37.5	61.0	0.0	N W	84 177	22 39	4 A 5 A	7 P 7 A	1.08	T ()
15 16	45 34	M = 0.A	30	12 P 12 1	27.0	63.0	5.4	W	166	15	JA	/ 1	0.00	()
17	22	M	11	12 P	16.5	67.0	5.4	W	133	18				
18	28	2 P	8	6 A	18.0	76.0	4.9	W.	87	22				
10	29 45	8 P	26	7 A 0 A	19.0 35.5	71.7	2.0	S	105 108	8 12				
20 21	57	2 P 2 P	35	2 A	46.0	64.3	5.1	SE	103	11				
22	52	2 P	37	6 A	44.5	86.0	3.3	SE	68	13	2 A	5 A	0.26	
22 23	48	L 0	19	12 P	33.5	57.3	1.7	W	295 112	22 14				
24 25	28 44	3 P 1 P	15 21	5 A 2 A	21.5	60.0 54.3	4.3	NW SE	138	10.5				
25 26	42	M	24	6 A	33.0	73.0	4.2	NW	104	12.5				
27	36	2 P	22	12 P	29.0	62.7	6.0	w	201	22	2 A	3 A	0.06	
28	39	M	22	0 A	30.5	84.3 98.3	0.0	W	75 49	12.5 5	1 P	4 P	0.04	
29 30	37 38	2 P 3 P	26 24	12 P 3 A	31.5 31.0	98.3	0.0	SE SE	10	4				
31	30	2 P	20	12 P	25.0	62.7	4.9	W	161	28				

^{*} Based on least time required to blow one mile.

MONTHLY SUMMARY

DECEMBED 1051	1	
DECEMBER 1953	Normal	Extremes
Barometer in inches (Readings reduced to sea level)		
Maximum 39,301, 24th Minimum 29,221, 14th Mean semi-daily 29,973	30.66 29.27 30.06	31.104, 1949 28.85, 1915
Range	1.38	1.78, 1895; 1.01, 1892
Air Temperature, in degrees F.		
Highest 59, 7th, 8th Lowest 8, 18th Mean 34,9 Range 51 Highest mean daily 48, 10th Lowest mean daily 16,5, 17th Mean maximum 43,8 Mean minimum 26,0 Greatest daily range 32,5th Least daily range 6, 1st	54.3 -1.9 27.5 56.2	65.5, 1908 —22.5, 1917 36.9, 1891; 17.1, 1917
Degree-days of heating load for month 932.5 Degree-days cumulative for season 2080.5	1162.5 2494.0	
Precipitation, in inches		
Precipitation 4.53 Snow 1.00 Maximum precip. in 24 hrs. 1.08	3.39 8.50	7.77, 1901; .58, 1943 26.50, 1092; T, 1891, 1943
Number of days with .01 or more 12	10	17, 1902; 4, 1892, 1943
Wind, in miles		
Total movement	4710 6.3	6694. 1925; 3239, 1918
Maximum velocity	31.4	66, 1949
Wind, direction		
Prevailing direction WNW North, days .2 Northeast, days — East, days .7 Southeast, days .3 Southwest, days .4 West, days .14 Northwest, days .5	WNW	
Weather		
Mean relative humidity, percent. 74.8 Mean cloudiness, percent 56.9 Number of clear days3 Number of fair days16 Number of cloudy days12 Number of hours bright sunshine 94.6 Percent of possible hours of bright sunshine33.5 Thunder and lightning First Snow	69.4 54.9 8 9 14 128 45.2 Nov. 6	71, 1929; 39, 1919 15, 1890; 0, 1948 16, '09; 4, '89, '30, '31, '36, '38 25, 1948; 7, '09, '23 172, 1896 27, 1948

Note—The first column in the above summary lists observations made during the month. The second column lists the averages based on observations made from 1889 to 1938, except that humidity records are based on observations made from 1929 to 1938. The third column lists extremes observed from 1889 to 1952.

ANNUAL, SUMMARY

ANNUAL	Normal	Extremes
Barometer, in inches (Readings reduced to sea level)		
Maximum 30,690, March 11th Minimum 29,039 Jan, 25th Mean semi-daily 29,988 Range 1,651	30.81 28.95 30.01 1.85	31.10, 1949 28.41, 1938 2.47, 1938; 1.38, 1933
Air Temperature, in degrees F.		
Highest 99, Sept, 2nd Lowest 7, Feb. 2nd Mean 51.1 Range 92 Highest mean daily 84.0, June 21st	95.7 -12.2 47.4 107.8	104, 1911 —26, 1904 51.7, 1949; 44, 1904
Lowest mean daily 15.5. Feb. 2nd Mean maximum 1.7 Mean minimum 40.5 Greatest daily range 42. Oct. 14th. 15th Least daily range 3. May 15th	57.8 36.5	
Degree-days 1952-53 heating season 5886	6698	
Precipitation, in inches		
Precipitation 19.76 Snow 27.3 Maximum precip, in 24 hrs.1.77 March 13th	43.70 47.78	59.00, 1938; 30.68, 1908 89.00, 1893; 24.50, 1919
Number of days with .01 or more12?	124	146, 1945; 96, 1924
Wind, in miles		
Total movement	52,223	63,571, 1908; 36,257, 1894 705, 1909
Mean hourly velocity 4.4 Maximum velocity 48. Feb. 15th 16th	5.8 39.5	80, 1938
Wind, direction		
Prevailing direction WNW North, days 39 Northeast, days 17 East, days 52 Southeast days 39 Southwest, days 17 West, days 73 Northwest, days 117	\\	
Weather		
Mean relative humidity, percent 71.7 Mean cloudiness, percent 51.0 Number of clear days 152 Number of fair days 136 Number of cloudy days 77 Number of hours bright sunshine 2373.1 Percent of possible hours of	69.9 51.7 116 123 126 2,353	77.9, '51; 64.9, '32 60, '98, '01, '02; 41, '08, '24 217, 1941; 59, 1927 182 1912; 64, 1936 179, '01, '02; 71, 1910, 1941 3038, 1941; 1864, 1902
bright sunshine 51.7 Last snow April 20th First snow Nov. 7th Last frost April 14th First frost Sept. 24th	52.8 April 15 Nov. 6 May 14 Sept. 21	Mar. 14 '10; May 11, '07, '45 Oct. 10, '25; Dec. 13, '41 Apr. 14, '43, June 8, '32 Aug. 22, '94, '95; Oct. 13, '09

